TABLE IV—Comparison of consultations in one calendar month for 10 pairs of trainees and trainers

	Mean No. of consultations in one month	
	Trainee	Trainer
Asthma	15.5	5.6
Depression	12.4	4.4
Diabetes	2.3	0.6
Epilepsy	2.4	0.2
Hypertension	21.4	1.1
Hypercholesterolaemia	20.5	0.2
Phlegm	2.0	1.1
Rheumatoid arthritis	4.2	0.4
Surgeons	112.6	18.7
Home	11.2	11.2

the practice mean list size of each principal, the mean number of trainee surgery and home visits a week, and the number of partners. There were no correlations. The percentages of patients seen twice or more for the five conditions after recruitment were plotted against the same variables for all full time trainees on 11 or 12 month attachments. Once again there were no correlations except a suggestion that the chances of a patient being seen twice or more after recruitment by the trainee was in inverse proportion to the number of partners. The mean percentage of patients seen twice or more was 15.7% for 10 trainees in practices with more than four partners and 31.8% for 24 trainees in practices with four partners or fewer.

Prescribing

As part of the checking procedure on the trainees' efficiency, 22 501 prescriptions (FP10) were examined. Prescriptions signed by the trainee but written by someone else were excluded. Large discrepancies were found between the mean rate per month for the trainees for such drugs as hypertensives, diuretics, antidepressants, bronchodilators, and anti-inflammatory drugs and the rates for unrestricted principals: table V shows some of these figures.

TABLE V—Mean prescribing rates per month for Oxford region trainees 1976-9 and unrestricted principals in England in 1978

Therapeutic class of drug	Mean rate per month	
	Trainees	Unrestricted principals
Preparations for treatment of asthma	5.4	44.6
Antidepressants	1.4	1.7
Anticonvulsants	0.6	12.7
Hypocarbamic agents including insulin	1.4	1.7
Diuretics	1.4	1.7
Anti-inflammatory drugs (excluding steroids)	7.5	19.0
Anticancer drugs	0.1	0.1

Source for unrestricted principals: Department of Health and Social Security.

These comparisons have to be treated with some caution. The figures for unrestricted principals include those from trainees and assistants but the latter are a small proportion only of the total. Of more importance, however, is the fact that the figures for unrestricted principals include all prescriptions dispensed, which include repeat prescribing and not just those written fully in the doctor's handwriting, as was the case for the trainees. Even if these factors are taken into consideration, however, the discrepancies are still large.

Discussion

It may be seen that most patients with these chronic problems that each trainee managed consulted him in the first six months. Table II shows that the mean number of patients new to the trainee was greater than the mean number of patients new to the practice compared with the second five or six month period. This pattern was apparent in virtually every analysis from each trainee for every condition. It seems therefore that the trainees did not have to be established in the practice for several months before these patients came to consult them.

The number of patients consulting per 1000 population did not differ greatly from figures in England and Wales in the second National Morbidity Survey.¹ Only for depression, osteoarthritis, pregnancy, and rheumatoid arthritis were there moderately large discrepancies. The range of numbers of patients seen by the trainees, however, does give rise to some concern. For 11 out of the 16 conditions there were some trainees on 12 month attachments who recorded no cases at all.

Numbers of consultations (as opposed to patients) showed larger discrepancies with national figures. This is confirmed in table IV in the comparisons with trainers. Table III shows the large percentage of patients seen once or twice only by trainees. These figures raise serious questions about the continuity of care by the trainees in the study—an activity thought to be a fundamental part of good general practice. For example, trainees on six month attachments saw 69% of patients with asthma, 72% of patients with diabetes, and 80% of patients with epilepsy only once. These figures might have been expected to some extent. But the same cannot be said of depression and pregnancy, where one would expect patients to be seen much more often. Trainees on six month attachments saw 52% of patients with depression and 62% of pregnant women once only.

Why should this continuity of care be lacking? Many trainees have said that they feel different about formally allocating patients to trainees and some trainees have said they feel different about asking patients to come back and see them again.

Conclusions

It is clear that the study could not relate what the trainees saw to how much they learnt. Nor can one deduce that there was a standard relation between the two or that there is a certain minimum number of cases that need to be seen for any particular condition.

As the main activity of trainees in training practices, however, is seeing patients, and as these patients seem to dominate what is learnt in tutorials, there are several questions that need answers. How far are trainees aware of what kind of patients their trainees are seeing? How do trainees learn effectively about the management of conditions if they have seen no patients with the condition in question? Some kind of clinical log book seems to be needed.

How feasible is it to change the trainees' clinical work if it seems that some areas are deficient? The percentages of patients returning to the trainees bore no relation to the trainees' overall workload, and therefore merely giving trainees more patients is unlikely to make any difference. On the other hand, Fournier and Savory showed that it is possible to steer patients to the trainee and appreciably increase his number of patients with chronic disease, with arrangements that were largely acceptable to the patients.²

What factors influence how trainees learn and teach in tutorials, and can trainers and trainees be helped to exploit individual consultations to achieve the maximum learning? In the Oxford region we have recently completed work on identifying priority objectives for vocational training (unpublished observations) in which we argue that the most important objectives for trainees to achieve in chronic disease are those related to general principles. These principles include those related to patient participation in management and treatment, and arrangements for monitoring clinical features and repeat prescribing. Any patient with chronic problems can provide the means for achieving many of these objectives, and can to some extent make up for the lack of patients with different chronic problems.

It is to be hoped that the results of this study will highlight the need for trainees to know what their trainees are doing and to use patient consultations to the maximum effect in their teaching.

It is a pleasure to thank the trainers and trainees in the Oxford

Despite this, I took the examination somewhat carelessly, relying on age and experience to see me through, so that when I sat the actual papers I was shaken by the amount of theoretical knowledge required. It was no surprise, therefore, when a few weeks later that taciturn letter dropped through the post box: "I know you are disappointed not to have gained enough marks ... etc. etc." Ego blows are not softer as one gets older, but one develops defence mechanisms. "It is for younger doctors"; "It is not a true test of anything if you have not passed"; "How can a good doctor be mediocre?" In truth I do not agree with these statements. I think that as far as it can be a good test and my failure was due to myself and a poor performance, and not to my stars. Therefore, I clutched my *Meditations of Marcus Aurelius*.

Minimum Standards for Training

Do standards improve patient care?

J J C CORMACK

The best government is that which governs least. J O Sullivan, 1837.

If all the major divisions or specialties in medicine, general practice is probably the most diverse in philosophy, tradition, and environment. The rapid development of general practice in the United Kingdom, which has flowed from the family doctor's charter and the establishment of the Royal College of General Practitioners and of university departments of general practice, has converted the cottage industry of the 1960s into the academic discipline of the 1980s. This development, linked with advances in treatment and greater awareness of patterns of human behaviour, has almost certainly improved the quality of primary patient care (difficult as this is to quantify), but has it begun to threaten the diversity, which is one of the glories of general practice, with a uniformity that could stultify further progress? This is a danger that should be considered by all who seek to lay down or suggest minimum standards of practice or of training.

The establishment of vocational training as a statutory prerequisite for independent general practice, with a heavy responsibility on the profession to provide suitable environments and tools to achieve the aims that are set. Would that responsibility not be met by simply appointing trainees to established practitioners and then letting them fend for themselves? Is it far to lay down basic standards for training practices when these standards may not be reproducible in the practices which the trainees will eventually enter?

I think that apprenticeship is not enough and that it is fair to expect certain basic standards of training practices. The danger of imposing a dull uniformity is illusory: variety of personality and circumstances will combat that. Standards are needed to meet the objective of providing "training experiences of wide opportunities and high quality," which is the declared aim of the responsible overseeing body.¹

Ladywell Medical Centre, Cornisthorpe, Edinburgh EH12 7TB

J J C CORMACK, MB, FRGP, general practitioner

a little closer to my heart. "Never say how unlucky I am that this should happen to me, do not feel bitter, say 'this is a misfortune, but to bear this worthily is good fortune'."

I may never become a member of that illustrious college, but that colleague who first put the idea into my head affected me in the same manner as the enigmatic raven affected Poe. And will I ever be free from ego-threatening examinations? Is there harm in Gilead? Quoth the colleague, "Nevermore." And as I am resisting the whole exam, my soul is still in the shadow of a presence, which, like the Raven, "still is sitting, still is sitting ... just above my chamber door."

(Accepted 25 October 1983)

Objectives

As Wright pointed out in a masterly review of what should be taught and learnt the objectives of training are the development of the specific skills and the acquisition of the specific knowledge on which general practice depends, along with the further development of the trainee's critical faculties.² Training is a preparation for specific forensic tasks and should build on and include elements of education. Education may be defined as the development of mental skills to enable the student to respond appropriately to unforeseen situations.

The environment in which such objectives may be satisfactorily achieved must encourage not only elements of apprenticeship ("do as I do") but also education ("but question why I do it as I do"). As Donald said: "If learning is to achieve its full educational value ... it must be supported by teaching."³

Minimum standards (table)

What then should be the minimum standards required of a teaching practice?

Minimum standards of a training practice	
Journal	Commitment
Texts with allied professionals	Commitment
Library	Library
Specialist interest	Specialist interest
Personal notes	Personal notes
Medical records	Medical records
Medical records	Medical records

(1) *Commitment*—It is of primary importance that the teaching unit should be the practice—not only the designated trainee but his partners, paramedical colleagues, and staff must be

region who took part in the study, and Dr C M Harris and Professor M P Vesey who advised on the original work.

This paper is derived from a thesis approved for the degree of MD for the University of London.

References

- Ronalds C, Douglas A, Perera Gray DJ, Selley P, eds. *Fourth national training conference*. London: Royal College of General Practitioners, 1981.
- RCGP Occasional Paper 16.
- Tate PH, Pendleton DA. Why not tear up the European aim? *J R Coll Gen Pract* 1980;30:743.
- Bain DJ. The first freedom. *J R Coll Gen Pract* 1969;19:29-35.
- Gilchrist IC, Mackay J. Personal points of view—the work of a trainee in general practice. *Health Bull (Edin)* 1973;33:721-5.
- Gibson JP. A study of general practitioner consultations and work-load in a training practice in South-West Ireland. *Journal of the Irish Medical Association* 1977;70:174-80.

- O'Hanlon PH. One trainee's clinical experience. *J R Coll Gen Pract* 1977;27:32-30.
- Stratton BA, Gowers JI. A comparison of trainee and trainer clinical experience. *J R Coll Gen Pract* 1979;29:47-52.
- Carver TA. Clinical experience in general practice. *J R Coll Gen Pract* 1979;29:40-4.
- Richardson JM, Morris JGR. A study of trainee general practitioners. *British Journal of Medical Education* 1972;6:26-31.
- Richardson JM, et al. A further study of trainee general practitioners. *J R Coll Gen Pract* 1974;24:501-5.
- Anonymous. Vocational training. [Editorial] *J R Coll Gen Pract* 1979;29:5-5.
- Office of Population Censuses and Surveys. *Studies on medical and population subjects no 28. Morbidity statistics from general practice and local authorities 1979-72*. London: HMSO, 1979.
- Fournier R, Savory JN. Steering patients with selected conditions to trainees. *J R Coll Gen Pract* 1980;30:729-33.

(Accepted 12 October 1983)

MRCGP Examination

Do you think, at your age, it is right?

RAYMOND E GOODMAN

To my shame, no sense of guilt, no fear of being out of date, not any overwhelming passion to update myself was sufficient to cause me to re-read the whole syllabus of medicine after a gap of 20 years. I had dabbled, reading the odd journal now and then, but only articles that interested me and little outside my special interests. The idea of a medical examination after so long awoke me, like Shelley's West Wind did the sleeping Mediterranean, which had been "lulled by the coil of his crystalline streams, Beside a pumice isle in Baia's bay." Since I took a textbook of medicine on holiday with me to that region, the simile is apt, both that sea and I "going grey with terror" at our respective impending encounters.

In the past I had always read the classics when revising for finals, so that, although I retook examinations regularly and in many different places, I read the works of Chaucer, Dante, and Gibbon. (Unfortunately, there was never a question on them in the subsequent papers.) For my recent postmenopausal fiasco with the membership of the Royal College of General Practitioners I read volume 1 of Proust's *Remembrance of Things Past* in the new Penguin edition. Proust describes how falling asleep in a strange bed leads him to drift back to earlier sleeps in other beds, to encounter again those night terrors of childhood. In the same way becoming a candidate again revived memories of examination horrors for me, those fraught confrontations that can shatter the ego so.

In some of my current nightmares I am late, I rush across London, only to be told that the exam is in oriental studies, or ancient Greece, and I awake sweating. No wonder De Quincey, who himself suffered terrible starvation and poverty in the

capital, projected this privation on to the streets themselves: "So then, Oxford Street, stony-hearted stepmother! thou hast listened to the sighs of orphans, and drunkest the tears of children." How many medical candidates would echo these feelings and thoughts about the roads leading to the royal colleges?

Is it worth it? I can speak only for myself. Meeting old friends again, like phlebotomy, conjunctivitis, Henoch-Schönlein purpura, and carcinoma tumour, or looking once more into Gray's Anatomy to see the stomach bed or the veins of the liver, as well as meeting many new friends like immunoglobulin subunits, and disseminated intravascular coagulation, has made it so. I shall never regret the time I spent reading the syllabus, even though I was tempted to renege. Davidson's *Textbook of Medicine: the Book of Revelations*, as it contained so much that I had long forgotten or was new to me. It was also a new experience to tackle multiple choice question papers. I had never encountered them before, having been only dimly aware of their existence through my children. I found this discipline both difficult and searching and, when revising, I developed a distressing tendency to pun everyone I met to the wall, asking questions like: With antihypertensive treatment, it is correct to say that such and such a phenomenon is likely or unlikely to happen, true or false? No wonder everyone asked me as much as possible.

I even went on a Sunday revision course prepared by the college. If I was depressed before, I was ready for electroconvulsive therapy on leaving. To begin with, everyone else was much younger than I, and that included most of the tutors. Young candidates, fresh from finals, kept leaving to their feet, enumerating long lists of causes of everything. Trial essays were discussed, having been marked earlier. I discovered that mine was next to the bottom. The examiner also remarked that he could not read my writing without great difficulty, and that I should take careful note of his lengthy comments at the foot of the page. Despite considerable effort, I could not decipher his handwriting, either. All in all, not a good day.

41 Gilwood Road, Northenden, Manchester M32 4BR
RAYMOND E GOODMAN, MSc, MRCP, psychosexual therapist and general practitioner

aware of and committed to the teaching programme. The trainee may undertake the bulk of the teaching programme or he may delegate it; he will probably be responsible for the administration and integration of the programme, though some of this work too may reasonably be shared, but he and the trainee must have the support and encouragement of the whole practice team.

(2) *Teaching time*—Without some formal teaching structure in the practice (in addition to day release programmes with other trainees) the training reverts to mere apprenticeship. Tutorials on specific clinical and administrative topics negotiated between trainer and trainee, random case analyses, and the discussion of difficult problems must be a basic part of the training programme and sufficient time set aside for this activity must be one of the team's first order of business.

(3) *Contact with allied professionals*—Practices differ in the make up of the teams they provide, and it should not be absolutely necessary for a training practice to have, say, attached health visitors. What is necessary is contact with other primary care agents—district nurses, health visitors, secretaries, receptionists, and social workers. The trainee must have the opportunity to observe the work of all these colleagues and of acquiring the communication skills needed for satisfactory team work.

(4) *Adequate records*—Records are basic tools of clinical practice and when the care of individual patients is shared, as is invariably the case where a trainee is concerned (the ultimate responsibility remaining with the trainer and his partners), the importance of adequate records is obvious and paramount. Adequacy does not necessarily mean A4 folders, although it is attained more easily with them than with the standard medical record envelopes. Minimum criteria for adequate records should include summary sheet or card with clinical, family, and social data, hospital letters and reports filed in chronological order, and, ideally, regularly "pruned", and recording of long term medication independently of continuation notes.

(5) *Library*—The written word is still one of the most durable, flexible, and economic of educational resources. The trainee should be encouraged to acquire his own library for study and reference, but the training practice needs to have a library of current textbooks and journals which the trainee can consult. The trainee should be able to use the larger reference libraries in local hospitals, postgraduate centres, or universities and should be encouraged to do so for project work, but he must at the same time have more sources of reference close at hand. These, then, it seems to me, are the five minimum standards that should be required of any training general practice. Such fundamental requirements are modest enough to meet O'Sullivan's call for minimal government.

Other desirable attributes

There are many other desirable attributes that are helpful but without which it would still be possible to provide an adequate if not an ideal training programme. Among the more important of these attributes are:

(a) *Regular meetings*—Members of a practice meeting together regularly to discuss both clinical and administrative matters may provide an intellectually stimulating forum for the exchange of ideas with immediate practical impact. Here theory and practice may interact and often provide valuable examples of mutual teaching and learning.

(b) *Special interest*—Not all training practices will have hospital attachments or run antenatal clinics or undertake paediatric surveillance, but such examples of special interests are useful and trainees should at least be aware of potential developments in these fields.

(c) *Membership of professional bodies*—It would be doctrinaire to insist that all trainees should be members or fellows of the Royal College of General Practitioners, but trainees should certainly be encouraged to consider the advantages of membership in helping to foster good morale and high standards and to combat isolationism. Such encouragement is obviously most effective when based on personal example. As the college is the academic arm of general practice, the BMA is its political arm, and all members of the profession who wish to influence (or indeed enjoy) negotiated remuneration and conditions of service should act from within rather than grumble from without. Participation in the affairs of local medical clubs and societies likewise help to provide balance and stimulus for trainees and trainers alike.

Challenge of standard setting

Freeling and Fitton recently suggested that all trainees should be attached to non-teaching practices for one or two weeks during their last three months of training so that they might have some ground for forming opinions as to whether or not the model of practice to which they had become accustomed provided the type of care patients need or want.⁴ This is certainly a question which cries out for an answer. There is a danger of imposing too many criteria for teaching practices and that is that the best could become the enemy of the good. Nevertheless, there must be minimum standards if general practice is to progress.

The environment of the training general practice cannot be "all things to all men" and it should be accepted that on completing his traineeship the trainee as prospective principal may be faced with very different conditions. To meet and work effectively in such conditions he must at least have had the opportunity to see what potential there is for development, and for this he needs basic reference standards.

The challenge to the profession, and especially to the academic leaders of the profession, is not only to establish such reference standards but to continue to evaluate them in an attempt to answer the tantalisingly difficult question: "Do such standards improve the quality of patient care?" The answer may always remain just beyond our grasp, but the question must be asked and continued pursued if we are to avoid the twin dangers of complacency and stagnation. We need standards, but they must be capable of justification.

References

- Joint Committee on Postgraduate Training for General Practice. *Criteria for selection of trainees in general practice*. JCPTGP, 14 Prices Gate, London SW7, 1980.
- Wright J. What is to be taught and learned? In: Cormack JJC, Marinker M, Morrell DC, eds. *Teaching general practice*. London: Kluwer Medical, 1981.
- Donald D. The teaching process. In: Cormack JJC, Marinker M, Morrell DC, eds. *Teaching general practice*. London: Kluwer Medical, 1981.
- Freeling P, Fitton P. Teaching practices revisited. *Br Med J* 1983;287:535-7.