PRACTICE OBSERVED

Practice Research

Evaluation of a course for general practitioners on muscles and joints

ALISTAIR K ROSS, WILLIAM A LAWTON

Abstract

We attempted to evaluate a course on muscle and joint problems for general practitioners using a practice log diary and a factual test paper, which were completed both before and after the course by the doctors attending. A significant improvement was shown in the group score in the test paper taken after the course ($p < 0.001$). Scores in the tests before and after had a negative correlation with age ($p < 0.001$).

In comparing the data in the second log diary with those in the first there was a significant reduction in the prescribing of drugs ($p < 0.001$), use of heat treatment ($p < 0.001$), and in "other" forms of treatment ($p < 0.001$). No appreciable change in investigations or number of hospital referrals was shown. Those doctors who had low scores in the first factual test referred appreciably more patients to hospital and requested more x-ray examinations ($p < 0.01$).

Introduction

A two and a half day residential course on muscle and joint problems was held at Stoke-on-Trent in October 1982 and was attended by 25 general practitioners. The format of the course

was similar to that described by Griffin and Barry and consisted largely of the clinical demonstration of patients to small groups of doctors.¹ Its aims were to improve skills in examination and diagnosis and to demonstrate techniques of treatment, including joint aspiration and local steroid injections. A further objective was to determine whether the course would alter the doctors' management of patients, particularly with regard to hospital referrals and investigations. A fuller description of the course is given elsewhere.²

Method

Before attending the course the doctors were sent a log book, in which they were asked to record the details of 20 consecutive cases of muscle and joint conditions in patients presenting to their practice. Information on each patient was sought on age, sex, the diagnosis or site of the problem, or both; whether the condition was new, old, or recurrent; details of investigations arranged; treatment prescribed, advised, or given; referral elsewhere—for example, hospital, private consultation, domiciliary visits, physiotherapy; and whether the doctor thought that physiotherapy was indicated. Six weeks after the course each doctor was sent a second log book for the details of a further 20 consecutive patients with muscle and joint problems.

At the beginning of the course each doctor completed a test paper consisting of 33 composite questions on rheumatology, most of them to be answered "true, false or don't know." These questions were selected from *Joint Studies in Rheumatology* by Dr Matthew Wilkinson.³ The test paper had a maximum score of 200, and four examples of the questions are given in the appendix. A questionnaire was also completed which gave the following information on the doctor attending: age, sex, past or present experience in orthopaedics or rheumatology; access to a physiotherapist, clinic, or a community physiotherapist; and whether he employed a practice nurse.

At the end of the course each doctor again completed the test paper. The accepted answers to the questions were issued after this test, and each doctor subsequently received by mail a copy of Dr Wilkinson's book. A third opportunity to answer the questions was offered six weeks after the course, when the paper was sent by post with the

averaging 11.5%, patient referrals compared with 22.5%, but this was not significant. Not surprisingly they also referred more patients for physiotherapy (12.1% compared with 4.4%).

To assess the doctors who attended the course, doctors were divided into three groups according to their scores on the first written test, and differences in patient management were sought by using the data from the first log diary, which was kept before attending the

course. The group with the top scores had requested significantly fewer investigations ($p < 0.001$) and given more local steroid injections and "other" treatment ($p < 0.05$). The bottom group had referred appreciably more patients to hospital ($p < 0.01$) and requested more x-ray examinations increased as scores generally decreased ($p < 0.01$) (table IV). These differences could not be attributed to the incidence of new patients in each group. The number of referrals to hospital was almost identical after the course for the high, middle, and low scores as before the course.

TABLE III—Patient profiles from the two log diaries completed by general practitioners

Date	Group before the course	Group after the course
Age (years)	1-93	1-93
Sex	47.4 (16.2)	47.1 (16.6)
Male	265	265
Female	245	245
Not recorded	80	80
No of cases	108	126
New	225	190
Recurrent	80	81
Not recorded	19	39
No of investigations per patient	27	27
1	123	107
2	13	13
3	4	4
4	1	1
5	1	1
6	1	1
7	1	1
8	1	1
9	1	1
10	1	1
11	1	1
12	1	1
13	1	1
14	1	1
15	1	1
16	1	1
17	1	1
18	1	1
19	1	1
20	1	1
21	1	1
22	1	1
23	1	1
24	1	1
25	1	1
26	1	1
27	1	1
28	1	1
29	1	1
30	1	1
31	1	1
32	1	1
33	1	1
34	1	1
35	1	1
36	1	1
37	1	1
38	1	1
39	1	1
40	1	1
41	1	1
42	1	1
43	1	1
44	1	1
45	1	1
46	1	1
47	1	1
48	1	1
49	1	1
50	1	1
51	1	1
52	1	1
53	1	1
54	1	1
55	1	1
56	1	1
57	1	1
58	1	1
59	1	1
60	1	1
61	1	1
62	1	1
63	1	1
64	1	1
65	1	1
66	1	1
67	1	1
68	1	1
69	1	1
70	1	1
71	1	1
72	1	1
73	1	1
74	1	1
75	1	1
76	1	1
77	1	1
78	1	1
79	1	1
80	1	1
81	1	1
82	1	1
83	1	1
84	1	1
85	1	1
86	1	1
87	1	1
88	1	1
89	1	1
90	1	1
91	1	1
92	1	1
93	1	1
94	1	1
95	1	1
96	1	1
97	1	1
98	1	1
99	1	1
100	1	1
101	1	1
102	1	1
103	1	1
104	1	1
105	1	1
106	1	1
107	1	1
108	1	1
109	1	1
110	1	1
111	1	1
112	1	1
113	1	1
114	1	1
115	1	1
116	1	1
117	1	1
118	1	1
119	1	1
120	1	1
121	1	1
122	1	1
123	1	1
124	1	1
125	1	1
126	1	1
127	1	1
128	1	1
129	1	1
130	1	1
131	1	1
132	1	1
133	1	1
134	1	1
135	1	1
136	1	1
137	1	1
138	1	1
139	1	1
140	1	1
141	1	1
142	1	1
143	1	1
144	1	1
145	1	1
146	1	1
147	1	1
148	1	1
149	1	1
150	1	1
151	1	1
152	1	1
153	1	1
154	1	1
155	1	1
156	1	1
157	1	1
158	1	1
159	1	1
160	1	1
161	1	1
162	1	1
163	1	1
164	1	1
165	1	1
166	1	1
167	1	1
168	1	1
169	1	1
170	1	1
171	1	1
172	1	1
173	1	1
174	1	1
175	1	1
176	1	1
177	1	1
178	1	1
179	1	1
180	1	1
181	1	1
182	1	1
183	1	1
184	1	1
185	1	1
186	1	1
187	1	1
188	1	1
189	1	1
190	1	1
191	1	1
192	1	1
193	1	1
194	1	1
195	1	1
196	1	1
197	1	1
198	1	1
199	1	1
200	1	1

* $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$.

TABLE IV—Cross tabulation of management of patients with doctors' scores on the factual test

	Group with top scores	Group with middle scores	Group with bottom scores
No of doctors	160	140	140
No of patients	225	225	225
Specific investigations requested	48	87	87
No of patients referred to hospital	181	12	31
No of x-ray examinations requested	28	43	43
No of local steroid injections requested	16	5	4
No of times "other" treatment was given	39	27	15

* $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$.

*Goodness of fit test used.

second log diary. The doctor was asked to answer the questions under the same conditions as for the previous tests, and to return the completed paper. Also enclosed with the diary was the doctor's score in the previous test papers, together with the group score, range, and mean on each occasion.

The data from the test papers and log diaries were coded, processed, and analysed by computer. Since the conditions under which the third test paper was completed were different from the first two the results are not as comparable but are given below.

An attempt was made to acquire a "control group" (of non-attenders on the course) by writing to the 27 doctors who had responded to the advertisement of the course but did not attend. They were asked (with inducements) to answer the factual test paper and complete a log diary. Only two doctors responded, and this part of the study was abandoned.

Results

GROUP PROFILE

Twenty-five doctors (20 men, five women) attended the course. The average age was 42.6 (range 29 to 65). Six (24%) doctors had experience in orthopaedics or rheumatology in hospital and two held current posts. Sixteen (64%) had direct access to a physiotherapist, clinic or a community physiotherapist, or both, and 21 (84%) employed a practice nurse.

FACTUAL TEST

Ten doctors (40%) scored less than 50% in the test before the course compared with two (8%) in the test after the course. Everyone had a higher score in the second test. The results of the two tests showed a significant increase in the group mean score in the second paper ($p < 0.001$, paired t test).

Doctors who had a practice nurse and access to a physiotherapist had higher average test scores before and after the course than those without such resources, but these differences were not statistically significant. The scores of those with past or present experience in orthopaedics or rheumatology did not differ greatly from the others (table I). There was a significant negative correlation ($p < 0.001$)

Scores plotted against age for general practitioners from tests taken before and after a course on muscle and joint problems.

of scores with age in both tests (figure). Table II gives the group scores on the individual questions where there was an appreciable increase in the score after the course and gives the clinical areas.

TABLE I—Results of factual test

	Before course*	After course*
Group mean	101.8 (51.9)	129.4 (64.7)
Standard error of the mean	8.1	8.1
Doctors with orthopaedics and practice nurse	111.1 (55.8)	133.3 (66.7)
Doctors with no physiotherapist and practice nurse	91.9 (46.0)	117.5 (58.8)
Doctors with rheumatology orthopaedics experience	100.2 (50.1)	133.5 (65.8)
Doctors with no rheumatology orthopaedics experience	101.0 (52.5)	128.7 (64.4)

*Maximum score = 200. ** $p < 0.001$.

TABLE II—Group mean scores on individual questions in which the scores on the second test showed a significant increase over the first

Question	Mean of score	Maximum possible score	Mean of score	Probability of occurring by chance	Clinical area
V	2.08	4	3.85	0.01	Joint aspiration
VI	1.32	4	3.62	0.05	Salicylate treatment of rheumatism
IX	1.18	5	2.78	0.01	Reiter's disease
XI	1.36	5	3.44	0.05	Salicylate treatment of arthritis
XVII	2.44	5	3.60	0.05	Physiotherapy
XXV	2.24	5	3.99	0.001	Physiotherapy
XXXII	1.04	10	6.27	0.001	Comparison of physiotherapy and physiotherapy

* p goodness of fit test used.

We thank Reckitt & Coleman Pharmaceutical Division for sponsoring the muscle and joint course; Dr Matthew Wilkinson for permission to use his questions on rheumatology in the test paper; all the doctors who took the tests and completed the log diaries; the research advisory committee of the North Staffordshire Medical Institute for the grant to carry out the evaluation; and Professor Richard Kempa, department of education, University of Keele, for criticism and advice on this paper.

Appendix

- Which of the following statements are true about aspiration of knee effusion?
  - It can safely be performed by a general practitioner in his surgery. T F D/K*
  - It may be more useful than serum uric acid estimation in the diagnosis of acute gout. T F D/K
  - It is best performed by inserting the needle into the space just behind the upper patella. T F D/K
  - It should be carried out only if a parallel tap can be elicited. T F D/K
- Which of the following features suggest a poor prognosis in a patient with rheumatoid arthritis?
  - Subcutaneous nodules. T F D/K
  - Vasculitis lesions. T F D/K
  - A weakly positive RA latex test. T F D/K
  - High erythrocyte sedimentation rate. T F D/K

- Which of the following are recognised features of Reiter's syndrome?
  - Mouth ulcers. T F D/K
  - Nail pitting. T F D/K
  - Conjunctivitis. T F D/K
  - Swollen tender toes. T F D/K
  - Arthropathy which sometimes responds to a course of tetracycline. T F D/K
- Which of the following drug pairs interact adversely?
  - Phenylbutazone-chlorpromazine. T F D/K
  - Allopurinol-cloxacillin. T F D/K
  - Aspirin-warfarin. T F D/K
  - Indomethacin-bendroflumide. T F D/K
  - Propafenone-salicylate. T F D/K

*T F D/K = True, false, don't know.

References

- Griffin GA, Barry SMK. Muscle and joint pain: design and evaluation of courses for general practitioners. *J R Coll Gen Pract* 1981;31:601-8.
- Ross AK. Practical aspects of muscle-joint problems. *Rheumatology in Practice* 1983;13:125-5.
- Wilkinson M. *Joint studies in rheumatology. A complete course for GPs*. Centre for Medical Education, University of Dundee in association with the Arthritis and Rheumatism Council, 1982.

(Accepted 15 December 1983)

Approval of trainers and training practices in the Oxford region: assessment

THEO P C SCHOFIELD, JOHN C HASLER

Abstract

This is the second paper of three on criteria for approval of trainers of general practitioners drawn up for the Oxford region. This paper describes assessment of trainers and training practices by a team of general practitioners who visit for one day.

Introduction

In the first paper (18 February, p 538) we listed recently accepted criteria for trainers and training practices in the Oxford region and how they were devised by a regional working party. The working party also had the task of recommending new methods of assessment that would allow performance in each criteria to be measured.

Until recently the assessment of trainers in the region had followed a fairly traditional pattern. Doctors applying for the first time were visited for two to three hours by two assessors from a panel of roughly 18 course organisers and senior trainers.

This was followed by an interview with an appointments committee. Reapprovals might require a visit or interview or both. It had become clear that much more information was gained from the visit than from the interview. The working party recommended therefore that for a pilot period of six months a much longer visit should be made for all reapprovals as well as initial approvals, and, as we shall describe in the third paper, such a visit is now made regularly in the region.

PURPOSE OF VISIT

These visits have two purposes. Firstly, they are used to assess both the trainer and his practice against the agreed criteria and to make recommendations to the appointments committee on approval. Secondly, and equally important, the purpose is educational, to assist the trainer and his practice to identify strengths and weaknesses, and to consider the steps that may be taken to build on these strengths and correct any weaknesses.

MAKE UP OF THE TEAM

There are usually three visitors and all trainers and course organisers in the region are invited to be members of visiting teams. Fifty trainers took part in the pilot study and subsequently only six trainers have declined the offer. It is, therefore, assessment largely by one's peers. The regional adviser or associate adviser will usually be included in the team visiting new prospective trainers, and they may also be included in other visits as well.

Medical School Offices, John Radcliffe Hospital, Headington, Oxford OX3 9DU.
THEO P C SCHOFIELD, MRCP, FRCP, associate regional postgraduate adviser for internal practice.
JOHN C HASLER, MRCP, FRCP, regional postgraduate adviser and clinical lecturer in general practice.
Correspondence to Dr Schofield.

TEAM LEADERS

The teams are led by a specially selected senior trainer or course organiser whose responsibility it is to arrange the timing of the visit with the practice and the members of his team and to ensure that they all agree the arrangements. The team leader also coordinates the preparation of the report after the visit.

ORGANISATION OF THE VISIT

The members of the team have a copy of the trainer's application form and the last trainee's report (if it is a reapproval). It is essential that the purpose and nature of the visit is explained to all members of the practice beforehand and a timetable needs to be agreed. This may vary from practice to practice to fit in with the availability of partners and staff. The observation of the practice premises and the way they function is best done at the beginning of the day when the practice is busy and the visitors do not need to be accompanied by the doctor during this time. Lunchtime may be used for the visitors to meet all the partners, or the visitors may prefer to spend this time on their own to consider the topics to be raised during the interview in the afternoon. A specimen timetable might be as follows:

9.00-10.30	Observation of the practice premises and organisation, equipment, and records. Interviews with health visitors and nurses.
10.30-11.00	Discussion with partners.
11.00-11.30	Discussion with trainee.
11.30-12.00	Preparation of report.
12.00-1.00	Lunch for the visitors on their own.
1.00-1.30	Doctor interview.
1.30-2.00	Preparation of report.

It is important that there is some discussion beforehand among the visitors about the details of what to do on the day.

Methods of assessment

Each of the separate methods of assessment provides information to evaluate a number of the criteria. For example, observing and discussing video recorded consultations not only provides information on clinical competence but also on professional values and preparation for and ability to teach. Information on the experience that the trainee obtains in the practice can come not only from the trainee but also from the practice staff, the trainer, the partners, and by direct observation. All the criteria therefore should be considered during each assessment.

OBSERVATION OF THE PRACTICE PREMISES AND ORGANISATION

During this phase the visitors are able to observe both the premises and the way the practice works at a busy time of the day. The visitors may work separately, looking at different parts of the practice, and this is both more economical and less intrusive. It is helpful if the practice staff are given time to talk to the visitors when they are not expected to be performing their normal duties. One visitor usually sits with the appointments clerk monitoring availability of appointments, while another may spend some time with the practice manager. The third may discuss with members of the health team their contributions to training.

Records.—The visitors look at a random sample of the records of the whole practice. The size of the sample varies with the size of the practice but is usually between 50 and 100 records, and the visitors also look at the records of one of the trainee's recent surgeries. They record the proportion of records that are in chronological order, that have clear, legible entries of each consultation, and that contain a completed problem list or summary. They also look to see if there is a record of regular treatment or anticipatory care—for example, blood pressures recorded, smoking histories, and cervical smears taken—when it is appropriate. Their findings are recorded (figure). They also

assess the efforts that the practice makes to improve the records and their overall value for patient management, teaching, and audit.

DISCUSSION WITH PARTNERS

The visitors meet all the partners to discuss the development of the practice as a teaching practice, the support they give the trainee, and their own participation in teaching. This is best done with all the partners sitting down together at a prearranged time rather than in casual encounters in the corridor, which are not really helpful. It also allows partners to ask questions of the visitors about training in general.

DISCUSSION WITH TRAINEE

If the practice is already a training practice the trainee can give a consumer's view of his training. He should be asked to comment on features that he particularly appreciates as well as areas of possible improvement.

VIDEOTAPED RECORDINGS

The visitors watch with the trainee several consultations that he has recorded in a recent surgery. This needs careful arrangement beforehand. Normally the trainee will have become familiar with the use of video equipment during his preparation for training and so the assessment visit should not be the first time on which he has used it himself.

Portable video equipment is available in all schemes in the region and several training practices have now installed their own. It is essential that the informed consent of patients is obtained before they are recorded and a specimen consent form is provided. It is our experience that most patients have no objection to being recorded.

The trainee needs to record a complete surgery, and it is helpful for him to make a list of the patients whose consultations have been recorded—their age, problem, the likely duration of the consultation, and the position on the tape of its beginning.

	Random sample of practice records (100)	Trainee's recent surgery
No. of sample		
Legible entry of each consult		
Contents in chronological order		
Completed summary or problem list	No appropriate	No completed
Complete record continued medication	On medication	Completed
Men and women aged 35-65	No blood pressure recorded	No blood pressure recorded
Women aged 35-65	No cervical smear recorded	No cervical smear recorded
Women aged 20-35	No Ruyter's immune recorded	No Ruyter's immune recorded

Specimen record analysis sheet.

and end. When the consultation is being discussed it is helpful to have the patient's records available for reference.

The discussion focuses not only on the effectiveness of the doctor in the consultation, using the approach by Pendleton *et al.*, but also on the issues that the consultation raises and how they might be used in teaching. If the doctor objects to the use of video tapes one of the visitors sits in a consultation session with him. This is not so useful and has happened only once.

TRAINER INTERVIEW

The purpose of the interview is to clarify and expand the aspects of the practice that have already been assessed and to explore new areas, particularly the trainee's approach to teaching, curriculum planning, and assessment. It is essential that adequate time is allowed for the interview and that it is conducted as a non-judgmental discussion among peers. The team leader will judge what feedback is appropriate at this stage.

The report

The report includes: a factual description of the way that the practice and the trainee achieve each criterion; mention of the particular strengths that have been identified; areas of potential improvement; recommendations for building on strengths and correcting weaknesses.

The report is agreed by all members of the visiting team and is then sent to the trainer and to the appointments committee. The appointments committee is made up entirely of general practitioners and consists of the chairman of the general practice subcommittee of the Oxford region, the two regional advisers, the local course organiser in charge of the scheme, the team leader, a representative of the candidate's local medical committee, and another general practitioner member of the general practice subcommittee. The appointments committee makes the decision about approval or reapproval in the light not only of the visitors' report but also of the reports from previous trainees in the practice. Sometimes the committee think that the information is incomplete or that there are areas that the members wish to discuss with the trainee, and it therefore may invite the trainee for interview before making a decision. This also gives the trainee the opportunity to discuss the report with the appointments committee.

Discussion

This kind of visit by three general practitioners, taking up most of the day and at some stage involving every member of the practice, seems at first sight daunting and time consuming. There are, however, some important points to make.

Firstly, all trainees now participate in making assessments rather than just a small group. So each trainee is being looked at by his colleagues and peers (who may learn as much from the visit as the doctor being visited).

Secondly, it is possible to obtain much more objective evidence on which to make decisions than was the case before. It is far to say that whereas previously we were comparatively short visits it was sometimes difficult to come to a conclusion about whether to approve or not we now believe that we can make that decision much more objectively and fairly.

Thirdly, we believe that the visit must have a strong educational component. Therefore, the visit report is sent to the trainee so that he and the rest of the practice can see what was thought to be good and where it was thought that improvements might take place.

Fourthly, we believe that trainees should know on what basis decisions are being taken. This is another reason why they see their reports (and also the reports their trainees write about them). The standard of report writing has much improved as a result of errors and recommendations based on no evidence are not usually acceptable to the recipients.

Fifthly, once a trainee is appointed he will have a major role in the lives of several, if not many, trainees and have an important influence on the local training scheme. Thus taking one day for the process of approval does not seem very long.

Conclusion

This method of assessment is similar in many ways to the methods described in *What Sort of Doctor*, particularly as it is a peer assessment based on direct observation of a doctor working in his own practice.¹ It differs, however, in two fundamental ways. Firstly, the assessment is based on criteria for a trainee and training practice rather than on just a doctor. Secondly, although the visits have much educational value for both the trainee and the visitors, the "bottom line" is whether the trainee is or is not approved. A degree of discomfort and threat is therefore inevitable and it is essential that the whole process is handled throughout with fairness, respect, and sensitivity.

References

- Pendleton DA, Schofield TPC, Havelock PB, Tate PH. *The consultation: an approach to learning and teaching*. Oxford: Oxford University Press (in press).
- Royal College of General Practitioners. *What sort of doctor?* *J R Coll Gen Pract* 1981;31:698-702.

(Accepted 9 December 1983)

In the third paper we will describe the results of our evaluation of these methods of assessment.

Diary of Urban Marks: 1880-1949

I engaged a boy called Robert Muffky, who was not a Russian, to clean the car and come in with me at a charge of 7 to a week. Muffky travelled me time by winding up the car when it had been stopped at a house. He was with me until 1913 when I took another boy called Kneath who at the time knew nothing about a motor. He had, however, a mechanical turn of mind and through the years taught himself all there was to know about a car. He has been with me now for 20 years and can take a car to pieces and rebuild it. He will have a few nuts and bolts left over but the car will go just as well as before it was dismantled. Kneath married one of my maids and his eldest son is also of a mechanical turn of mind and is now engaged in engineering. Cars are not Kneath's only specialty. He can dismantle a watch as easily as he can a car but the final result is not always so successful. Still, he is a useful man to have around.

A Difficult Case

A trainee patient

S J GILLAM

Seated round a table allocating the morning's visits over coffee, my trainee thrust at me a usually dull and dog-eared folder in my direction. "Ophelia," he said with a playful gleam in his eye, "now she has a number of interesting problems. A good one for you." What, I wondered, were the other partners sniggering about? Ophelia lived in a three room flat on the grim housing estate that comprised a third of our catchment. My memories of this visit haunt me now—the peeling wallpaper, the olfactory mingling of sweat and damp, the assorted bodies lying idly around in the front room, and, above all, Ophelia lying miserably on the only piece of furniture in the barren room she shared with her daughter and grandson.

I had encountered this widowed 54 year old woman several weeks before by proxy. Her daughter Goneril had appeared asking for painkillers for her mother who was confined to bed on account of arthritic pain and "didn't want to bother the doctor." I argued disingenuously that her mother should attend herself unless she really did need a visit. Goneril sighed and left, but my satisfaction was shortlived.

That evening Ophelia generously lumbered into the surgery. I dutifully sought details of her unusual epigastric pain and noted her consumption of every conceivable type of analgesic. She was grossly overweight and had hypertension, and I fear that she even received a little little return on the hazards of obesity. She consented, anyway, to barium meal examination.

Two days later she returned; the abdominal pains were better; she was now suffering chest pains on the left side. Various instincts suggested electrocardiography; they were promptly repressed. She proceeded to see me for half an hour on Goneril's problems: her financial difficulties, her violent unemployed husband, their son, the future.

Over the next few weeks I learnt much about Ophelia's other nine grown children, some of whom were emotionally and materially dependent on her; Ophelia junior with her housing difficulties, Derek in trouble with the law, David, and the rest. I emerged from these dreary meetings as depressed as she was. What was I supposed to do to help her face such insoluble problems? Why me? For Ophelia was one of three or four hopelessly afflicted regular attenders at surgery who provoked a number of questions: Why do certain patients choose to consult the trainee? What, if anything, do they have in common? How do we deal with them?

The answer to the first question seemed simple. Flattering though it was to have patients return to me, sadly my greatest asset was not charisma, but novelty. Reading over Ophelia's case notes showed that she had attended my predecessor still more often and my successor bears witness to comparable

devotion. The cyclical process—embrace of new practitioner, investigation, reconciliation or rejection or both, re-embrace—may be gratefully accepted by trainees whose patience they have temporarily exhausted or whose responses they now anticipate.

Much has been written on "problem patients," how they are perceived, and what characteristics they share.¹ They suffer combinations of physical (often musculoskeletal), social (often related to isolation), and psychological (often depressive) problems. Presenting with intractable physical symptoms ("organ rectitis"), they are often taking a large number of superfluous drugs. Groves has colourfully described four categories of "hateful patient": entitled demanders, manipulative help rejecters, self destructive deniers, and dependent clingers.² My patients fell smoothly into all four groups simultaneously but at least elicited a common sequence of responses: initially anxiety that a diagnosis had been overlooked, then irritation with feelings of inadequacy, and finally weary antipathy with a dash of true compassion.

What was I doing for such patients? The short answer is almost anything. Enthusiasts may subject them (along with the taxpayer) to unnecessary hardship trying out their latest diagnostic and therapeutic philosophy. Major intervention is, however, not desired or required and therefore unlikely to be of more than temporary benefit. More appropriate psychotherapeutic approaches will have been suggested and met with equal resistance. Ophelia has been told many times that her "problems are psychological" and, indeed, concedes her capacity to deny personal conflicts and to project these on to her children. She is dimly aware of the ways in which, from the centre of her web, she has prevented her children "learning to deal with their own problems." While no new insights will alter the facts of her material deprivation, I am sceptical of her own proposed solution: migration to that mecca for the maldesired, a new town.

Lesson learnt

What did I do for Ophelia? The short answer is almost nothing. I offered regular opportunities for indirect expression of her despair but was unable to get her to voice some of the disappointment with herself as wife and mother, which I suspected underlay her depression. I helped her to identify some of the positive aspects of her life but changed nothing. She is more accepting, but also more acquiescent, in status quo. Though temporarily relieved, she is still prone to recurrence of real pain—and I failed to spare her an unnecessary barium meal.

What did I learn from Ophelia and from other "trainee patients"? Firstly, negative feelings constitute important clinical data about patients' psychology—my frustration mirrored hers. Errors in diagnosis and treatment result if such feelings are denied. In the best of circumstances, the psychology of problem patients are remarkably uniform and that neither age nor length of time since qualification greatly affects them.³

46 Linden Avenue, Wembley, Middlesex
S J GILLAM, MB, DCH, trainee general practitioner

Among other things this presumably reflects the prevailing protestant ethic reinforced by unimaginably consistent selection criteria for medical school. How inadequately does medical education prepare you for some of the most demanding of doctor patient transactions!

Secondly, such people provided an early opportunity to exercise essential counselling skills. Different schools of psychotherapy cherish different approaches to the problem of re-orientating consultations, but those who champion simple sympathetic, non-judgmental, and non-directive approaches figure heads or jargon may, I think, claim comparable results. Finally, zealots benefit from exposure of their limitations.

- References
- Klein DJ, Naiman J, Kohnman A, Munro C. Patient characteristics that influence general practitioners from family physicians. *J Fam Pract* 1982;14:801-8.
  - Stimson GV. General practitioners' "troubles" and types of patients. In: Stimson GV. *The sociology of the NHS*. Sociology Review Monograph No 22. University of Keele, 1976.
  - Groves JE. Taking care of the hateful patient. *N Engl J Med* 1978;298:198-7.
  - Malcolm R, Foster RH, Smith C. The problem patient as perceived by family physicians. *J Fam Pract* 1977;5:361-4.

(Accepted 13 January 1984)

What Annoys Me Most

The kerbside consultation

HUGH M BAIRD

The kerbside consultation always takes place away from the safety and sanctity of the surgery and tends to occur in such places as the street, restaurant, or club. It seems almost as if a doctor's appearance in these places is sufficient to stimulate half formed memories of ailments and ill health, which, when recalled, so dominate the mind as to demand instant action. The perpetrator is nearly always someone who could well have come to the surgery and whose problem has probably been present for some time. Very few of these problems are genuinely urgent and when, rarely, some emergency does present in this way, the doctor is seldom asked for advice or requests for advice or information or to discuss hospital reports and the result of investigations, but it is still surprising that an appreciable number of people seem to want to discuss very intimate matters, and though they seem happy to recount details of their piles, ruptures, and menstrual and reproductive disorders, I am certainly not prepared to do so in public.

The kerbside consultation has many variations, all of which are equally annoying. For instance, there is "While you are here doctor would you mind looking at . . ." Or there is the proxy call, which is a request for a home visit left not at the surgery with the receptionist but at a neighbour's house which is thought the doctor may visit later that day or week or month. As a result, not only may the doctor find himself with extra and unplanned house calls, but the message can so easily be forgotten and when subsequently remembered will often result in a late call to an area already visited. Yet another variation is the request for a home visit without actually implicating the person who

made the request. Although the patient may be someone who has not been seen by the doctor for several years, the suggestion will invariably be made to "Tell him you were just passing and have dropped in for a chat."

Management of the kerbside consultation is difficult and often disappointing. There is usually no escape once it has occurred. The sarcasm intended by offers of instant examination in public is not recognised as such and at worst may even be accepted. Bad temper and sheer rudeness seem to have little if any deterrent effect. Prevention is really the best answer, and perhaps if we made ourselves more accessible to the general public—on the telephone, for instance—some of these incidents might not occur at all. Be this as it may, a general practitioner can always try to walk at a brisk pace and look as if he is perpetually on his way to yet another emergency. "Developing situations" may sometimes be recognised early and avoided. For instance, persons seen loitering with intent to waylay may occasionally be dodged, while those who wait at the kerbside with hand outstretched, as if in the act of stopping a bus, may be given a cheery wave in reply as you drive past. For the rest of that day, however, you will probably be left wondering what was really wanted. The car, too, may be left ready for a quick getaway, and some cynical gentlemen once remarked that it was easy to spot a general practitioner's car as it was nearly always reversed into a parking place for just such a purpose.

Least anyone should think that these suggestions are intended to be taken seriously or that I do not enjoy my work as much as I really should, let me hasten to add that they are written very much with tongue in cheek. The kerbside consultation is an occupational hazard that is not necessarily restricted to the medical profession but which must be experienced by lawyers, dentists, vets, and members of many other professions. Undoubtedly, the best way of dealing with it is to employ those basic tools of the trade without which no doctor is adequately equipped for the rough and tumble of general practice. These are patience, tact, diplomacy, self restraint, and, above all, a sense of humour.

Four Winds, Glenalee, Wiltshire DG8 6PU
HUGH M BAIRD, MB, DORSTCOG, general practitioner