PRACTICE OBSERVED

Practice Research

Detection and management of hypertension in general practices in north west London

KARIM H KURJI, ANDREW P HAINES

Abstract

An examination of the practice notes and attached correspondence of 900 patients aged 30 to 65 years in a random sample of 18 general practices in north west London showed that 340 (47%) of 716 patients consulting in a 10 year period had no blood pressure readings in their records. The blood pressure was equal to or above 160 mm Hg systolic or 95 mm Hg diastolic, or both, in 115 (31%) of those whose blood pressures were recorded; 18 (16%) of these were not followed up.

Seventy four patients were being treated for hypertension. Diuretics were the most commonly prescribed drugs. Treatment was started after one blood pressure reading in 34 (46%). Nine of those who had an initial raised blood pressure reading were normotensive on follow up. A further 14 patients had subsequent raised blood pressure readings but were not treated. Sixty one (89%) of the 88 patients with hypertension did not have a blood pressure recording after diagnosis for one or more periods exceeding 12 months. Of 64 hypertensive patients with complete records, 62 (74%) apparently had had no physical examination performed by the general practitioner and 61 (72%) did not seem to have had any

investigations initiated by the general practitioners. Fifteen (35%) of 43 patients taking oral contraceptive pills apparently had no blood pressure recordings during the time they were taking these. The results of this study suggest that there are still deficiencies in the detection and management of hypertension in general practice.

Introduction

There is clear evidence that early detection and adequate treatment of moderate and severe hypertension in the middle aged population can appreciably reduce the risk of stroke, renal damage, and heart failure.¹ The effects of controlling hypertension on mortality from ischaemic heart disease are less clear. Nor is it clear whether controlling mild hypertension carries appreciable benefits. Despite the known benefits of treatment, the results of population surveys have shown that in most cases hypertension is either untreated or inadequately treated.²⁻⁴ The reasons for this deficiency seem to be mainly the lack of adequate screening or case finding, poor patient compliance, and inadequate follow up.

Hypertension may be detected either by mass screening or case finding, the latter being defined as using the consultation as an opportunity to measure blood pressure. Case finding is more economical in general practice, where about 90% of the patients on the practice list may be seen within five years.⁵ The working party of The Royal College of General Practitioners in reporting on *Health and Prevention in Primary Care* recommended that general practitioners measure the blood pressure of all their patients over 30 years of age. Several authors have expressed concern at the quality of blood pressure control in general practice.⁶⁻⁸ Rose and Heller reported that only 38% of hypertensive patients were treated, most with

Of the total of 88 hypertensive patients, 53 (60%) had had blood pressures recorded in the past 12 months, thus allowing mean blood pressures to be determined for this period. Of these, 34 (64%) had a mean blood pressure lower than 160/95, while 19 (36%) had a mean blood pressure higher than 160/95. Sixty one (69%) of the 88 had periods exceeding one year when there were no blood pressure recordings in either the hospital correspondence or the practice records. Advice on one or other form of non-drug treatment (reduction of weight, reduction of salt, transcendental meditation) was given to 28 hypertensive patients in drug treatment. Non-drug treatment alone was used in only one patient. Smoking habits were recorded in nine of the 88 hypertensive patients, all of whom were also on drug treatment.

PHYSICAL EXAMINATION

Of the 88 hypertensive patients, assessment of physical examination could not be made in four because their records were incomplete. Of the remaining 84 patients, 62 (74%) had no evidence in the practice notes of physical examination having been made in the practice apart from checking the blood pressure. Fourteen (17%) had had only their heart or lungs, or both, auscultated, and a further eight (10%) had had other combinations of examinations. Of those patients seen by hospital staff, 15 (40%) of 38 patients had had an examination of the chest, abdomen, femoral pulses, and fundi recorded in the correspondence, and only five had no examination findings recorded in the correspondence.

INVESTIGATIONS

Of the 84 hypertensive patients for whom data on investigations could be assessed, 61 (72%) had had no investigations initiated by the general practitioners, three had had urinalysis, three had had chest x ray examinations requested, and a further 17 had had combinations of other investigations, including one intravenous pyelogram.

SECONDARY HYPERTENSION

Of the 88 patients, nine (13%) had conditions that might have been causally related to hypertension. Eight of the nine patients had renal lesions, while one had an endocrine disorder (acromegaly). Eight of these patients were seen by hospital specialists, and one was not referred. Seven of the nine patients apparently had no investigations initiated by their general practitioners; the two patients with investigations had an intravenous pyelogram and a 24 hour urine sample measured for vanillylmandelic acid respectively.

INTERVIEWS WITH GENERAL PRACTITIONERS

Twelve (67%) of the 18 general practitioners who participated in the study were interviewed. All said that they would always record all the blood pressures, examinations, and investigations that they did, including those with negative findings. Six of the 12 general practitioners had an active policy of case finding for hypertension. Only two of the 12 general practitioners interviewed had not had their sphygmomanometers checked in the past two years, and six had had them checked less than six months ago. There was some variation in the level of blood pressure at which treatment would be started. Ten of the 12 general practitioners were in favour of a specially trained nurse being provided by the district health authority to assist in screening for hypertension.

Discussion

Eighteen separate randomly selected general practices were surveyed; a somewhat larger sample than in previous reports.⁹⁻¹¹ As all the practitioners interviewed (12 of 18) said that they would always record all the blood pressure measurements, examinations, and investigations that they did (including those with negative findings), the results are likely to be a correct reflection of practice. The prevalence of hypertension based on

one raised reading in this study was 31% (115 of 376 patients). Rose and Heller found in their survey of general practitioners notes in one part of inner London an initial raised blood pressure in 45% (74 of 164) of those with blood pressure readings in their notes, despite using the higher cut off point of 160 mm Hg diastolic.⁹ A high proportion of patients (46%) did not have subsequent readings before starting treatment with anti-hypertensive drugs. Hart found that only about 40% of those who initially register a high reading will be classified as hypertensive after three sets of readings.¹²

The proportion of patients consulting their general practitioners in the past 10 years was 81%, for either sex; this figure is lower than that reported (90%) by Ritchie and Currie for a five year period.⁹ Rose and Heller found that half of the patients made contact in one year and two thirds in three years. They suggested that not all contacts are recorded in the notes as they would be by practices that undertake prospective studies. Brent is known to have a high population mobility, with as much as a 20% annual turnover (personal communication) and it is likely that some patients may have left the district but continued to be registered with general practitioners in Brent.

Less than half of those who attended their general practitioners had a blood pressure recording—the proportion was smaller in men than in women. Men in the age group 30-39 are much less likely to have their blood pressures recorded than women in the same age group. This difference might be attributed to the family planning and antenatal services that women of this age group receive, although 15 of 43 women who took oral contraceptive pills had no evidence of a blood pressure measurement in their records.

Sixty nine per cent of hypertensive patients did not have any blood pressure recordings in their practice notes nor in their hospital correspondence for periods longer than one year, suggesting that there is room for improvement in the monitoring of blood pressure in hypertensive patients. Of the 115 patients with an initial raised reading, 74 (64%) were started on treatment. This compares favourably with the 38% (28 of 74) found by Rose and Heller in inner London and 28% (32 of 115) found by Ritchie and Currie in north east Scotland.^{9,13} Most of those on drug treatment were started by their general practitioners. Unlike Rose and Heller, we found no evidence that tranquillisers were being used as antihypertensive drugs. As found by Ritchie and Currie, the most commonly prescribed anti-hypertensive drug was a diuretic. About two thirds of patients on antihypertensive drug treatment had their drug regimen altered, suggesting active participation by the practitioner in patient management. Less than a third of those started on drug treatment, however, were on continuous treatment according to their medical records. Of those patients who had blood pressures recorded in the past year, more than a third had a mean blood pressure above 160/95 mm Hg.

The prevalence of secondary hypertension in a general practice population is likely to be low.¹⁴ Although routine intensive investigation for secondary hypertension by general practitioners is unwarranted since curable causes are rare, particularly in those over 40, it seems reasonable to check a mid-stream urine specimen and plasma urea and electrolytes and to see whether there is evidence of retinopathy or cardiomegaly. Most hypertensive patients in our study had no recorded investigations (72%) or physical examination (74%) by their general practitioners. Hospitals still play a major role in the care of hypertensive patients, with as many as 19% (17 of 88) being referred primarily for this.

Half of the general practitioners interviewed had an active policy of case finding on hypertension screening. The interviews also showed much variation in the threshold level of systolic and diastolic blood pressures above which individual general practitioners would start treatment, which no doubt reflects uncertainties about the value of treating patients with mild hypertension. The results of this study suggest that there are still deficiencies in the detection and management of hypertension in general practice.

tranquillisers. We therefore decided to review the management of hypertension in general practice locally.

Methods

A list of the 172 general practitioners who serve Brent District was obtained from the Brent and Harrow Family Practitioner Committee. Twenty three general practitioners were randomly selected; of these 18 (78%) agreed to participate.

A uniform system of sampling 50 records per practice was devised. The practice list size L was divided by 50 to yield every L/50th record to be examined. When the record selected fell outside the age range 30 to 65, the next record in line would be selected. This was designed to minimise disruption of normal practice activities, the record removed being temporarily replaced with a marker. The practice notes, together with attached correspondence, were examined and relevant data extracted in a standardised fashion by one of us (KK). Information extracted included age, sex, completeness of records, whether the patient had consulted in the 10 year period (1 August 1972 to 1 August 1982), prescription of the oral contraceptive pill, number of blood pressure recordings, and place of recording (general practice or hospital). For those patients who had had blood pressures recorded in the notes, the following data were noted: the duration since the last recording, whether a raised reading had been obtained, the rechecking of a raised reading, and the highest blood pressure readings before treatment.

Data collected on patients with hypertension comprised: the number of readings between the initial raised reading and the start of treatment, details of physical examination and investigations recorded, smoking habits, type of drug and non-drug treatment, changes and continuity of drug treatment, the longest period for which there was no blood pressure reading, the number of and mean of blood pressure measurements in the past 12 months, and whether or not the patient was referred to a hospital.

A brief standard interview was administered to general practitioners to obtain information on their record keeping, their usual practice for the examination, investigation, and treatment of hypertensive patients, the servicing of their sphygmomanometers, and the presence of any screening programme for hypertension in their practice. The data were analysed manually using standard recording forms. Raised blood pressure was defined as a systolic blood pressure of ≥ 160 mm Hg or a diastolic blood pressure of ≥ 95 mm Hg, or both, in keeping with the definition of the World Health Organisation.¹⁵ This does not necessarily imply that all patients fulfilling this definition require treatment but that they at least require follow up and monitoring of their blood pressures.

Results

The sample comprised 900 patients aged between 30 and 65 years; 425 (48%) were men and 462 (52%) women. The sex of 13 patients was not recorded and these were excluded from further analysis. Table 1

TABLE 1—Consultation with general practitioners in the past 10 years by age and sex

Age (years)	Men		Women	
	No consulting (%)	Not consulting (%)	No consulting (%)	Not consulting (%)
30-39	144 (81)	34	157 (84)	31
40-49	96 (75)	32	104 (75)	33
50-59	143 (81)	32	170 (80)	39
60-65	39 (95)	2	36 (90)	4
Total	342 (81)	63	374 (81)	88

shows the proportion of patients who consulted their general practitioners in the past 10 years by age and sex. There was no difference between the proportion of men and women.

DETECTING AND RECOGNISING HYPERTENSION

There were blood pressure recordings in the notes of 300 (34%) of 887 patients in the five year period 1 August 1977 to 1 August 1982,

TABLE 1—Blood pressure recordings by age and sex in 18 general practices

Age (years)	Men		Women		Men and women	
	No who had blood recorded (%)	No who had not recorded (%)	No who had blood recorded (%)	No who had not recorded (%)	No who had blood recorded (%)	No who had not recorded (%)
30-39	144 (81)	34	157 (84)	31	135 (45)	115 (40)
40-49	96 (75)	32	104 (75)	33	121 (41)	115 (40)
50-59	143 (81)	32	170 (80)	39	140 (48)	115 (40)
60-65	39 (95)	2	36 (90)	4	53 (17)	115 (40)
All ages	342 (81)	63	374 (81)	88	716 (38)	115 (40)

patients by age and sex who had had blood pressure recorded in the 18 general practices.

Women were more likely to have had a blood pressure reading than men ($p < 0.005$), particularly in the younger age groups. Of the 376 patients with blood pressure recordings, 73 (19%) had a recording noted only in correspondence from a hospital, 219 (58%) had recordings only in correspondence from a hospital and a further 82 (22%) were recorded in both. Two further patients had blood pressures measured elsewhere—once by a private doctor and one at the Department of Health. Of those with a recorded blood pressure, 115 (31%) had a systolic blood pressure ≥ 160 mm Hg or a diastolic blood pressure ≥ 95 mm Hg. A subsequent recording appeared in the notes of 97 (84%) of these 115 patients. Nine of those with an initial raised blood pressure reading were normotensive on follow up and were not treated. Eighty eight (91%) of the 97 patients who were followed up after subsequent raised readings or were normotensive but receiving treatment with antihypertensive drugs. This group of patients included a few who were started on treatment before 1 August 1972. The records of six of the 48 recognised hypertensive patients were deficient in one or more areas; this accounts for the slight variation in denominators used for calculating proportions. Fourteen patients had subsequent raised readings but were not treated. Of the 43 women identified as having been on the oral contraceptive pill at some time during the 10 year period, 15 (35%) had no evidence of a blood pressure recording anywhere in the notes.

MANAGING HYPERTENSIVE PATIENTS

Of the 88 recognised hypertensive patients, 74 (84%) were on treatment with antihypertensive drugs. Thirty one (42%) of these 74 patients were on diuretics alone, 16 (22%) on β blockers and diuretics, 12 (16%) on β blockers alone, and 15 (20%) on other combinations. Only two patients were treated with a tranquilliser, one for associated anxiety and the other for symptoms of alcohol withdrawal. One hypertensive patient was receiving drugs for weight reduction. Treatment was started by general practitioners in 50 (68%) of 74 patients treated, and by hospital staff in eight (11%) of the 74 patients. In the case of seven patients (9%) it was not clear who had started treatment.

The number of readings made between an initial reading of $\geq 160/95$ and starting drug treatment was two or more in 26 (35%) patients and one in 11 (15%) patients. Thirty four (46%) of the 74 patients did not have subsequent readings before starting treatment with antihypertensive drugs. For a further three patients it was not determined with certainty as their records were incomplete. Of the 74 patients on antihypertensive treatment, eight could not be assessed because the quantities of drugs prescribed had not been entered in the notes. Of the 66 patients who could be assessed, 22 (33%) had stopped drug treatment altogether, 10 (15%) had intermittent lapses of fewer than three months' duration, 14 (21%) had intermittent lapses of more than three months' duration, and 20 (30%) were apparently on continuous treatment according to their medical records. A change in drug treatment, whether by altering dosage or type of drug, occurred in 48 (65%) of the 74 patients.

We are grateful to the Brent and Harrow Local Medical Committee and the general practitioners of Brent Health District for their cooperation. We also thank Professor David Miller, Dr R F Heller, Dr L J Edwards, and Dr M A Newton for their advice and encouragement.

References

1. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Effects of treatment on morbidity in hypertension. Results in patients with diastolic blood pressures averaging 115 through 125 mm Hg. *JAMA* 1967;202:1028-54.
2. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Effects of treatment on morbidity in hypertension. II. Results in patients with diastolic blood pressures averaging 114 mm Hg. *JAMA* 1970;213:1143-52.
3. Schoenberg J, Stamler J, Skeels RB, Skeels S. Current status of hypertension control in an industrial population. *JAMA* 1972;222:559-62.

4. Heller RF. Detection and treatment of hypertension in an inner London community. *Br J Prev Med* 1978;30:208-12.
5. Health and prevention in primary care. Report from General Practice 18. London: Royal College of General Practitioners, 1978.
6. Barlow DH, Bevers DG, Hawthorne WM, Watt HD, Young GAR. Blood pressure measurement at screening and in general practice. *Br Heart J* 1977;38:2-12.
7. Heller RF, Rose G. Current management of hypertension in general practice. *Br Med J* 1973;1:1442-4.
8. World Health Organisation. Arterial hypertension. Report of a WHO Expert Committee. Technical Report Series No 628. Geneva: WHO, 1978.
9. Ritchie LD, Currie AM. Blood pressure recording by general practitioners in north-west Scotland. *Br Med J* 1978;2:1220-3.
10. Hart JT. Semi-continuous screening of a whole community for hypertension. *Lancet* 1970;1:223-6.
11. Berglund G, Anderson G, Wilhelmsen L. Prevalence of primary and secondary hypertension: study in a random population sample. *Br Med J* 1976;1:554-6.
12. Anonymous. Looking after patients with high blood pressure. [Editorial.] *J R Coll Gen Pract* 1976;26:235-6.
13. Accepted 6 January 1984.

Quality of care in managing hypertension by case finding in north west London

GERALD MICHAEL

Abstract

The records of patients who were registered with general practices in inner London and outer London (both training and non-training practices) were examined for details of blood pressure recordings. A blood pressure recording was found in the past five years in only 22% of the notes of inner London practices but in 45% of notes in outer London practices, there being no difference between training and non-training practices. In 31% of the notes with a blood pressure recording the blood pressure had been raised ($\geq 160/95$) at some time. In 65% of these notes for patients aged 65 or more and in 82% of these notes for patients aged under 65 treatment had been given at some time. Thirty eight per cent of patients in the older group and 38% in the younger group were being treated at the time of the survey, this being the most commonly used drugs. Thirty four per cent of all those with a raised blood pressure at some time had a latest recording of under 160 mm Hg systolic and 95 mm Hg diastolic.

There seems to be a continuing need for general practitioners to detect and treat hypertension, and it is suggested that general practitioners should consider using systematic case finding for hypertension in their practices.

Introduction

Many studies have shown that hypertension is a symptom free condition.¹ Since treatment reduces mortality and morbidity,² it is important to check the blood pressure of every adult and to

give effective management when the level is raised. Obviously patients should be inconvenienced by the regimen and its side effects as little as possible. In 1974 Heller and Rose looked into general practice in inner London to determine the number of patients whose blood pressure had been recorded, and for those patients who had been found to have hypertension the quality of care that they received.³

This paper presents the results of a further inquiry done in 1982 in a similar area of inner London, but the study also included practices in outer London. The inquiry was to examine, firstly, any differences in the recording rates for blood pressure in practices in inner London and outer London, and secondly, the differences in recording rates between training and non-training practices.

Methods

Six practices in each of three groups were invited to participate. The first was a group of six training practices in outer London—Barnet, Brent, and Harrow family practitioner areas—comprising all the training practices that had trainees from the vocational training scheme at Edgeware General Hospital. The second was a group of six practices in the same area of outer London matched as far as possible for type and size of the training group. The third group was six inner London non-training practices chosen by an inner London course organiser for vocational training, who was asked to provide a list of practices containing a mixture of sizes from singlehanded to group. Two specially trained research assistants carefully examined all the contents of the record envelopes, including day to day continuation cards, antenatal cooperation cards, family planning cards, and any letters. Any record of a blood pressure measurement in the past five years was noted.

A pilot study in one single handed non-training practice in outer London was performed to estimate the sample size required to show a 10% difference in the rate of recording blood pressures between each of the three groups of practices. The results showed that 2300 records of patients over the age of 24 had to be examined overall. A systematic sample of each of the participating practices was drawn (each 48th

42 London Road, Stanmore, Middlesex HA7 4NU
GERALD MICHAEL, MB, MRCP, general practitioner

recorded, so that the same proportion of notes was sampled in each practice. It was found that the 48th record was that of someone younger than 24, the 49th record was examined and then subsequently the 47th, to keep the rate steady at one in every 48.

The records were also studied to see if either the systolic or the diastolic pressure had been recorded as being $>160/90$. It was noted how many hypertensive patients had reached a diastolic pressure of less than 90 at the time of their last recording. Statistical analysis of the difference between the proportion of notes with a blood pressure recording in each practice group was made using the χ^2 test. Comparisons were not made between the different practice groups for raised blood pressure, treatment, and control of hypertension.

Results

No training practice refused to take part. One outer London practice with two partners refused, one inner London practice refused, and another inner London practice was reluctant, so was not measured.

BLOOD PRESSURE RECORDED

Of the 2374 records examined, 909 (38%) included a blood pressure recording in the past five years (table 1). The differences between training and non-training practices in outer London were small, but,

TABLE 1—Tendency to record blood pressure (BP) in different practices in the past five years

Outer London training practice	No in sample	BP recorded	
		No	%
1	295	122	41
2	109	66	61
3	129	84	65
4	119	53	45
Total	652	325	50
Non-training practice			
1	74	26	35
2	105	50	48
3	129	84	65
4	119	53	45
Total	427	213	50
Inner London			
1	83	11	13
2	105	50	48
3	129	84	65
4	119	53	45
Total	436	208	48
Grand total	2374	909	38

overall, patients in outer London were more likely to have had their blood pressure recorded than those in inner London ($\chi^2=82.96$, $p<0.05$). One inner London practice had only one blood pressure recording in 83 notes studied, and the proportion of notes with a blood pressure recording was lower in each of the four inner London practices than in any outer London practice. Patients who had been registered for four or five years had a greater chance of having had their blood pressure recorded.

There was a change in the recording pattern with age: 30% of those under 45 had a blood pressure recording in the notes, rising to 55% of those 65 or more. Young women were more likely than young men to have had a blood pressure recording.

RAISED BLOOD PRESSURE

Of the 909 blood pressure recordings where the blood pressure was recorded, the pressure was raised (160 mmHg or more systolic or 90 mmHg diastolic) in 268 (31%) patients at some time. The proportion of blood

pressure recordings which were raised increased with increasing age, from only 4% in the age group 25-34 to 60% of those aged 65 or more.

TREATMENT

Ninety (62%) of the 143 patients with raised blood pressure aged 65 or over and 74 (52%) of the 141 patients under 65 had been treated at some time; 54 (38%) patients in the older group and 47 (33%) in the younger group were still being treated at the end of the survey and were receiving a total of 164 different prescriptions (table 1). Thiazide diuretics were prescribed most often.

TABLE 1—Drugs prescribed to 101 patients being treated for high blood pressure

Drugs	No.
Thiazide diuretics	22
Other diuretics	11
Other antihypertensives	11
Other hypertensive treatment, such as methylglutamine	20
Sedatives or tranquillisants	5

Note: It was impossible to determine whether the sedatives and tranquillisants were being used to treat the hypertension or whether they were prescribed to treat other conditions.

CONTROL OF BLOOD PRESSURE

Table 111 gives the latest recorded diastolic pressure. Overall, in 34% of the hypertensive patients the recording was under 90 mmHg diastolic, and in those younger than 65, 33%, had a blood pressure recording of under 90 mmHg diastolic.

TABLE 111—Latest recorded blood pressure in 286 patients who at some time had hypertension ($>160/90$)

Age (years)	Diastolic pressure	Total
	90-100	110+
65-75	51	85
75-85	65	64
85-95	15	10
Total	97	179

Discussion

Heller and Rose reported that in 1974 in one part of inner London 24% of patients aged 20 and over had had their blood pressure recorded.⁴ Eight years later there seems to have been no change in inner London, although more than 60% of patients registered with practices in outer London had a blood pressure recording in the notes. Although the longer a patient has been registered with a doctor the more he or she is likely to have had the blood pressure recorded, the differences are not great enough to account for the low rates in inner London, both in 1974 and now, just on a high rate of turnover.

Fleming and Lawrence examined the records of self selected doctors who did a practice activity analysis before attending a residential course in 1980 and found that between 1975 and 1979 some 51% of patients' records had a blood pressure noted.⁵ Because of the nature of this course it is likely that the doctors' recording figures were atypical and it might be assumed that because of the problems of inner cities the results in outer London are lower than average. It may be, therefore, that our results from outer London are the more usual—that is, about 43% of the adult population has had the blood pressure recorded.

There was no difference between training and non-training practices in outer London. One of the main criteria for selecting a trainee is that he has a high standard of practice, and it was

Minimum Standards for Training

Top 10 points for trainers

DENIS DURNIO

Over the past 10 years, as a result of much hard work by trainers, advisers, course organisers, and many official bodies, vocational training for general practice has earned its place as a professionally challenging postgraduate career. Now, for many reasons, the organisations most closely concerned in setting standards—trainers, the Royal College of General Practitioners, the General Medical Services Committee, and the Joint Committee for Postgraduate Training in General Practice—are anxious to review and, if necessary, improve the criteria for the selection of trainers in training practices.

I have carefully reviewed the criteria in my own region and cast the fairest eye a regional adviser might have over some of the most recent publications from 14 Princes Gate (the Royal College of General Practitioners) and offer what can be defined only as a personal view of essential and desirable criteria for selection as a trainer. Although I suggest 10 essential features for a trainer, they are not in rank order, but in my view the most important characteristics of a good trainer are enthusiasm, time to teach, and structure. It was, however, encouraging to find

Royal College of General Practitioners) is considered to be the normal way to assess clinical competence. Though some would wish to challenge this view, offering an alternative method of assessment is less easy. The document *What Sort of Doctor*, published by the college in 1981, goes some way towards a solution.¹ It suggests ways of looking at the performance of established practitioners, particularly in regard to professional values, accessibility, clinical competence, and an ability to communicate. One advantage of the idea is its flexibility for regions—for example, the examination of a sample of video tapes of consultations and the discussion between staff and the visitors. The controversy over whether the MRCPG should be a requirement of prospective trainers may be resolved soon, as most of our new applicants are former trainees most of whom sit the college examination—and eventually pass it.

Courses for trainers

I have been slower than most to agree that a course for trainers should be completed before an applicant is accepted as a trainer. Applicants who have attended a course make a better impression at the interview. They are more familiar with how to structure time for teaching and translate their experience as general practitioners into effective teaching. Visits to prospective training practices by representatives of the trainer selection subcommittee is an important step in assessment. One quickly senses how welcome the building and staff make you feel. Cups of tea or coffee are usually offered; certainly their absence is always noted. I will be more than ready to declare a bias against inadequate premises. There can be no excuse for poor premises, even less for a touch of paint, tidiness, or paper towels on the couch. It is now over four years since my practice moved into a purpose built health centre. The last memory of our move was the way in which the increased space made everyone seem more relaxed. This was in addition to the obvious surge in enthusiasm which we all felt for our work.

The statement of the Joint Committee for Postgraduate Training that all records in teaching practices should have their contents in chronological order by the end of 1984 is a modest request. Records are still far from being satisfactory in teaching practices. Trainers are most concerned to see more summaries, drug sheets, a note of occupation and telephone numbers, and notes written up for any consultation. To my knowledge only two of our training practices have all their records summarised.

I am sorry only that more trainees do not have the opportunity to make such summaries. Fortunately, some of our training practices are working with computers and more trainees should be given the opportunity to see some of the advantages and disadvantages at first hand.

Some teaching practices have an arrangement whereby all partners and attached medical staff meet early each day for about half an hour. Problems from the previous day's work are discussed, including any night calls, mail is read, and certain parts of the day's work allocated. This is excellent—what I call

my subjective view so well supported by the recent publication from the college, *The Influence of Trainers on Trainees in General Practice*.²

Quantifying enthusiasm is not easy. Indeed the enthusiasm shown during an interview of a prospective trainer may never be sustained during the stress of teaching a trainee for a whole year. At the interview of a prospective trainer I am often struck by my very early reaction of "He'll be OK" or, perhaps, "...not OK", and the rest of the interview is spent trying to find evidence to support my intuitive feeling.

In some regions having the MRCPG (membership of the

Foresterhill Health Centre, Westburn Road, Aberdeen AB9 2AY

DENIS DURNIO, MR, FRGP, regional adviser in general practice, Grampian region

expected, therefore, that some trainees would have faced the problem of case finding, if not screening, for hypertension.³⁻⁵ All that may be said is that training has not distracted trainers, and their partners from taking and recording blood pressures.

If our figures for outer London are typical it is worrying that 57% of adults seen never to have had their blood pressure recorded. Fifty-two per cent of patients with a raised blood pressure recorded who were aged under 65 were offered treatment at some time, but only 33% of patients are still having treatment, and only 33% achieved a diastolic pressure of <90 mmHg. Not all patients with a blood pressure $>160/90$ need treatment—for example, neither those whose readings settle spontaneously on second and subsequent readings, nor many elderly patients whose levels are only moderately raised.

It was reassuring that nearly all the regimens were considered to include standard effective drugs for hypertension. Although β -blockers are widely thought to give better cardiac protection than thiazides, many more thiazides were prescribed.⁶⁻⁸ This may be because many patients were controlled with thiazides before the advantages of β -blockers were known. It will be interesting to see if there is a movement from thiazides to β -blockers in the future.

The blood pressure recording rate needs to be improved, because it is a simple procedure that leads to the detection of a dangerous but treatable condition. Probably the easiest way to improve detection rates is by cascading.⁹ This means that arrangements are made for every patient who attends surgery to have the blood pressure taken and recorded in the notes at least once every five years. This may be done by the doctor, nurse, or a trained recorder. It needs to be recorded in such a way that it is known to have been done and therefore is not unnecessarily repeated every time the patient attends. It is hoped that studies evaluating the benefits of such case finding in general practice might be set up to give practical guidelines so that the detection rate of hypertension may be improved.

My research assistants were Mr R Fine and Miss Nikki Gale. The project was supported by a grant from the North West Thames

Regional Health Authority. I thank Dr R Heller of the department of community medicine, St Thomas's Hospital for advice and encouragement.

References

1. Armit J, Pratt JH. Nature of the symptoms associated with essential hypertension. *Br Med J* 1981; 1: 875-81.
2. Weiss NS. Relation of high blood pressure to headache, epistaxis and selected other symptoms. *Br Med J* 1982; 1: 101-3.
3. Traftord JAP, Horn CR, O'Neil H, et al. Five year follow-up of hypertension in mild and moderate hypertension. *Br Med J* 1981; 2: 111-15.
4. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Effects of treatment on morbidity in hypertension: results in patients with diastolic blood pressure averaging 115 through 129 mmHg. *JAMA* 1982; 202: 102-34.
5. Veterans Administration Cooperative Study Group on Antihypertensive Agents. Effects of treatment on morbidity in hypertension: results in patients with diastolic blood pressure averaging 115 through 129 mmHg. *JAMA* 1970; 213: 1143-52.
6. Heller RB, Beevers DG. Current management of hypertension in general practice. *Br Med J* 1977; 1: 1442-4.
7. Fleming DM, Lawrence MSA. An evaluation of recorded information about prescriptive measures in practice. *Br J Clin Pharm* 1981; 21: 615-20.
8. Hunter JR, Swift J. The history of vocational training for general practice. *Br J Clin Pharm* 1979; 28: 24-32.
9. Irvine D. *Teaching Practice*. Report from General Practice No 15. London: Royal College of General Practitioners, 1981.
10. Joint Committee on Postgraduate Training for General Practice. *Criteria for the selection of trainers in general practice*. London: JCGTP, 1979, reprinted 1980.
11. Lambert DM. Effect of propranolol on mortality in patients with angina. *Postgrad Med J* 1979; 55: 57-60.
12. Beevers DG, Johnson JH, Larkin H, et al. Clinical evidence that beta-adrenergic blockers prevent more cardiovascular complications than other antihypertensive drugs. *Drugs* 1983; 25 (suppl 2): 326-30.
13. Stewart IM. Complicated evidence of his myocardial infarction in hypertensive patients under treatment containing propranolol: excluding beta-receptor blockade. *Clin Sci Mol Biol* 1978; 81: 509-11.
14. Barber JH, Beevers DG, Fife R, et al. Blood pressure screening and supervision in general practice. *Br Med J* 1979; 1: 843-6.

(Accepted 23 December 1983)

Diary of Urban Marks: 1880-1949

Not many of the practitioners of Swansea volunteered for active service at first but under the army's 100 000 scheme nearly every one of us volunteered when called on to do so. But it was found necessary by the government to veto the medical profession joining en bloc and an order was issued that at least one of the partners in each practice taken in the army unit was to be called on to do so. Through the British Medical Association a medical war committee was set up in every division of the BMA. Dr Beag, who happened to be chairman of the division at the time, and myself, who was secretary, were appointed to those positions on the war committee. Whenever the government requested medical men to replace those who were on active service each division throughout the country had to give up its particular quota. The committee met every month and went through the list of names in the area. The needs of each district were considered, the age, and family responsibility of each medical man, together with a consideration of their health as to whether they could stand up to active service and if not to what they could do at home in regard to helping the medical service in or around their own district. Gradually a list was evolved in order of merit to speak, and every man next due for service was warned. As the war progressed it so happened that I knew the town and its needs very well and Beag as a consultant knew the district well, in consequence of which the committee were very largely dependent on us. In March 1917 there was a great slaughter of medical men and nurses on the western front. Some hundreds of medical men were put hors de combat at one swoop. The government sent out an urgent order for replacements

and our committee called on every man in the area to attend its meeting on a certain Sunday at the hospital. It was the largest meeting of medical men I have ever seen. The courtyard was jammed with cars. We started our investigations at 3 pm and did not finish until midnight. It was not a question then of considering the public needs at home but merely a question of supplying the army and letting the public take care of itself. Every man under 40 years of age was considered. Eventually my turn came to be considered. I told the committee what work I was doing and left the room for them to consider. On my return, Dr Beag told me that the committee had decided to add my name to the list of those available for active service.

ONE HUNDRED YEARS AGO

We learn from the *Philadelphia Medical News* that General Grant, in a fall on the ice on December 24th, struck heavily upon the outer and upper portion of his left thigh. The pain was extreme, and the inability to use the limb complete. The fall, the age of the patient, and the subjective symptoms, suggested probably a fracture of the neck of the femur, and the complete examination of the limb, which alone was possible at first because of the pain, did not entirely exclude this diagnosis. After the lapse of thirty-six hours, the pain subsided, and then a thorough examination, made by Dr. Fordyce Barker and Dr. L. A. Stimson, failed to discover any objective signs of fracture. The injury appeared to have been a severe contusion of the outer and upper portion of the left thigh, with possible rupture of muscular fibres or attachments in the effort to avoid the fall. The surgeons in attendance confidently anticipated an early recovery. (*British Medical Journal* 1884; 1: 177.)

good general practice—but really it is only an adjunct to time set aside for teaching which allows for uninterrupted sessions between the trainer and trainee. The amount of time for this may be negotiable between regions or indeed in the same practice between trainers as their needs vary. If I had to set a golden rule for training in general practice, however, it would be that two to three hours was the norm in all teaching practices. This commitment in our region is underlined by inviting trainers to provide a detailed timetable for teaching. This statement linked to trainer reports suggests that we have a satisfactory arrangement of training in general practice.

As training for general practice develops and with more visits by the Joint Committee to gauge more importance has been placed on the availability of good current books and journals on general practice in teaching practices. Despite this worthwhile development, I suspect that this facade of availability may still conceal a lack of consistent use of books and articles as a teaching resource.

Support for trainer

The importance of partners and ancillary staff in a teaching practice being sympathetic to training is well established. Indeed, on occasions it may be necessary to nominate a partner to be responsible for a trainee when a trainer becomes unexpectedly ill. Partners will also be expected to share special knowledge with trainees—for example, about police work, paediatric surveillance, family planning, or geriatric assessment. Some practices that I know take this stage further and partners regularly take a day's teaching each week or even a full month's supervision of the trainee, though under the watchful eye of the trainer. Should the partners of trainers attend courses for trainers? This is worth considering for several reasons, but especially because some practices assume that a partner will succeed automatically a senior trainee who retires. It is customary, especially early on in a training year, that a trainee spends time with receptionists at work and sees at first hand the skills of the nurses and health visitors. Nothing will upset a trainee year more than friction between these colleagues, indeed, feedback from these people is invaluable for assessment. At a recent residential course for trainers in our region we looked at assessment in some depth in groups of five or six trainers, all of whom felt that the subject was difficult. The aim of the group work was to construct an assessment sheet that

was relevant to general practice. Each group came to the group leader not knowing what the previous group had agreed. At the end of six sessions we were left with what was in effect a copy of a Manchester rating. Trainers reported that using a Manchester rating with their trainees may take up to two hours to complete. If repeated twice in a year, which allows any change to be noted, this may be a useful exercise.

The greater use of portable video equipment in teaching practices is a major advance in measuring and understanding the needs of trainees in the consultation. Our invitation to trainers to make video recordings of teaching sessions has proved less successful, but this is a topic which must develop soon. Tapes of teaching sessions might be useful for the reapproval of trainees.

Any prospective trainer should have a working knowledge of teaching styles and methods. Portable video equipment is widely available but audio tape recording has a place and is simpler to use, provided a good quality microphone is used. Also, sitting in with a trainee and observing him at work gives instant and useful pointers, which may be of great practical use to the trainee. Those wishing to be trainers would benefit from reading the sections on teaching styles and group work in *The Future General Practitioner*, still one of the most helpful publications in general practice in the last 10 years.

My list of "desirable criteria" are self explanatory. Perhaps our list is worth a comment in suggesting that a prospective trainer should be enthusiastic, and perhaps it is appropriate to end by saying that I also believe that a trainer should enjoy most of what he does. His influence on young trainees will be considerable—let us hope he finds it fun most of the time.

As a regional adviser I hope to remain close enough to the grass roots to have views that are little but not too far ahead of the consensus view on training. You will, I hope, feel that my feet are still on the ground!

References

1. Freeman J, Roberts J, Metcalfe D, Hillier V. *The influence of trainers on trainees in general practice*. Occasional Paper 21. London: Royal College of General Practitioners, 1982.
2. Working Party Report. *What sort of doctor*. London: Royal College of General Practitioners, 1981.
3. Royal College of General Practitioners. *The future general practitioner—learning and teaching*. London: BMA, 1972.

ONE HUNDRED YEARS AGO. A report by Mr. George Turner to the Buntingford Sanitary Authority contains an account of an epidemic of scarlet fever, which was spread in a remarkable manner through the agency of a family van in which the first case appeared. Early in the spring the van stopped at Thelfield, where there were cases of scarlet fever. The man's child, who was then a year old, was ill with a sore throat, next the man himself, his brother, and a son, suffered in a similar manner. The van proceeded to Croft, the village in which the proprietor lived, and there the disease quickly spread, and the man called on his sister, living at Croft, to see his daughter. This family subsequently suffered from scarlet fever, and communicated the disease to those living near them. The van went next to school and club feasts at Yardley, on Whit-Monday, and several cases occurred there in consequence. On May 30th it went to Chipping, and on June 1st some children who had visited the van at the fair were attacked by the fever. These were the first cases of which the authority received any information, and they were visited by Mr. Turner on June 8th. Next day Munden was seen next destination of the van, where it stayed from June 1st to June 8th, at the result of which several cases of scarlet fever appeared in this village also. During the next three days the van remained at Wade's Hill, where the disease made its appearance, but Mr. Turner was unable to trace any definite connection between the cases in this and adjoining places and the visit of the van. The mischief caused by this family did not cease here, for a relation

in the same line of business travelled with them, and so a separate family living at Doe Bridge suffered in the same way from scarlet fever. The van then went to school and feasts at other villages. The sanitary authority properly decided to prosecute the man, but the proceedings unfortunately fell through, owing to the fact that when it came to the medical officer to inform his patient at the outset of the nature of the disease. With a view of preventing similar occurrences, Mr. Turner strongly advocates the abolition of fairs, which when effected, would be followed by the disappearance of the travelling vans. The small local fairs are deemed as nothing more or less than public nuisances, no business worthy of the name is transacted, and of such a nature as to be detrimental to the health of the community. Mr. Turner suggests that the van be restricted to converting a quiet village into a pandemonium for a day or two. A similar circumstance, though, unfortunately, not of such an extensive nature, is mentioned by Mr. Welsby, where a case of smallpox was imported by a Whit-Monday holidaymaker. In referring to the incident, Dr. C. E. Saunders observes that there is scarcely any public holiday in the county, and that the disease was introduced in some part of his extensive district by a case of infectious disease, due to the visit of fairs, or persons themselves visiting infected districts. The congregation of large numbers of persons at fairs, and the consequent dissemination of zymotic disease, is no new evil, and efforts have been made in some parts of the county to stop these gatherings, though so far with very partial success. (*British Medical Journal* 1884; 1: 438.)