- ³ Gillie O, Chisholm J. *The "Sunday Times" self-help directory*. London: Times Newspapers, 1976.
- ⁴ Gee JL, Gee RC. *1983 GP guide*. Petersfield: Asgard Publishing Co, 1983.
- ⁵ Smith R. Part time agony aunt in trousers. *Br Med J* 1983;287:1029-32.
- ⁶ Richardson A, Goodman M. *Self-help and social care: mutual aid organisations in practice*. London: Policy Studies Institute, 1983.
- ⁷ Guardian. *Directory of pressure groups and representative organisations*. Epsom: Bowker Publications, 1979.
- ⁸ Knight S. *Help! I need somebody. A guide to national associations for people in need*. 3rd ed. London: Henry Kimpton, 1980.
- ⁹ Moorehead C. *Helping, a guide to voluntary work*. London: Macdonald and Janes, 1975.
- ¹⁰ Patients' Association. *Self help and the patient*. London: Patients' Association, 1982.
- ¹¹ Share Community. *Guide to self-help groups*. London: Share Community Ltd, 1980.
- ¹² Thames Television. *Health help*. London: Thames Television, 1978.
- ¹³ Todd J. *Someone to talk to. Directory of self-help and support services*. London: Mental Health Foundation, 1982.
- ¹⁴ Gillie O, Price A, Robinson S. *Sunday Times self-help directory*. London: Granada, 1982.

What carbohydrate foods should diabetics eat?

Before the days of insulin and oral hypoglycaemic agents little could be offered to diabetic patients other than dietary advice. Drug treatment heralded an era of diminishing enthusiasm for the role of diet in the management of diabetes—to the extent that in the early 1970s standard medical textbooks devoted no more than a few paragraphs to diet¹ and many hospitals did not have a dietitian to advise diabetic patients. Around that time West reviewed the attitudes of diabetic patients in North America towards the dietary advice they had been given.² Some 22% claimed not to have received any advice, and 25% said that they had received some advice but had made no attempt to follow it. Only about half of all American diabetics claimed to be following a prescribed diet, and much the same was true in Britain.³

During the mid- to late 1970s attitudes began to change. Epidemiological evidence suggested that the standard diabetic diet, low in carbohydrate and relatively high in fat, might actually increase the risk of cardiovascular disease, the most frequent cause of death among diabetics.⁴ Furthermore, several studies suggested that diets relatively high in fibre rich carbohydrate reduced hyperglycaemia in diabetic patients.⁵⁻⁸ Though there remains some disagreement about the precise quantity of carbohydrate which should be included in the diabetic diet,⁹ the diabetic associations of many Western countries, including the United States,¹⁰ Britain,¹¹ Canada,¹² Australia,¹³ and Finland,¹⁴ have affirmed the importance of diet in the management of diabetes and recommended an increase in unrefined carbohydrate (especially fibre rich carbohydrate) with a reduction in the fat intake. Nevertheless, restriction of monosaccharides and disaccharides remains a cornerstone of diabetic dietary policy and clear advice is needed on which carbohydrate foods are the most suitable for diabetic patients.

Two approaches may be used to answer this question: short term experiments examining the response of the blood glucose concentration to various foods and longer term studies. Most investigators have used the first approach. The greatest number of foods has been studied by Jenkins and his colleagues,¹⁵ who are particularly well known for their work showing the reduction of postprandial glycaemia by the gel forming fibre guar.¹⁶ In a comparison of 15 carbohydrate containing foods they found that dried leguminous seeds—for example, kidney beans, red lentils, black eyed peas, and chick peas—produced

a glycaemic response substantially lower (44% to 71%) than that of wholemeal bread. These findings are not surprising since longer term studies have shown a substantial improvement in diabetic control when gel forming fibres from such leguminous sources have been added to the diabetic diet in substantial quantities.⁷ Glycaemic responses for rice, spaghetti, and All Bran were in an intermediate category, being higher than those for beans but below those for bread. Surprisingly there was no difference between white and wholemeal bread. In these circumstances All Bran appeared to be a relatively "good" food despite the fact that one third of its total carbohydrate is made up of disaccharides.

The work of Crapo and her colleagues has aroused interest because they found that potato has a glycaemic response similar to that of dextrose—both worse than bread; white rice and corn were better than bread.¹⁷ In the light of these observations and others in non-diabetics^{18 19} the recent report by Bantle and coworkers should not have caused much surprise.²⁰ In a well conducted study they found that sucrose, fed as part of a mixed meal, did not produce higher plasma glucose concentrations than when carbohydrate was derived chiefly from the unrefined sources of potato and wheat. Meals containing fructose resulted in lower glycaemic excursions.

Nevertheless, taken together these studies raise some questions concerning two fundamental aspects of the new dietary recommendations. Firstly, are all starchy carbohydrate foods rich in dietary fibre suitable for diabetics? Secondly, is restriction of sugar really of crucial importance?

On the first issue undoubtedly foods rich in gel forming fibre (found particularly in various types of cooked dried beans) are especially useful in the diabetic diet. The results of long term studies⁷ have confirmed those of the shorter experiments.¹⁵ Furthermore, these foods provide a useful source of protein, allowing the diet to contain less animal protein, which is so often high in saturated fat, and they also lower serum cholesterol concentrations.²¹ The main reason for this beneficial effect of beans compared with all other carbohydrate foods seems to be delayed digestion.¹⁵ Malabsorption of carbohydrate may occur, but not to any great extent, and it does not appear to produce any long term problems.²²

Advice about other starchy carbohydrate foods is, however, less clear cut. The results of test meal studies suggest that the amount of dietary fibre is not the only determinant of the glycaemic response: the response to wholemeal (high fibre) and white (low fibre) bread is the same,¹⁵ and wheat in pasta elicits a lower response than wheat in bread.²³ Whole rice also results in a much flatter blood glucose curve than rice flour,²⁴ and glucose and insulin responses to whole apples or apples as purée differ,²⁵ which suggests that particle size and integrity of structure may be important. The results of laboratory studies of digestion with human digestive enzymes show that the rate of starch hydrolysis correlates well with the biological response to ingested food,²⁴ and such techniques might therefore be used to predict physiological response to food.²⁶ Does this mean that wholemeal bread and potato (already identified according to their glycaemic responses as less satisfactory than several other foods in the test meal experiment) should be eliminated from or at least discouraged in the diabetic diet, or that suitability of foods for the diabetic may in future be determined in the laboratory?

The answer to both questions must be no, at least in the light of present knowledge. Wholemeal bread and potatoes have formed a substantial part of several experimental diets which have over weeks or months shown an appreciable improvement in diabetic control.^{27 28} Furthermore, a recent

study compared diabetic control in insulin dependent diabetics in which the only dietary intervention consisted of replacing low fibre white bread with bread enriched with either wheat bran or guar gum: the mean blood glucose concentrations improved to an appreciable and similar extent.²⁹

What is the explanation for the difference between the test meal experiments and the longer term (and essentially more relevant) feeding studies? Apparently the benefits of a diet high in starchy carbohydrate are not necessarily conferred by a reduction in postprandial glycaemia. Indeed, in studies in which diurnal blood glucose profiles have been examined^{6-8 27 30} the main benefit seems to be a reduction of background blood glucose concentrations with little change in postprandial excursions giving a resultant overall decrease in average blood glucose concentrations, possibly via an effect on insulin sensitivity.³¹ Even starchy carbohydrates low in fibre appear to confer some benefit.³² Thus although the test meal and laboratory digestion experiments are of interest, they cannot replace longer term feeding studies in determining the best foods for diabetics. At present cooked dried leguminous seeds seem to be the best bet, with foods containing rice and pasta a good option too, but there seems no good reason for not including wholemeal bread, potato, and indeed other cereals or vegetables in a sensible diabetic diet.

What about sugar? Sucrose is not an essential food, and the overweight diabetic will need to restrict energy intake from sucrose as well as fat. Bantle and his colleagues concluded that in non-obese diabetics sucrose in controlled amounts as part of nutritionally balanced meals may enhance palatability and so increase overall dietary compliance without ill effect.²⁰ Given as part of a mixed meal, sucrose does not produce a greater glucose response than comparable amounts of potato or wheat starch. In this study, however, glucose concentrations were unacceptably high after all the test meals, which were low in dietary fibre. Mixed meal studies, such as those carried out by Bantle and colleagues, are undoubtedly preferable to tests of individual foods because of the likely effects of protein and fat on absorption and insulin secretion, but definitive advice concerning sugar can be given only once the longer term effects have been examined. The same reservation applies to the suggestion that fructose might be preferable to sucrose.

Finally, every commentary on dietary advice for diabetics should restate the most important principles: energy intake should be adjusted to maintain the ideal body weight and in the insulin dependent diabetic a regular meal pattern should be established to match the injected insulin.

J I MANN

University Lecturer in Social and
Community Medicine, and
Honorary Consultant Physician,
Radcliffe Infirmary,
Oxford OX2 6HE

- ¹ Mason AS, Boucher BJ. Diabetes mellitus. In: Scott RB, ed. *Price's textbook of medicine*. 12th ed. Oxford: Oxford University Press, 1978: 435-48.
- ² West KM. Diet therapy of diabetes: an analysis of failure. *Ann Intern Med* 1973;79:425-34.
- ³ Tunbridge R, Wetherill JH. Reliability and cost of diabetic diets. *Br Med J* 1970;ii:78-80.
- ⁴ Mann JI. Diet and diabetes. *Diabetologia* 1980;18:89-95.

- ⁵ Kiehm TG, Anderson JW, Ward K. Beneficial effects of a high carbohydrate, high fibre diet on hyperglycemic diabetic men. *Am J Clin Nutr* 1976;29:895-9.
- ⁶ Simpson RW, Mann JI, Eaton J, Carter RD, Hockaday TDR. High-carbohydrate diets and insulin-dependent diabetics. *Br Med J* 1979;iii: 523-5.
- ⁷ Simpson HCR, Simpson RW, Lousley S, et al. A high carbohydrate leguminous fibre diet improves all aspects of diabetic control. *Lancet* 1981;ii:1-5.
- ⁸ Kinmonth A-L, Angus RM, Jenkins PA, Smith MA, Baum JD. Whole foods and increased dietary fibre improve blood glucose control in diabetic children. *Arch Dis Child* 1982;57:187-94.
- ⁹ Reaven GM. How high the carbohydrate? *Diabetologia* 1980;19:409-13.
- ¹⁰ Nuttall FQ. Dietary recommendations for individuals with diabetes mellitus, 1979: summary of report from the Food and Nutrition Committee of the American Diabetes Association. *Am J Clin Nutr* 1980;33: 1311-2.
- ¹¹ Nutrition Sub-committee of the British Diabetic Association's Medical Advisory Committee. Dietary recommendations for the 1980s—a policy statement by the British Diabetic Association. *Human Nutrition: Applied Nutrition* 1982;36:378-86.
- ¹² Special Report Committee of the Canadian Diabetes Association. 1980 guidelines for the nutritional management of diabetes mellitus. *Journal of the Canadian Diabetes Association* 1981;42:110-8.
- ¹³ Winton R. Diet and diabetes. With care, diabetics can enjoy a normal, balanced diet. *Med J Aust* 1982;ii:245.
- ¹⁴ Huttunen JK, Aro A, Pelkonen R, Puomio M, Siltanen I, Akerblom HK. Dietary therapy in diabetes mellitus. *Acta Med Scand* 1982;211:469-75.
- ¹⁵ Jenkins DJA, Wolever TMS, Jenkins AL, et al. The glycaemic index of foods tested in diabetic patients. A new basis for carbohydrate exchange favouring the use of legumes. *Diabetologia* 1983;24:257-64.
- ¹⁶ Jenkins DJA, Goff DU, Leeds AR, et al. Unabsorbable carbohydrates and diabetes: decreased post-prandial hyperglycaemia. *Lancet* 1976;ii:172-4.
- ¹⁷ Crapo PA, Insel J, Sperling M, Kolterman OG. Comparison of serum glucose, insulin, and glucagon responses to different types of complex carbohydrate in noninsulin-dependent diabetic patients. *Am J Clin Nutr* 1981;34:184-90.
- ¹⁸ Crapo PA, Reaven G, Olefsky J. Postprandial plasma-glucose and insulin responses to different complex carbohydrates. *Diabetes* 1977;26:1178-83.
- ¹⁹ Jenkins DJA, Wolever TMS, Taylor RH, et al. Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* 1981; 34:362-6.
- ²⁰ Bantle JP, Laine DC, Castle GW, Thomas JW, Hoogwerf BJ, Goetz FC. Postprandial glucose and insulin responses to meals containing different carbohydrates in normal and diabetic subjects. *N Engl J Med* 1983;309: 7-12.
- ²¹ Bingwen L, Zhao Feng W, Wanzhen L, Rongjue Z. Effects of bean meal on serum cholesterol and triglycerides. *Chin Med J* 1981;94:455-8.
- ²² Jenkins DJA, Wolever TMS, Taylor RH, et al. Slow release dietary carbohydrate improves second meal tolerance. *Am J Clin Nutr* 1982;35: 1339-46.
- ²³ Jenkins DJA, Wolever TMS, Jenkins AL, Lee R, Wong GS, Josse R. Glycaemic response to wheat products: reduced response to pasta but no effect of fibre. *Diabetes Care* 1983;6:155-9.
- ²⁴ O'Dea KO, Snow P, Nestel P. Rate of starch hydrolysis in vitro as a predictor of metabolic responses to complex carbohydrate in vivo. *Am J Clin Nutr* 1981;34:1991-3.
- ²⁵ Haber GB, Heaton KW, Murphy D, Burroughs LF. Depletion and disruption of dietary fibre. Effects on satiety, plasma-glucose, and serum-insulin. *Lancet* 1977;ii:679-82.
- ²⁶ Jenkins DJA, Ghafari H, Wolever TMS, et al. Relationship between rate of digestion of foods and post-prandial glycaemia. *Diabetologia* 1982;22: 450-5.
- ²⁷ Simpson RW, Mann JI, Eaton J, Moore RA, Carter RD, Hockaday TDR. Improved glucose control in maturity-onset diabetes treated with high-carbohydrate-modified fat diet. *Br Med J* 1979;ii:1753-6.
- ²⁸ Anderson JW, Ward K. Long-term effects of high-carbohydrate, high-fibre diets on glucose and lipid metabolism. A preliminary report on patients with diabetes. *Diabetes Care* 1979;1:77-82.
- ²⁹ Vaaler S, Hanssen KF, Dahl-Jorgensen L, et al. Improvement in long-term diabetic control after high fibre (bran and guar) diets. *Diabetologia* 1983; 25:200.
- ³⁰ Aro A, Uusitupa M, Voutilainen E, Hersio K, Korhonen T, Siitonen O. Improved diabetic control and hypocholesterolaemia with guar gum in type 2 insulin-independent diabetes. *Diabetologia* 1981;21:29-33.
- ³¹ Ward GM, Simpson RW, Simpson HCR, Naylor BA, Mann JI, Turner RC. Insulin receptor binding increased by high carbohydrate low fat diet in non-insulin dependent diabetics. *Eur J Clin Invest* 1982;12:93-6.
- ³² Simpson HCR, Carter RD, Lousley S, Mann JI. Digestible carbohydrate—an independent effect on diabetic control in type 2 (non-insulin-dependent) diabetic patients? *Diabetologia* 1982;23:235-9.