Clinical Algorithms

Syncope

ANTHONY D ORMEROD

Transient loss of consciousness may be due to syncope, epilepsy, cardiac arrhythmia, and, rarely, transient cerebral ischaemic attack. It is important not to attribute syncope to cerebrovascular disease when the true cause is a transient disturbance of cardiac rhythm which is amenable to therapy. The differences between epilepsy and syncope will usually be apparent from a carefully taken history and whenever possible from an eye witness account.

Syncope is sudden, brief loss of consciousness due to transient impairment of cerebral circulation, from whatever cause, usually occurring in the absence of organic brain disease or cerebrovascular disease. Typically the onset is sudden or develops over a few seconds with blurred vision, dizziness, cold extremities, and perspiration. These features may give enough warning for the sufferer to take evasive action such as sitting down; otherwise the patient becomes limp and sinks to the ground with sighing respirations, pallor, and sweating. If venous return is impaired by the patient being propped up, or if the attack is prolonged, muscle twitching and convulsions may occur. These are rare but may confuse the diagnosis. The patient with syncope may complain of funny spells or dizzy turns in which consciousness is altered but not lost. The setting in which the attack occurs may help to distinguish the simple faint from epilepsy. For example, faints commonly occur in warm weather, when standing for prolonged periods, or after venepuncture.

In epilepsy the patient is usually rigid rather than limp; the onset is more abrupt, with injury and sometimes incontinence; and recovery is slow with confusion and drowsiness. The presence of an aura or perceptive or motor disturbances may all point to epilepsy. Fits and faints can both be precipitated by emotion and can both be related to the menstrual cycle. In transient ischaemic attacks consciousness is not usually lost unless the basilar artery territory is affected or there is global ischaemia. These patients will usually have focal neurological symptoms such as vertigo, visual disturbances, or ataxia.

Investigations

History and examination are the most important factors in determining the cause of syncope. In those patients for whom a diagnosis is not apparent on initial assessment further investigation is often fruitless and needs to be carefully directed to ensure the exclusion of sinister causes.¹ Most patients have a good prognosis but those over 70 years and those with a cardiac cause are at a much

higher risk of death or major morbidity.² In the young patient with a history suggestive of a simple faint no further investigation is required. In the older patient this diagnosis should be made with great caution as cardiac dysrhythmia is more likely. In the elderly drugs are frequently to blame and many factors may operate together.

In all but the common faint an electrocardiograph is mandatory. During the electrocardiogram carotid sinus massage will reveal carotid sinus hypersensitivity and may uncover an underlying conduction defect. This is best done with the patient lying at 45° for five seconds only, on one side only, and with facilities for resuscitation at hand.³ A full blood count may help to exclude anaemia or blood loss; serum electrolyte concentrations may indicate dehydration, Addison's disease, or hypocalcaemic and uraemic causes of convulsions; chest radiographs might show evidence of intrinsic cardiac disease or pulmonary embolism.

In a review of 121 patients with syncope in whom the diagnosis was undetermined by initial assessment a definitive cause was found in only 13.⁴ By far the single most useful test was 24 hour continuous ambulatory electrocardiography, although careful interpretation is required to decide whether the abnormalities found explain the patient's symptoms. In the population studied computed tomography, radionuclide scanning, skull radiography, lumbar puncture, and glucose tolerance tests did not aid the diagnosis in any of the patients. Echocardiography, cardiac catheterisation, and electrophysiological studies were of intermediate value. Recent evidence suggests that electrophysiological studies are the most sensitive indicator of cardiac arrhythmia. In 149 patients with negative results from exhaustive investigations, including 24 hour continuous ambulatory electrocardiographic monitoring, 63% showed positive results, and with appropriate therapy 87% of these patients became free of symptoms⁵; the technique reveals a high incidence of unsuspected paroxysmal ventricular tachycardia not detected by non-invasive methods. It is, however, more expensive and invasive than 24 hour electrocardiographic monitoring and requires a skilled operator. There may be a place for empirical permanent pacing of patients with a suspected conduction abnormality not confirmed by investigation.⁶

References

- ¹ Critchley EMR, Wright JS. Evaluation of syncope. *Br Med J* 1983;286:500-1.
- ² Day SC, Cook EF, Funkenstein H, Goldman L. Evaluation and outcome of emergency room patients with transient loss of consciousness. *Am J Med* 1982;73:15-23.
- ³ Davies AB, Stephens MR, Davies AG. Carotid sinus hypersensitivity in patients presenting with syncope. *Br Heart J* 1979;42:583-6.
- ⁴ Kapoor WN, Karpf M, Maher Y, Miller RA, Levey GS. Syncope of unknown origin. The need for a more cost-effective approach to its diagnostic evaluation. *JAMA* 1982;247:2687-91.
- ⁵ Camm AJ, Levy AM. Evaluation of syncope. *Br Med J* 1983;286:895.
- ⁶ Gulamhusein S, Naccarelli GV, Ko PT, et al. Value and limitations of clinical electrophysiologic study in assessment of patients with unexplained syncope. *Am J Med* 1982;73:700-5.

Syncope 1

Syncope 2