Clinical Algorithms

Memory loss

ROBERT A WOOD

Memory loss covers a wide range of medicine, neurology, and psychiatry. This account touches on various conditions in which memory loss is only part of the picture, but I have emphasised pure amnesic states or others in which amnesic symptoms can occur early in the course of the illness. The importance lies in the early recognition of the treatable and in preventing deterioration where the cause is not inexorably progressive. The main subdivision is between amnesia of sudden onset, in which the outcome is usually favourable, and slow onset amnesia—often with global dementia—where it is not.

Memory

Defects of memory are either in the registration of new experience (antegrade) or the recall of past experience (retrograde). In gross dementia the patient may lose the ability to repeat a series of numbers he has just heard. This immediate type of defect is also seen in dysphasia, or whenever attention or concentration is distracted—as in depression or acute confusional states. Short term memory deficit is tested by inviting the recall of a few numbers and two or three names after five minutes of other conversation. The limbic system, including the mamillary bodies and hippocampus, must be intact for new experience to be registered and recent events recollected. In Korsakow's amnesia, where the main damage is in the mamillary bodies, antegrade amnesia—learning ability—is permanently damaged and several years of experience are lost. After head injury an antegrade defect is noted but this is temporary. The retrograde amnesia shrinks until minutes or seconds before the accident. Deep memory, including childhood events, can be retained and recalled when limbic structures no longer function. The long term memory is tested by asking for details of the patient and his family and about important people or events. Moments of great personal pleasure or poignancy can be recalled most clearly.

Alcohol related amnesia

When the blood alcohol concentration is very high the memory may be blacked out. This is impossible in inexperienced drinkers, who will pass out before they become amnesic. Many episodes of acute alcoholic amnesia lead on to delirium tremens, the withdrawal syndrome of alcohol tolerance, which is characterised by motor excitation, tremor, and hallucination; it clears within two to three days. Treatment is with chlormethiazole or haloperidol to reduce the excitation and with parenteral vitamin B complex to prevent the development of Wernicke's encephalopathy. Alcohol and its subsequent withdrawal are important causes of seizure in epileptics. Alcohol and head injury are often associated, and intracerebral clot is hard to diagnose in intoxicated or delirious alcoholics.

When prolonged drinking gives way to confusion, the presence of ataxia and diplopia or disorders of gaze indicate the development of Wernicke's encephalopathy. This confusional state is caused by thiamine deficiency. It should be suspected if the confusion of withdrawal does not clear in the usual time. If treatment with parenteral vitamin B complex is omitted or delayed a dense and permanent Korsakow's amnesia will become apparent when the confusion clears. No treatment is worth while at this stage, and fewer than a tenth of patients show any improvement in the long term. Institutional care is the usual end result.

Chronic alcoholic dementia is more common than Korsakow's amnesia. The early features are occasionally amnesic, but generally judgment and social projection are impaired first. It is a long term toxic effect, usually in the older age groups, and the older the patient the poorer the prospect for improvement. Computed tomography usually shows cortical atrophy and ventricular enlargement; these may improve with abstinence, but the dementia itself seldom disappears entirely, and improvement is maximal after a month or two. Abstinence is the only beneficial treatment. This is the fourth most common progressive dementia.

Malingering and hysteria

Hysterical amnesia is rare. The patient may claim to remember nothing or forget only specific personal events. Feigned amnesia is common, often to avoid arrest or conviction: the subject who "disappears" after a car crash and turns up uninjured hours later has been hiding with the purpose of metabolising alcohol. It should be distinguished from the brief, complex, automatic activity seen in some temporal lobe seizures. The conscious non-dysphasic patient who cannot give his name does not have an organic diagnosis. Hysteria resolves in time.

Epilepsy

After a major seizure consciousness is recovered quickly but antegrade amnesia lasts for up to an hour. The patient may also appear muddled. Epilepsy is high on the list of possibilities in a patient who has been briefly amnesic and who has unaccountable injuries. In the known epileptic alcohol and failure to comply with prescribed drugs are common provocative factors. Epilepsy may occur in most of the organic syndromes in which amnesia occurs and both it and the medication used to control it may give a false impression of the severity of amnesia.

Post-traumatic brain syndrome

After head injury a classical amnesic syndrome is seen with severity and duration relating to the severity of injury. Most patients recover fully, though it may be days or weeks before short term memory is restored. The retrograde amnesia, initially quite long in many patients, shrinks progressively until it is just seconds or minutes before the injury; this contrasts with the lengthy retrograde amnesia in Korsakow's syndrome.

Memory loss

The importance of the amnesia is that it confirms brain injury. Skull radiographs are useful, although a fractured skull is no predictor of the severity of the neurological injury. Computed tomography has revolutionised the management of head injury. Unconscious patients need specialised supervision in units where the brain can be decompressed and clot evacuated. Dexamethasone may reduce the duration of the organic brain syndrome after head injury or neurosurgery. Opiates and sedatives are unsuitable because of possible effects on eye signs and consciousness.

Cerebrovascular syndromes

Pure amnesia may be due to cerebrovascular disease. Transient global amnesia is common; memory for recent events is lost, usually for a few hours during which the patient can carry out various routine activities but has to be reminded that he has done so. Embolism or transient ischaemia in the basilar distribution is the likely cause; this may lead to temporary left hippocampal dysfunction. Full recovery occurs without treatment.

Rarely a cerebral infarct in the left hippocampus may produce a pure amnesia with short term memory loss and the usual retrograde and antegrade components. Prognosis is reasonable provided the other hippocampus is intact, but maximum improvement may take up to three months. Some events in the basilar distribution produce amnesia associated with florid confabulation and sleep disturbance. Sometimes a variety of visual disturbances are associated, such as cortical blindness, visual disorientation, field defect, and defects of gaze (often upwards). Amnesia with visual defects occurs after resuscitation from cardiac arrest or bypass surgery. This condition, variously known as borderzone or watershed ischaemia, refers to the fringes of the basilar circulation, and has a variable prognosis; there is no treatment. Temporary amnesia may follow vertebral angiography. Any established lesion should be assessed by computed tomography so that prognosis can be given.

Multiple cerebral infarction can lead to dementia. The syndrome is unpredictable, but it can usually be distinguished from Alzheimer type dementia by the stepwise deterioration, often with relative preservation of insight and personality. Lability of mood and pseudobulbar features are pointers, as are sudden onset pyramidal tract or parietal features. The commonest cause is hypertension, but any vascular disease affecting the left heart, aorta, or carotid vessels may be a cause. The treatment is the identification and removal of the cause; it is not enough to hope that patients will not have recurrent cerebrovascular events. The most effective prevention is good antihypertensive treatment, avoidance of smoking, and early surgery in carotid atheroma.

Depression

Mild depression may present with memory loss, and severe depression in the elderly may masquerade as dementia. The problem is that poor memory and concentration may be more obvious than the abnormal mood, depressive thought content, and sleep disorder. The depressed patient tends to be uninterested and inattentive but does not confabulate. Up to 10% of elderly patients thought to have dementia may make an excellent recovery with antidepressant treatment. Such opportunity must not be missed, and the presence of some cortical atrophy on tomography should not preclude a trial of such treatment.

Some patients with early or established dementia are depressed. Insight disappears early in Alzheimer's disease but may be preserved in multi-infarct dementia. Some patients with multi-infarct dementia recover much of their intellectual function when given an antidepressant.

Cerebral tumour

Rapid progression of focal signs, the onset of epilepsy, and papilloedema support the diagnosis of cerebral tumour. Slowly growing and suitably sited tumours may cause slowly progressive dementia,

but headache is almost invariably present. Rarely a pure amnesic syndrome is seen in primary or secondary cerebral malignancy. Pure amnesia can develop as a non-metastatic manifestation of bronchial carcinoma, in association with encephalitic features in the cerebrospinal fluid. Tumour is diagnosed from focal electroencephalographic features, computed tomography, and angiography.

Pseudotumour cerebri

Amnesia or pseudodementia is rarely seen in pseudotumour cerebri (or benign intracranial hypertension), a diagnosis made when serious causes of raised intracranial pressure have been excluded. There are many causes, but most patients are young women. When headache is severe and amnesia present a shunt will generally be required to allow some of the excessive cerebrospinal fluid to drain into a jugular vein or the peritoneal cavity. Prognosis is excellent.

Subdural haematoma

Half of the patients with chronic subdural haematoma do not recall the injury, which may have been mild. Headache and mental change (including amnesia) are the usual complaints but both may be missed when there is confusion. Focal signs and epilepsy are pointers. Very gradual tentorial herniation can lead to unobtrusive features such as minimal ptosis and paresis of upward gaze. Subdural haematoma is diagnosed by computed tomography, may be bilateral, and should be considered in alcoholics and epileptics. Chronic subdural haematoma causing pressure symptoms requires urgent surgery.

Boxer's dementia

The tragic syndrome of dementia with ataxia, dysarthria, and parkinsonian features is rare but is a compelling argument against boxing. There is speculation that forms of dancing involving violent head movement may have the same long term result.

Occult hydrocephalus

Though rare, occult hydrocephalus is important because early treatment produces good results. The clue is the history of meningitis, intracranial haemorrhage, or head injury. The features are dementia with a very early amnesia, incontinence, and apraxia of gait. This is a slow, unstable, wide based, fixed footed gait with correction so poor that falling is common. There may be no motor or sensory features on examining the legs.

Neurosyphilis

Although rare, neurosyphilis has a good prognosis if treated early. The common presentation is simple dementia, not the florid grandiose confabulation that is so well known. The early features may be irritability and personality change as in some cerebral tumours. Headache is not unusual. As amnesia is often late, the differential diagnosis is from other causes of personality deterioration. Quite often there are Argyll-Robertson pupils. Diagnosis is made by examining blood or cerebrospinal fluid. Positive haemagglutination and fluorescent antibody test results in blood do not confirm active disease, but in cerebrospinal fluid a positive Venereal Diseases Research Laboratory test and raised protein concentration and lymphocytosis are confirmatory. Treatment is with penicillin, initially under corticosteroid cover.

Disorders of movement

The early signs of Huntington's chorea are irritability and anxiety-depression, with amnesia and choreiform movement later.

Most patients retain insight for some time. In both Parkinson's disease and Wilson's disease intellectual deterioration is late and the underlying disease will not be suspected because of amnesia. The dementia of advanced Parkinson's disease, although sometimes an indication of cerebral disease, can be related to drug treatment, especially excessive levodopa.

General medical disorders

Hypothyroidism, vitamin B₁₂ deficiency, and hypercalcaemia may all present as dementia with prominent amnesic features and in the absence of other more usual symptoms. Routine screening in dementia should always include estimation of thyroid stimulating hormone, mean cell volume, and serum calcium concentration.

Drug treatment

A full drug history must be obtained in every patient who has memory defect. Chronic dependence on sedatives, benzodiazepines, or phenothiazines and certain anticonvulsants may lead to a chronic pseudodementia in which memory failure is prominent. Digoxin and corticosteroids are less well known causes. Sulphonylureas and insulin, while they may obviously cause acute hypoglycaemia, can more unobtrusively lead to chronic hypoglycaemia, to which the patient becomes partly acclimatised. The pseudodementia usually recovers when the cause is removed; a few patients possibly sustain permanent intellectual impairment.

A wide variety of drugs cause toxic confusional states, either alone or with other causes. The list includes sedatives, anticonvulsants, antidepressants, antiparkinsonian medication, antihistamines, corticosteroids, digoxin, atropine, opiates, cycloserine, isoniazid, hypoglycaemics, and hypotensive agents. The treatment consists of reducing or withdrawing the offending drug, replacing it if necessary, correcting other contributory factors, and giving chlormethiazole or haloperidol to control overactivity or other florid features.

In younger patients solvent abuse can produce an apathetic pseudodementia. The florid hallucination of the bad lysergic acid trip is well known in casualty departments.

Toxic and metabolic confusional syndrome

In the older age group the responsibility is to identify the underlying disease in confusional syndromes, commonly chest or urine

infection, myocardial infarction, or cerebral infarction. Generally the treatment is that of the cause, but with medication to control the more florid manifestations. Sedatives should be avoided in chronic lung disease, severe heart failure, and acute intermittent porphyria, where the disease may be intensified. Cranial arteritis is not easy to diagnose; if suspected corticosteroid treatment should be urgently considered. Drugs are an important cause.

Encephalitic illnesses of any severity produce striking memory impairment although mingled with other features. The recovery may not be complete and the antegrade amnesia may persist longer than after head injury, and with a deeper retrograde amnesia.

Alzheimer's disease

Alzheimer's disease is responsible for up to 80% of cases of dementia. Cases occurring in people aged under 65 are described as presenile. There may be early insight, with some anxiety or depression, but this shortly disappears. A poor and deteriorating memory may be the presenting problem. The other main features are deterioration of intellect, personality and behaviour change, and focal neurological symptoms or signs. These include subtle dysphasic errors, along with apraxia and agnosia, which are recognised from an inability to dress, find the way around a familiar environment, or carry out simple tasks with everyday objects. Focal and major epilepsy is seen later in the illness along with primitive reflexes such as sucking and tonic grasp and parkinsonian features. The progress is inexorable, and death is from inanition and pneumonia.

The diagnosis is strongly suggested by dementia associated with mild organic signs but with no evidence of intracranial pressure increase or headache. Computed tomography confirms diffuse cerebral atrophy with ventricular enlargement, and brain biopsy (seldom carried out) shows fibrillary tangles and granulovacuolar degeneration. Despite biochemical evidence of certain deficiencies in the brain, there is no treatment.

Benign forgetfulness

This is the "condition" in which many middle aged and older patients are inclined to forget the names of persons or places—often temporarily. Mood and affect are normal, and formal tests of memory, intellect, and personality show nothing abnormal. Fatigue and worry may contribute. This condition is not a precursor of Alzheimer's disease and may be the commonest "amnesia" presenting to the medical profession. Reassurance is required.

MATERIA NON MEDICA

Not enough books—too many bibles

A recent *Materia non Medica* article celebrated first and second class travel on the Nairobi-Mombasa railway. It certainly is an excellent train, but perhaps a more memorable experience awaits the traveller who passes by the 10 or 12 upper class carriages and travels in one of the two third class carriages. At least half of the passengers travel in this part of the train. You are not allowed to pay £4 for a four course meal in the upper class dining car; instead you have two options: fill yourself up with maize and beans for fifty pence in the snack bar, or buy samosas. These are pieces of curried meat or vegetable wrapped in pastry. The railway through the outskirts of Nairobi is lined with people waiting to sell them through the windows of the slow moving train. I recommend the samosas—but make sure you have the correct change.

One benefit of third class travel is the people you meet. My neighbours were two brothers returning home for their father's funeral. One of them told me stories of how as a child he had guarded his father's cows from wild animals. The secret of killing a lion, he told me, is to delay your spear thrust until the lion is upon you. He was a member of the same tribe as the

president, the Kalenjin. His ancestors had certainly fought lions to protect their cattle but I had some doubt about the claims of this grey suited man.

Paradoxically, I found reading a book a great conversation starter. Both men cast envious eyes on my book—Paul Theroux's *Great Railway Bazaar*, the story of a train journey from London to Japan via India, returning through Russia. They bemoaned to me the difficulty of getting books to read—both were perusing bibles, which they said were the only books available. This is not because of any censorship or restrictions but simply because the literacy rate in Kenya is growing so fast that there are not enough other books to go round. One brother asked me to send him the *Oxford Dictionary of Idiomatic English*.

It saddened me to remember that the prize for best pupil nurse at our hospital never varied—it was always a bible. Mission schools require every pupil to have a bible. Every hotel room contains a bible. Kenya is not a heathen country. I wonder how long it will be before the churches feel they can stop this book propaganda? Must the bible be given to everyone like Chairman Mao's little red book? Is there no alternative prize for good pupils?—JIM THORNTON, Chogoria Hospital, Kenya.