Paul Griffiths for advice on computing, and the doctors, staff, and particularly the patients of the study practices, without whom the project could not have been conducted.

Appendix

Suggestions concerning advice to be given.

As general practitioners and their patients show wide individual variation, it is not surprising that exactly the same anti-smoking message can be delivered to each patient who smokes.

It is to be hoped, however, that each doctor will be able to discuss briefly most or all of the following points with each patient.

- (1) *Risk*—Smoking is known to cause a large number of diseases which together account for much of the sickness and premature death in the community.
- (2) *Cost*—The habit costs the average smoker well over £200 a year. Unborn children and young children suffer if exposed to tobacco smoke.
- (3) *Group up*—All smokers should give up the habit if possible. Eight million Britons have already given up.
- (4) *Problem of giving up*—These last two for three weeks for most people. Any weight gain tends to be temporary.
- (5) *"Let's hazard our smoking"*—If you cannot give up smoke fewer cigarettes, take fewer puffs, inhale them less deeply, and leave longer butts. But try to give up completely.

"We were a little optimistic, but there is a stable number of patients who find it surprisingly easy to give up smoking once they make a firm decision to do so."

Diary of Urban Marks: 1880-1949

After Jarvis, an old man from Gloucester was obtained. He carried on the work well but as he suffered from urinary trouble he smelled rather badly. He used to apologise for this and like Groves he took meals by himself. He carried on until during the last few months of my absence Soady, whom I had with me in Mesopotamia, took charge. This had been arranged between Soady and myself and he was acting for me when I returned. During these six months he worked hard and was very popular. He pulled the practice together and when I arrived home everything was working smoothly and normally.

Just a few words in connection with Arthur Soady. He was an Irishman from Dublin and with his father had been accountants for a big railway in Ireland. Soon after he qualified Arthur felt the call of medicine and having taken a medical degree joined the army for 12 months, as all Irishmen had that option. I did not see him at Blackpool but I was allotted a cabin with him on the boat on which we went out to Mesopotamia. He had a charming manner with him and curiously enough had not a good word for the Irish. For the rest, he had a curiously shaped hooked nose and was about 5 ft 5 in in height. He did not possess a brogue.

On my return to civil practice I tried to persuade him to come into partnership with me and open out in Port Tennant on the other side of the Swansea River, where I had a large number of patients owing to my connection with the Baldwin Works. Unfortunately, for myself and for him, he had an aunt in Dublin from whom he had great expectations. Arthur managed all her affairs and she insisted that if he settled down he must be somewhere near her. In vain he pointed out to her that he could be in Dublin in a few hours from here. It was useless, and Arthur for some time did a series of locum tenenships all over the country. He went to the West Riding of Yorkshire to take over a practice for a man who was dying from the effects of

References

1. Mauser JS, Mauser B, Rial WY. The influence of a physician on the smoking of his patients. *Am J Public Health* 1968;58:46-53.
2. Pincheffe G, Wright HB. Smoking habits of business executives: doctor variation in reducing cigarette consumption. *Practitioner* 1972;205:209-12.
3. Porter AMW, McCullough DM. Counseling against cigarette smoking. *Practitioner* 1982;229:9-9.
4. Handel S. Change in smoking habits in a general practice. *Postgrad Med J* 1973;49:679-81.
5. Burnum JE. Outlook for treating patients with self-destructive habits. *Ann Intern Med* 1981;94:387-93.
6. Russell MHE, Wilson C, Taylor C, Baker CD. Effect of general practitioners' advice against smoking. *Br Med J* 1979;1:231-5.
7. Stewart PJ, Rosser WW. The impact of routine advice on smoking cessation from family physicians. *Can Med Assoc J* 1982;126:1091-4.
8. Sheikh K, Mattingly S. Investigating non-response bias in mail surveys. *J Epidemiol Community Health* 1981;35:204-5.
9. Jamrozik K, Fowler G. Anti-smoking education in Oxfordshire general practices. *J R Coll Gen Pract* 1982;32:140-53.
10. Laranjeira J, Guha HB, Van Vunakis H. Nicotine and its metabolites: radioimmunoassays for nicotine and cotinine. *Biochemistry* 1981;20:255-60.
11. Office of Population Censuses and Surveys. *OPCS Monitor*. General Household Survey GHS 80:1. London: OPCS, 1981:11-5.
12. Fowler G. Practising prevention: smoking. *Br Med J* 1982;284:1306-8.
13. Siller RW, Wilson MB, Makin RE, Ball KP. Descriptions among smokers. *Br Med J* 1979;1:1185-9.
14. Wilson R, Hughes J, Roland J. Verification of smoking history in patients after infarction using urinary nicotine and cotinine measurements. *Br Med J* 1979;1:1026-8.
15. Burt A, Hingworth D, Shaw T, Thornley P, White P, Turner R. Stopping smoking after myocardial infarction. *Lancet* 1974;1:946.

(Accepted 14 February 1984)

alcohol. After this doctor's death he stayed on until the practice had been disposed of and the next thing I heard was that he had married the widow. They naturally could not stay in the neighbourhood and he settled down in London in one of the east end suburbs. But the London fogs played havoc with his chest. He had never been in good health since the time he arrived in Mesopotamia. On his arrival there he had been posted to the front line well up the country. As soon as he reached there he went down with dysentery and was invalided to Baghdad. There he developed asthma and finally in the October of 1917 was sent to No 2 BGH. I remember the colonel asking me if I would take charge of an invalid officer who was fit for next to nothing. I was greatly surprised to find that my charge was my old friend, Soady. I put him in charge of the dysentery wards. When I took over the accounts of the mess at the end of 1917 it was largely due to the accountancy of Soady that we put the mess on a sound footing. He remained with us until March of 1918 when he left Mesopotamia for home. It had been arranged between us that he should take over my practice until I returned home. When he arrived in England his old trouble asserted itself and he went down with pneumonia and it was not until late in 1918 that he was able to take charge of my practice.

He sold his first practice in London at a large profit and then went to Skeneburg but the cold winds of the winter did not suit him and he had to retire from the place. Just about this time his aunt died and left him over £10,000. He then decided to take up a seafaring life and joined the P & O Company as a ship's surgeon. Meanwhile, his wife went to live in London in a private house. For some years he went backwards and forwards to Australia in the boats and on several occasions when the ship put into Swansea for cargo I had an opportunity of seeing him. He was always very delighted to come to Swansea and was never tired of telling people how I saved his life. Finally he tired of the sea and with his wife settled down into a country practice.

Fig. 2—'Slapped cheek' disease

10. 3.—From "Rice Pudding" in *When We Were Very Young* by A. A. Milne, illustrated by H. Shepard, published by Methuen Children's Books Ltd. and reproduced with the permission of Curtis Brown Ltd.

captures something of this in his poem "Rice Pudding" when Mary Jane is again presented with rice pudding, the resulting tantrum cannot be contained within the family. So the family doctor is called upon to "cure" Mary Jane (fig. 3), whose family problem may not be appreciated by its grown up members. In adults the underlying problem may be marital disharmony or alcoholism in the spouse, or the stress induced by caring for an elderly parent. The patient may not necessarily be fully aware of the nature of the problem, and there is little conscious attempt to conceal the state of affairs.

The cover story

Peggy E, a 16 year old schoolgirl, presented with a complaint of "catarrh". It was only after this had been dealt with that the true agenda was revealed—she wanted to go on the contraceptive pill. Such "collateral problems" may come to the surface as a parting shot—the "while I'm here" syndrome, and after the patient has had an opportunity to assess the doctor and judge his likely response to a second, more "serious" offer. The experienced general practitioner learns to anticipate and facilitate the disclosure by such a phrase as, "Is there anything else we should discuss?"

The frequent attender

Most practices contain a few patients who generate a disproportionately large volume of work for the doctor—well illustrated by Dickinson and Farrow.¹ Such patients are characterised by the high frequency of contact, and by the variety of their minor but baffling symptoms. The frustration they engender in the doctor resides not simply in the high demand but in the fact that they persistently reopen issues which the doctor has attempted to close. Freeing has coined the term "recidivist" for such a patient, recognising the pejorative nature of the term, but unable to suggest a better one.²

Minor medicine—a classification

So what at first appears to be a simple triviality is sometimes an elaborate affair. The proposed classification (fig. 4) is not intended to be used to stick labels on people; rather it is an attempt to clarify thinking on a difficult subject.

Fig. 4—Classification of minor presentations in general practice

Recognition—and triage

In addition to applying common sense epidemiology along the lines mentioned above the doctor deploys many skills in dealing with this numerically important component of his work. They include pattern recognition, a sound knowledge of the natural history of diseases (both major and minor), the use of time as a diagnostic tool, and, above all, a knowledge of his patients and their illness behaviour. With experience, general practitioners acquire a kind of rank order of seriousness to apply to any given presentation: thus Hodgkin records diarrhoea as 100 trivial to 1 serious, and haematemesis as 1 trivial to 2 serious causes.³ Laboratory investigations usually have little to offer, with the exception of the erythrocyte sedimentation rate; the finding of an otherwise unexplained raised value calls for a further assessment of the apparent minor presentation.

Much of minor medicine is not amenable to diagnosis in the sense of applying labels that indicate clear cut pathology and imply a prognosis. When diagnosis is not clear at an extreme the doctor's natural response might be "I don't know exactly what this is, but it's not serious"; at the other, "This patient requires prompt admission to hospital." Perhaps a more suitable term is "triage"—hitherto reserved for the sorting of casualties of war or major disaster. *Blackburn's Gould Medical*

Rethinking Established Dogma

Trivia, triage, and treatment

JAMES D E KNOX

Some years ago senior medical students at Dundee were invited to define a general practitioner and to describe his work. From among the replies it was possible to define a common theme, best summarised by one response, which read, "The general practitioner is a triavologist: a signpost to hospital."

From the patients' point of view their presentations are seldom, if ever, trivial. In comparison, however, with say, Mr A's acute myocardial infarction, Tommy B's sunburn pales to insignificance. By the same token, Ina C's presentation of tiredness and churning in the stomach—coupled with a request for a tonic—could be regarded as humdrum, an everyday occurrence in general practice. Ina C's request was

in fact, the presentation of what some months later was seen to be overt exophthalmic goitre. Thus, perhaps the term "trivial" ("litter discarded at the crossroads of life") needs to be replaced by another, less emotive phrase such as "minor medicine." By this I mean "a perceived deviation from the normal, associated with temporary interference with usual functions or a greater incapacity of short or several days' duration." Such a definition still begs a number of questions. Ezra D, aged 75 years, attended the doctor to obtain his signature to enable him to claim from a private pension fund. The doctor noted a large swelling of the right side of his face—almost certainly a mixed tumour of parotid. Ezra declined any treatment for a condition he had had for over 40 years, and which, he claimed, had never bothered him. The definition of "minor medicine" used in this paper thus requires further qualification: the deviation as perceived by the patient and presented to the doctor.

Epidemiology of minor medicine

By such a definition I classified as "minor" approximately 90% of 649 contacts, initiated by patients and seen by me at the Dundee Medical School teaching practice from August 1982 to July 1983. Applying the same definition selectively to 439 children under 16 years of age in the same practice, the proportion of minor medicine was 80%.

Jacob applied this approach, somewhat arbitrarily, to the diagnostic labels attached by 24 family doctors to 6017 first contacts, in part of a larger study of the work of Dundee general

practitioners.¹ Analysis of the distribution by age group and place of contact showed that the proportion of "non-serious" disease decreased with age and home contact compared with consulting room contact (fig. 1).

Fig. 1—Non-serious first contact as a percentage of all contacts (6017 consultations). Source: Dundee Health Centre Study, Phase 5, March 1979, total of 16 291 contacts.

Content of minor medicine

Minor medicine covers a wide spectrum and includes many fascinating disease states. Some of these are only now becoming more clearly defined—for example, "slapped cheek" disease (fig. 2) in which human parvovirus infection has only recently been incriminated as an important aetiological agent.² Such examples relate to relatively straightforward conditions, of interest in themselves, and, of course, far from trivial to the sufferer. Most minor presentations in general practice are of this "single problem" type. In a small proportion, however, the patient's condition is less easily defined and may be only part of more complex situations. It is possible to classify such presentations into subgroups, depending on the doctor's further knowledge of the patient and his setting. Such knowledge is often acquired over time and depends to some extent on the doctor's willingness and ability to perceive the person and the family behind the presentation.

"Mary Jane" phenomenon

In a few presentations the apparent patient, usually a child, is in reality merely an index of a disturbed family. A A Milne

Dickinson, however, has extended the definition to include the sick as well as the wounded.³ The general practitioner in the National Health Service occupies a key position as a determinant of hospital use—an increasingly expensive prescription. It is important to the nation's economy that this resource is appropriately deployed. In addition to economical and efficient use of expensive hospital resources, appropriate triage secures the effective use of scarce specialist skills, ensuring that they are deployed to best advantage. Visitors from the United Kingdom to foreign countries lacking a strong primary care system are often struck by the extent to which the work of highly trained specialists is "diluted" by the inclusion of inappropriate case material.

Treatment

At the level of the individual patient recognition and triage of minor medicine usually have to be complemented by treatment. In this paper it would be inappropriate to attempt to give details of treatment for each condition ranging from acne to zymotic diseases. Some generalisations may be made, however.

Single minor condition

The "seriousness" of a condition is often inversely proportional to the meaning the symptom holds for the sufferer: the patient rolling in agony with colic is concerned with the immediate relief of his pain, while the patient with vague epigastric unease may well be living in fear of having cancer of the stomach. It follows that management of minor medicine has to take psychosocial considerations fully into account. Sometimes the patient feels guilty about "bothering the doctor," and a delicate patient has to be tried to save the patient's face and at the same time, when appropriate, encourage self care in the future.

One of the greatest therapeutic dilemmas facing the general practitioner dealing with minor illness, especially upper respiratory tract infections, is an absence of a rational basis for prescribing antibiotics.⁴ Perhaps an acknowledgment of this fact and its wide acceptance by the medical profession would lead to a reduction in what seems to be widespread overuse of antibiotics.

A proportion of single minor problems require the skills of the nurse rather than the doctor. More could—some would say "should"—be delegated to such professional staff, provided they are suitably trained for these tasks, and nursing education and training need to be more fully geared for this to happen more widely.

Mary Jane

The underlying problem with "Mary Jane" is likely, to be complex, requiring the assessment and management skills of other health professionals such as the health visitor, social worker, or marriage counsellor. The need here is for greater teamwork, including better communication.

Collateral

The doctor sometimes merely needs to listen and to use counselling skills. The time spent in dealing with the "while

I'm here" presentation is, in my experience, usually well worth while, despite the irritation it may cause the doctor at the time.

Recidivist

Such people resemble psychotic patients because prolonged contact and attempts to communicate rationally are commonly unrewarding. I usually give myself a time limit on such consultations and will occasionally seek the help of consultant colleagues (suitably briefed) to secure temporary respite from the recidivist's importunities. This also allows for constant vigilance for the early presentation of major disease.

Minor presentations of major illness

It is usually necessary to consult a specialist as early as possible. This decision to refer sometimes requires positive effort: after all, some 90% of contacts do not call for such action, and often it has to be taken in the face of slender evidence. The patient's well being must take pride of place over the doctor's sensitivity concerning his reputation as a diagnostician with consultant colleagues. Among the peculiar intellectual satisfactions of general practice is the justification of such referral action in the light of subsequent events.

Triage and medical education

As long as medical education continues to be based almost exclusively on the teaching hospital, the undergraduate has little opportunity to acquire the knowledge, skills, and attitudes necessary for efficient, effective, and efficacious recognition and management of a very large proportion of health problems of contemporary society. And many of these problems remain underresearched. University departments of general practice are striving to correct this imbalance, but can we be sure that a high proportion of final year medical students will not continue to dismiss as "trivial" many of contemporary society's health problems?

References

1. Jacob A. Dundee Health Centre study. *J R Coll Gen Pract* (in press).
2. Anderson MJ, et al. Human parvovirus, the cause of erythema infectiosum (fifth disease). *Lancet* 1983;1:1178.
3. Milne AA. *When we were very young*. London: Methuen Children's Books, 1931:51.
4. Dickinson KG. A difficult case. *Pandora's box*. *Br Med J* 1983;287:951-5.
5. Farrow D. A difficult case. *Lancet*. *Br Med J* 1983;287:1114-5.
6. Freeling P. In: *A workbook for training in general practice*. Bristol: Wright PSG, 1983.
7. Blackburn J. *Towards earlier diagnosis*. 4th ed. Edinburgh: Churchill Livingstone, 1978.
8. *Blackburn's Gould Medical dictionary*. 3rd ed. New York: McGraw-Hill, 1973.
9. Howe JGR. Clinical judgement and antibiotic use in general practice. *Br Med J* 1979;1:1063-4.

(Accepted 22 December 1983)

Correction

A "Taylor made" practices

In the article by Dr S Williams (14 January, p 116) the year that Dr Henry Pearson Taylor died should have read 1945.