PRACTICE OBSERVED

Practice Research

Mobile patients in an inner London practice

FRANCES HANSON, MICHAEL WILKS

Abstract

In an area with the most highly mobile population in Britain 10% of the consultations in one practice were with temporary residents and a further 10% with newly registered patients. We analysed data gathered in 1980 about these two groups and about established patients that included demographic characteristics, the diagnoses made, and the actions taken by the doctors. Temporary residents caused a considerable amount of administration, but they were relatively straightforward to deal with medically and the practice received compensation for the work; with new patients problems were more time consuming, both clinically and administratively, and there was no financial compensation. A practice with a high turnover of its registered patients is at a considerable disadvantage compared with one in a more settled area.

Introduction

It is widely accepted that a highly mobile population causes many problems for the primary health services. The Acheson report and two publications from the Royal College of General Practitioners described the problems in detail,^{1,2} and in one survey 1705 out of 1802 general practitioners from all over the

LONDON W8
FRANCES HANSON, MD, research assistant
MICHAEL WILKS, MB, general practitioner

Correspondence to: Frances Hanson, Department of General Practice, St Mary's Hospital Medical School, Lioness Grove Health Centre, Grafton Street, London NW8 5EG.

United Kingdom identified high mobility as one of the most important social factors that influenced the pressure under which they worked.³

An analysis by the Office of Population Censuses and Surveys of the 1981 census showed that there is a wide range of mobility in different parts of Britain.⁴ Using migration into the district as an indicator of mobility, the lowest rate was found in Afan in South Wales, where only 5% of the population had moved in the year 1980-1. The highest rates tended to be in inner London, in large new towns, and in areas with a sizable proportion of service personnel or students. The highest of all was recorded in Kensington and Chelsea: 21% overall, rising to 30% for people aged 25-34 and to 43% for 16 to 24 year olds. The average for Britain was 9.6%.

Through their accident and emergency departments hospitals provide primary care services for some mobile patients—those who do not register either permanently or temporarily with a general practitioner, and those who cannot do so because they have no fixed abode. In the main, however, these services are met by general practice.

The impact of patient mobility on a practice depends largely on the number of temporary residents who attend the surgery, the rate at which patients register with and leave the practice (turnover rate), and the nature of the problems that these mobile patients present. We have analysed data relating both to temporary residents and to newly registered patients in one practice in the most mobile area of the United Kingdom and have drawn conclusions about some of the effects that mobility has on the doctor's work.

We can find no comparable published research about temporary residents in an inner city. Studies using different approaches have been done on newly registered patients, mostly in new estates and new towns. Whitfield, and Bain and Philip, showed a higher consultation rate in the first year of registration, while Goodman and Crombie stated that "only in patients reporting new episodes (of illness) within six months of registering was the episode rate noticeably high."⁵⁻⁷

It has been recommended that practices with a high turnover

1580

of patients should be compensated for the extra workload by increasing support services⁸ or by an increase in the capitation fee in the first year⁹—an idea echoed most recently in the Acheson report.

The setting

The surgery is situated in the W8 postal district of London, taking patients from Kensington and Earl's Court in the area of the Kensington, Chelsea and Westminster Family Practitioner Committee. There is no large housing estate locally. A few temporary residents come from a hostel run by MIND and attend a nearby psychiatric unit as day patients. The practice was expanding and the original principal had recently taken on a part time partner; on 1 July 1980, 3018 patients were registered with the two doctors. Anyone who settled in the practice area and wished to register was accepted; anyone who stayed in the area for no more than three months was taken on as a temporary resident if eligible to be one.

The age and sex distributions of the patients are unusual by national standards, though they reflect closely those of the local population in having a high proportion of young adults and a high ratio of women to men. There are many students. In 1980, 9618 consultations were held; about 10% of these were with temporary residents and another 10% were with newly registered patients as defined below.

The temporary residents are typified in two ways. Firstly, in the United Kingdom generally, about 40% of temporary residents sign on for three months rather than two weeks (information provided by the Statistical and Research Division, Department of Health and Social Security); in the area of the Kensington, Chelsea and Westminster Family Practitioner Committee in 1980 the figure was 82.6% (information provided by the family practitioner committee); and in the practice it was 75.4%. Secondly, the distribution of demand over the year is very different from that which would be found in, say, a seaside resort: we noted from the practice's quarterly statements from the family practitioner committee that the number of temporary residents was highest in the first quarter of the year. This is again typical of the area (information from the family practitioner committee). Clearly, these temporary residents are not people taking their summer holidays and, indeed, their consultations are least frequent from May to September.

Method

The practice was one of a group of seven, five of which were in inner London, participating in research with the department of general practice, St Mary's Hospital Medical School, and had computer facilities for storing and analysing the registration data and consultations of all its patients. The data that we analysed related to consultations in 1980.

The registration data for each patient included age, sex, civil status, social class (according to the Registrar General's classification), and date of joining the practice. The consultation data included up to two diagnoses for each consultation; whether or not a prescription, a certificate, or a follow up appointment was given; and details of up to three investigations and one referral for each diagnosis. The diagnoses were coded in the practice according to the ninth revision of the International Classification of Diseases¹⁰; for the other items the research group developed its own classification codes.

Each registered patient was allocated a unique number that formed part of the registration data and was recorded on a consultation sheet at every consultation. This number enabled us to link the two sets of data and to analyse the data from consultations for individual patients. Temporary residents had no registration data file, and though similar demographic information about them was recorded on the consultation sheet at each of their consultations, the absence of a unique number means that we cannot tell which patient participated in which consultation. Payment was received from the family practitioner committee in 1980 for 500 temporary residents,

BRITISH MEDICAL JOURNAL VOLUME 288 26 MAY 1984

but we do not know how often each consulted, or if any signed on more than once in the year. Comparisons with registered patients have therefore been made on the basis of the first consultation.

One of our aims was to compare the consultations of newly registered patients with those of the temporary residents. As more than three quarters of the temporary residents signed on for three months, we defined a "new patient consultation" as any consultation that took place in 1980 within three months of the patient registering with the practice. Consultations with patients who had been registered for more than three months were identified as "established patient consultations." We were thus able to compare the consultations of the temporary residents and with those of the patients with those of the established patients and with each other.

Results

The three tables show our findings for the three groups and indicate where differences between each pair are significant (using the χ^2 test). Table I gives the sex, age, marital state, and social class of the patients consulting. Table II lists the diagnoses made in terms of the major categories of disease according to the International Classification of Diseases. The number of specific diagnoses in the two smaller groups was usually too low for reliable comparison, but where they are high enough this is noted in the text. Table III shows what the doctors did as outcomes of the consultations.

Using the definition given above, there were 514 new patients who consulted. Of the registrations in 1980, 77% of them first consulted in the calendar month of registration, 89% within one month, 99% within two months, and 100% within three months. Their average consultation rate was 2.0, and over one year it was 4.0. Patients who were registered for more than a year and consulted had a consultation rate of 3.8 in 1980. The number of patients who registered with the practice in 1980 was 649 and the number leaving was 396. The turnover rate of registered patients was therefore 27%.

Comparisons between the three main groups of patients are as follows:

TEMPORARY RESIDENTS AND ESTABLISHED PATIENTS

Temporary residents were more likely to be aged between 10 and 20 years (the peak age being 21) and less likely to be under 10 or over 30. They are more often single and in social class II (which includes students). Their most frequent complaint is of a minor infection—vaginosis in category I, upper respiratory infections in category 8, and gastroenteritis in category 9 are all relatively more common. They have fewer mental disorders, either psychotic or non-psychotic (with the exception of alcohol dependence) and less

TABLE I—Patients consulting

	Established patients (n=7760)	New patients (n=514)	Temporary residents (n=960)
Sex			
Male	2212	295	279
Female	5548	219	681
Age (years)			
0-9	290	38	82
10-19	2360	381	82
20-29	1503	198	189
30-39	562	96	99
40-49	747	99	46
50-59	1305	172	61
60 and over	4647	619	710
Marital state			
Single	2254	298	193
Married	5506	216	767
Separated, divorced, widowed	203	27	26
Unknown	105	21	24
Social class			
I	169	21	38
II	2158	362	389
III	2443	387	381
IV	443	88	80
V	100	16	13
Unknown	473	62	61

*The 7760 consultations include 36 for which age, sex, marital state, and year of registration are missing and which have been excluded from other data in this table (n=7760).
†p<0.01; ‡p<0.05; §p<0.001; ¶p<0.005 significantly different from new patients.

TABLE II—Diagnoses made (from International Classification of Diseases, 9th rev.—minor surgical procedures)

Category	Established patients (n=7760)	New patients (n=514)	Temporary residents (n=960)
(1) Infectious and parasitic diseases (except cancer)	487	68	56
(2) Endocrine, nutritional, and metabolic diseases	80	0	11
(3) Endocrine, nutritional, and metabolic diseases	170	17	112
(4) Diseases of blood and blood-forming organs	80	0	6
(5) Mental disorders	857	18	64
(6) Diseases of the nervous system and sense organs	755	78	90
(7) Diseases of the circulatory system	277	28	26
(8) Diseases of the respiratory system	151	139	180
(9) Diseases of the digestive system	563	58	56
(10) Diseases of the genitourinary system	680	70	100
(11) Complications of pregnancy, childbirth, and the puerperium	27	0	3
(12) Diseases of the skin and subcutaneous tissue	434	65	73
(13) Diseases of the musculoskeletal system and connective tissue	712	73	84
(14) Congenital anomalies	24	0	2
(15) Certain conditions originating in the perinatal period	1	0	0
(16) Symptoms, signs, and ill-defined conditions	992	102	163
(17) Injury and poisoning	507	32	45
(18) Supplementary classification of diseases	45	0	7
(19) Supplementary classification of factors influencing health status and contact with health services	1632	168	204
(20) Minor surgical procedures	87	0	7
Total	9741	1000	1230

*p<0.01; †p<0.02; ‡p<0.05 significantly different from established patients.
§p<0.01; ¶p<0.05 significantly different from new patients.

TABLE III—Outcome of consultations

	Established patients (n=7760)	New patients (n=514)	Temporary residents (n=960)
Two diagnoses made	2081	272	232
Certificate given	366	48	51
Referral made	142	13	15
Follow up appointment made	661	60	65
Specific referral made	30	0	4

*p<0.01; †p<0.02; ‡p<0.05 significantly different from established patients.
§p<0.01; ¶p<0.05 significantly different from new patients.

disease of the nervous system and sense organs—otitis media and was in the ears in particular are rare.

The consultations of temporary residents are much less likely to be in the patient's home and are not as likely to produce two diagnoses or referral to a specialist, but investigations—largely bacteriology and simple haematology—are ordered more often and follow up consultations are more often arranged.

NEWLY REGISTERED PATIENTS AND ESTABLISHED PATIENTS

New patients are more strongly represented at all ages under 30 (especially children under 2), but they are less frequent over the age of 50; more are single and fewer are married, and though more are from social classes I and IV, fewer are from social class III. They are less likely to be diagnosed as having a mental disorder or to fall into the small category of "endocrine, nutritional, and immunity diseases and metabolic disorders." In category 19 pregnancy is more common, and category 16 shows that more diagnoses are left at the symptomatic level. The newly registered patients are accorded two diagnoses or even certificates less often, and investigations and follow up appointments are arranged more often.

1582

Conclusions

The data illustrate many of the effects of a highly mobile population on the work of a practice and show that it is primarily the high turnover rate of registered patients that causes the greatest clinical and administrative problems—with no extra remuneration. A practice in which this rate is high is therefore at a considerable disadvantage compared with one in a more settled area.

We thank Dr Carole Wigg for her part in recording the data, and Dr Conrad Harris for the help of the department of general practice, St Mary's Hospital Medical School, in setting up the recording system and for advice in the preparation of this paper.

References

1. London Health Planning Consortium, Primary Health Care Study Group. Primary health care in inner London. London: DHSS, 1981. (Acheson report.)
2. Royal College of General Practitioners. A survey of primary care in London. Occasional paper 16. London: RCGP, 1981.
3. Royal College of General Practitioners. Inner city. Occasional paper 17. London: RCGP, 1981.

4. Jarman B. Identification of underprivileged areas. *Br Med J* 1983;286:1705.
5. Davis F. People changing address. 1971 and 1981. In: *Population trends*. London: OPCS/HMSO, 1983:19-22.
6. Whitfield MJ. The relationship between year of registration and morbidity in a general practice. *J R Coll Gen Pract* 1982;32:699.
7. Bain DGG, Philip AE. Going to the doctor—attendances by members of 10 families in their first year in a new town. *J R Coll Gen Pract* 1975;25:821.
8. Goodman M, Crombie DL. Reported illness and change of residence. *J R Coll Gen Pract* 1982;32:699.
9. Miles DFB, Rowley A. Why not pay for high turnover? *J R Coll Gen Pract* 1980;30:40.
10. Office of Population Censuses and Surveys. Census 1981. County report. London: HMSO, 1982.
11. World Health Organization. Manual of the international statistical classification of diseases, injuries and causes of death (ninth revision). London: HMSO, 1978.
12. Harris CM. Mental illness in inner London. *Br Med J* (in press).

BRITISH MEDICAL JOURNAL VOLUME 288 26 MAY 1984

*Royal College of General Practitioners. A survey of primary care in London. Occasional paper 16. London: RCGP, 1981.

†Royal College of General Practitioners. Inner city. Occasional paper 17. London: RCGP, 1981.

‡Jarman B. Identification of underprivileged areas. *Br Med J* 1983;286:1705.

§Davis F. People changing address. 1971 and 1981. In: *Population trends*. London: OPCS/HMSO, 1983:19-22.

¶Whitfield MJ. The relationship between year of registration and morbidity in a general practice. *J R Coll Gen Pract* 1982;32:699.

‡Bain DGG, Philip AE. Going to the doctor—attendances by members of 10 families in their first year in a new town. *J R Coll Gen Pract* 1975;25:821.

§Goodman M, Crombie DL. Reported illness and change of residence. *J R Coll Gen Pract* 1982;32:699.

¶Miles DFB, Rowley A. Why not pay for high turnover? *J R Coll Gen Pract* 1980;30:40.

‡Office of Population Censuses and Surveys. Census 1981. County report. London: HMSO, 1982.

‡World Health Organization. Manual of the international statistical classification of diseases, injuries and causes of death (ninth revision). London: HMSO, 1978.

‡Harris CM. Mental illness in inner London. *Br Med J* (in press).

(Accepted 23 February 1984)

Pertussis: what percentage of children can we immunise?

D M JELLEY, A G NICOLL

Abstract

The immunisation records of 584 children who were born between 1978 and 1982, in a general practice of average social class distribution, were examined: 3.5% of the children would have been excluded from starting a course of vaccination including pertussis using contraindications established by the Department of Health and Social Security. A further 3.5% had reactions to immunisation that were judged severe enough to prevent completing the course of vaccination. In 1981 and 1982 13% of parents refused pertussis vaccination, considerably fewer than from 1978 to 80. Consequently, immunisation against pertussis rose from 51% to 84% over the five year period. Given the incidence of contraindications and the level of parental refusal, it is concluded that a pertussis uptake of 80% would be a reasonable target for any population.

Introduction

The incidence of pertussis in the United Kingdom, already declining in the early decades of this century, decreased further after the introduction of an effective vaccine in the late 1950s. In the years immediately before immunisation was begun between 30 000 and 90 000 cases were notified annually, with a case fatality of one per 100. In 1972, when vaccination rates were approaching 80%, there were only 2000 cases and two deaths in

England and Wales.¹ During the 1970s, however, anxieties about safety and efficacy of the vaccine arose after reports of vaccine induced convulsions and encephalopathy.² By the mid and late 1970s most infants no longer received protection against whooping cough, with immunisation falling to around 30%, and the illness began to re-emerge on a large scale with 66 000 cases notified in England and Wales in 1978.³

In response to these alarming developments the National Childhood Encephalopathy Study was set up to assess the relative risk of serious neurological disorders associated with immunisation in early childhood. The results of analysis led to an estimation of lasting neurological damage after pertussis vaccination of 1 in 310 000 single injections, much less than that after the illness itself.⁴ The Joint Committee on Vaccination and Immunisation has continued to recommend giving the pertussis vaccine, with due regard to stipulated contraindications.⁵

In our health authority there is a move towards measuring the percentage of uptake of immunisation in service units in the community such as health clinics or caseloads of health visitors. There are, however, no published data available on the frequency with which the state contraindications to pertussis immunisation occur in the community, which would be helpful in establishing realistic target figures for vaccination coverage. Thus we decided to examine the uptake of pertussis immunisation in a general practice area over the past five years and to identify the incidence of contraindications to vaccination.

Method

The practice that we studied lies on the outskirts of Nottingham and has five general practitioners, two health visitors, and other workers. The practice provides a primary medical care for a population of 10 000, which includes the mining village in which it is located and two smaller villages. It is the only general practice serving this population. The vaccination cards held by the practice of children born in the years 1978 to 1982 were examined. Data were collected for each child

Queen's Medical Centre, Nottingham

D M JELLEY, BA, assistant medical student

A G NICOLL, MRCGP, lecturer, department of child health

Correspondence to: Dr A G Nicoll, Department of Child Health, Floor E, East Block, University Hospital, Nottingham NG7 2UH.

on whether a full course of triple and polio vaccination had been given. When the pertussis vaccine had not been given any contraindication recorded by the health visitor was noted. Ambiguous statements on the vaccination cards were checked against the health visitor's records and the medical records when necessary. Children who did not receive pertussis vaccination who had no stated contraindications were assumed not to have had the vaccination by parental request. For the years 1980 to 1982 additional information was collected for the children from the notification of birth certificates concerning the occupation of the parents.

Results

A total of 584 vaccination cards were examined, which represented 93% of the children in the practice age-sex register who had been born between 1978 and 1982. The data concerning contraindications to pertussis were analysed into three categories in accordance with Department of Health guidelines.¹

Absolute contraindications—(i) history of severe local or general reaction to preceding vaccination; (ii) history of cerebral irritation or damage in the neonatal period; (iii) history of fits or convulsions in the child.

Relative contraindications—(a) Parents or siblings with a history of idiopathic epilepsy; (b) children with developmental delay thought to be due to a neurological defect or with neurological disease.

Inaccurate contraindications—This group includes children who were considered ineligible for pertussis vaccination according to now outdated criteria or to inaccurate interpretations of these criteria.

A fourth group of children consisted of those whose parents refused the vaccination in the practice age-sex register. This group included two adopted children whose family history was unknown and the medical opinion was not to vaccinate against pertussis. These two children were excluded from the analysis.

Table I shows the uptake of pertussis vaccination and the reasons for non-vaccination. The uptake of pertussis was 63% over the five years, but this figure conceals a rise from 51% in 1978 to 64% in 1982. There was also a concomitant decrease in the number of parents who refused vaccination, from 28% to 7% over the same period, a mean of 13% in 1981 and 1982. Table II gives data concerning the social class of the families. This did not differ appreciably from the distribution by social class in England and Wales.

TABLE I—Uptake of pertussis vaccination and reasons for not vaccinating children in one general practice between 1978 and 1982

Year	Received pertussis vaccine	Absolute contraindications		Relative contraindications		Inaccurate contraindications		Parents refused	
	No.	No.	%	No.	%	No.	%	No.	%
1978-82 (n=582)	371	63	28	5	10	2	40	7	133

TABLE II—Distribution of practice population by social class compared with distribution in England and Wales

Social class	I		II		III		IV		V	
	No.	%	No.	%	No.	%	No.	%	No.	%
Practice (n=582)	20	8	71	28	10	7	42	31	12	9
England and Wales	55	18.5	120	37.5	160	50	180	55.5	210	65.5

Table III shows the breakdown of the various groups of contraindications. Reaction to a previous dose of vaccine was the commonest reason for not completing the full course of vaccination. The commonest inaccurate contraindication was epilepsy in a second degree relative. It should be noted, however, that Department of Health guidelines on this point have been revised by the practice since the study. Until May 1982 the guidelines stated that epilepsy in any family member could be taken as a contraindication.

Discussion

The data collected in this study raise several issues relevant to the debate on pertussis vaccination. In this practice the rate of pertussis immunisation has climbed steadily since 1978 and now comprises almost 90% of the population. This has been achieved through a combined initiative of the health visitors and general practitioners, who were more and more concerned at the high incidence of pertussis locally. This is advantageous not only in terms of the protection thus offered to individuals but also by so raising the level of herd immunity in the community the vulnerable non-vaccinated infants aged 3 months or under run a much lower risk of contracting the disease.

The incidence of absolute contraindications to pertussis vaccination was 5%, but for two thirds of these children the

TABLE III—Contraindications to receiving pertussis vaccination during 1978-82

Type of contraindication	No.
Absolute (n=28)	
Severe reaction to first or second dose	19
Fit in child	4
Relative (n=10)	
First degree relative with idiopathic epilepsy	10
Inaccurate (n=40)	
Epilepsy in first degree relative	18
Severe allergic reaction to pertussis vaccine	1
Severe allergic reaction to other vaccine	1
Child had had pertussis	1
Parental refusal	1
Normal incidence	1
Severe respiratory illness	1
Gastro-oesophageal reflux	1

reason was a "severe reaction" to the first or second triple vaccination. This indicates that in a given population only 3.5% (1.5% absolute and 2% relative contraindications) should be advised not to start pertussis immunisation. Thus from the results of this study we estimate that in a given population 7% of children will either be advised not to start the course of pertussis immunisation or will fail to complete the course owing to reaction to the vaccine. A further 13% of parents (in 1981 and 1982) continued to choose not to have the vaccination, although the number is decreasing.

Thus a target of 80% uptake of pertussis immunisation seems to be achievable. Questions still remain, of course, as to how a "severe" reaction should be defined, and by whom this should be done. At present the decision seems to lie with the health visitors. There might be some value in clarifying the definition of "adverse reaction" by joint consultation between the doctors and health visitors. This is particularly important in view of the high incidence of minor reactions after routine immunisations—up to 15% using the adjuvanted vaccine.

Perinatal difficulties also cause problems of definition. Hull has illustrated the differences of opinion among professionals on several of these issues.² In the interpretation of contraindications further difficulties arise with respect to the precise definition of fits and neurological problems in the immediate family and with the need to exclude from consideration, for instance, febrile convulsions and metabolically caused fits. Allergic tendencies in the child have been taken as a contraindication and are still cited as such by the manufacturers of the vaccine. Moreover, manufacturers state "family history of epilepsy" as a contraindication, without distinguishing between first or second degree relatives, or between idiopathic epilepsy and epilepsy due to other causes. We welcome the news that one company (Wellcome) is to bring its package inserts into line with Department of Health recommendations for epilepsy. A recent study has questioned whether even the established criteria of neurological disability or developmental delay should in fact be contraindications to vaccination.³ It is interesting that in the United States the only

either: The sleeplessness, and the colic and the panic of that, and being left alone at home to cope and feeling totally trapped... and I started to get extremely distressed and I was fragile he was and how easy it would be to hurt him. Frightening that I should feel that violence, and be able to rationalise it... but feeling very scared and upset by it.

Linda: I have never been so tired in all my life. I can remember being like a walking zombie, and I couldn't relax. She was a very small baby and I was so scared that something was wrong with her because of the bad pregnancy and the bad birth... and because you're tired (the worries) build up to enormous proportions.

None of this is extraordinary. The existence of mild or transient postnatal depression is common and well known to both the medical profession and lay public. The more subtle, and long term, disruption caused by the arrival of a first child, however, may still be largely unrecognised. As Ann Oakley has pointed out, whereas in a previous generation women gave up work on marriage, now they do so in large numbers only when they become mothers.⁴ Usually it is retirement or redundancy that makes doctors consider the effects of not having a job, but all of these effects were experienced by this group of mothers. Some regretted the loss of financial independence. All of them lost the loss of everyday social contact with other adults was keenly felt.

Alice: I was thinking about that the other day, with Christmas I was just thinking if I was at work we'd all be going down to the pub.

Ruth: It was an awful blow leaving work. That came as a real shock to me because I never realised I'd miss (the people) so much.

Many, but by no means all, mothers already had friends living nearby and unrelated to work, so that the isolation was minimised. For two, however, the isolation was made more worse by a move of house shortly after the birth.

Sarah: It was a horrible feeling—when you are used to having some support group and suddenly find yourself completely on your own with a baby.

Ellen: When we moved here I just stayed in the house all day, so I went nuts... one of the things I couldn't cope with, ever, about being at home is that you try and do something constructive, or you try and do some work. After you've been doing it for about two hours you think, well I'll sit down and have a cup of coffee now or a break, and there isn't anyone to talk to. I'm sure that's shared by loads of housewives.

Not all of these mothers regretted the loss of the job itself and the change of role it entailed. Some were only too happy to give up jobs which brought them little or no reward, and for them becoming a mother brought a distinct improvement in their enjoyment, and perhaps their status.

Margaret: I like being at home, seeing all my friends with their children.

For others, however, the loss of a career to which they had previously made a substantial commitment made adjustment to motherhood particularly difficult.

Alice (ex laboratory technician): You just lose your identity, don't you? I mean you're just the wife, and you fill in a form and what's your occupation? housewife.

Linda (ex geophysicist): I felt trapped six months later when the realisation grew that I was not going to get back to my job. That triggered a depression was that the company came back to me and said your maternity leave ends in a month... I had to say I can't go back, I'm still feeding the child.

Pam (ex assistant manager of a building society office): I object to my status being taken away. If I ever go back to work I've got that status to get back, which will take a long time.

Possibly the most insidious effect, and the hardest to explain, was the feeling that they had somehow lost their independence.

Alice: This thing was so dependent on me, that's what I couldn't bear; I couldn't get away from her.

Pam: I feel we never used to go out before we had children, but don't you miss it when you can't go out.

Joan: We all want a little bit of independence. I could have it, that's the thing that annoys me. My husband plays golf on a Saturday morning and he says "If you want to go and do something, you can do it, I don't mind." But you know more people and more things going on when you worked, and I say "I don't know what to do."

Pam: (What stops me going out) is the tiredness, and the [misconception isn't there. You feel your responsibility is still with your child.

The point is made not because it is impossible for mothers of young children to go out in the evening on their own. But it has become more difficult to do so more because of looking after their children all day makes them too tired, or because of the powerful tie they feel to their children.

What can general practitioners do?

Can general practitioners do anything to help first time mothers? Most doctors and patients would agree that recognising and treating severe or psychotic postnatal depression is a doctor's responsibility. There remains, however, considerable doubt, both inside and outside the medical profession, whether doctors should become concerned in matters that fall outside the traditional medical model of diagnosing and treating disease. Few of this group of mothers had discussed their problems with health visitors or doctors, perhaps because they felt professional people had little to offer.

Esther: I want to go and get it medicalised. I didn't need to go to some medical jerk.

Linda: I'm not going to have Valium... that's an admission of failure.

Whatever their intentions, it seems unlikely that doctors will have any opportunity to help women to adjust to motherhood if they are perceived as concerned only with treating disease. Whether you agree with the Royal College of General Practitioners that general practitioners should undertake more responsibility for promoting health, in this area at least health visitors might claim to be more suited to the task.

There are, however, two ways in which either of these groups might offer practical help. On becoming a mother a woman is faced with the demands of a dependent infant while simultaneously adjusting to the loss of status and independence that comes with giving up a job. While the medical profession, and difficult is the gap between a woman's individual experience and the conventional view of motherhood as a happy and fulfilled state, embodying what many mothers feel is expected of them by the rest of society. The idea of a loving mother who has plenty of time to satisfy all her child's needs, as well as keep a home spotlessly clean and produce hot meals at regular intervals. (Note that it entails constant sacrifice of the woman to the needs of others.) Some mothers manage to achieve this and get a lot of fulfillment from succeeding, but for many others it is an impossibly remote goal to aim at, and their

1584

stated absolute contraindication to pertussis vaccination is the occurrence of convulsions, encephalitis, local neurological signs, or collapse after the first or second dose.¹

Clarification of the issues surrounding contraindications is important for parents and health visitors alike. This requires local and national initiatives, with doctors and health visitors discussing together the specific details and difficulties concerning definition of contraindications. It is also important to discuss the criteria for determining whether a child has a severe enough reaction to vaccination to warrant stopping the course of immunisation. To facilitate closer monitoring community immunisation records might be reorganised so that uptake rates are readily available not only for health districts but for units such as nurseries, schools, clinics, or health visitor lists. Such measures should take us closer to the target of 80% uptake of pertussis immunisation.

We thank the health visitors and general practitioners of Calverton practice, particularly Dr Tom Venables for support and encouragement, and June Summers for preparing the text.

Life Changes

Birth of the first child

M D JEWELL

The rewards are immeasurable really. She's a super little kid. We couldn't have hoped for anything better, Jean.

It's been disastrous. My husband thinks my whole personality has changed. I've lost a sense of myself, Alice.

Unlike other major events that influence our lives, becoming parents is now for most of us a matter of choice. It is an event of which one is always warned and for which some preparation might be made. Yet interviewing women who have had their first babies in the last few years about their experiences has reminded me of some of the sacrifices entailed in parenthood. I talked to 10 women, whose first children are now aged between 22 months and 5 years. Their ages at delivery were between 25 and 30 years. There is no pretence that they constitute anything other than a selected group: because of my interest I chose to interview some whom I knew and had some experiences, as well as some with happier memories. They were predominantly middle class (this partly reflects the composition of my practice), and in particular all were working full time before delivery. At the time of interview two were working full time, three part time, and five were not working at all, although only one of those now working had returned to work within a year of the birth.

All these mothers had found many rewards in having children. None of them regrets the decision to become a mother, but some of them had taken a long time to adapt to and enjoy motherhood. Arriving home with a small baby demanding full

time care is an event for which some women were unprepared. The words "surprise" and "shock" were frequently used to describe their reactions:

Sarah: I think a lot of people like us, who have their kids later, are just not prepared for it.

Ellen: I remember thinking I was going to leave (the baby) in the hospital. I was very surprised to be bringing it home because I just hadn't thought of myself with a child.

Conventional wisdom ascribes part of this problem to the predominance of small families: women who have contact with babies before they have their own find initial handling easier. One woman felt this was true for her:

Pam: (The baby) never got me down. I put it down to because I had come from a big family and I always knew how to handle children.

Similarly, there was support for the belief that the initial period of adjustment may be made more difficult by the sense of failure from an abnormal delivery:

Sarah: Immediately he was born everything collapsed because he was in intensive care. First reaction is total panic... there's something wrong with the baby and they won't tell me. A lot of women who have caesareans feel a phenomenal sense of failure at not having given birth naturally. I know you're not supposed to do it but I think quite a lot of people do.

Loss of sleep was mentioned as a problem in the early stages, but effects ranged from mere tiredness to a more profound mental disturbance.

1586

shortcomings only induce a sense of failure and guilt. Under these circumstances doctors may be able to help by clearly distancing themselves from the ideal, by helping mothers to understand that they are not alone in having such problems, and by giving them permission to express negative feelings to their children without guilt. What some mothers need is to be encouraged to think of their own needs, to understand, as one put it, that "mother and child" means "mother" as well as "child". For instance, many described the occasional need to do something without their children, and there is no reason why doctors shouldn't encourage mothers (and fathers) to arrange it. This sort of attitude may be adopted, I think, without either detracting from the sense of success enjoyed by "perfect mothers" or encouraging child neglect.

If these women were unwilling to share their worries with professionals, several accounts underlined the value of good close support: from a mother living nearby, paradoxically from an unemployed husband being constantly at home, or from self help groups, which in one case had been set up by a health visitor.

Esther: The best therapy was other women saying "I'm glad you said that" or "Yes, I remember feeling like that too," and what is awful is that some women feel that they cannot open about it for fear of the medical reaction.

Jean: The group provided me with other mothers with children of a similar age who lived near me that I could discuss problems with without going to a more specialised service.

It has been argued that informal contact with other mothers is the main value of well baby clinics, but formal postnatal groups could provide such help more reliably: perhaps general practitioners could ensure that they were offered to every first time mother in their care. Helpful as such groups may be, it may be dangerous to assume that women will always be able to share problems when all they have in common is motherhood.

Linda: The trouble with going to things like National Childbirth Trust classes is you meet the perfect mums there and I used to come home feeling more inadequate. They were obviously very content with motherhood and childrearing.

If such meagre suggestions seem to offer cold comfort to mothers facing major disruption to their lives that is because the problem is a political and social one rather than a medical one.

Changes

I recently had cause to ponder the changes that have taken place in my practice on the Isle of Lewis over the past 40 years or so, although I have been here for only a quarter of that time. The people on the island—the Macdonalds, the MacLeans, the MacIntoshes, MacInnans—have remained almost unchanged for centuries, but communication on the island and patients' medical expectations have changed dramatically.

My predecessor's first car was an early Morris, fitted with its engine and suspension against the hills and potholes of the unmetalled roads stretching 35 miles from end to end of the practice. Breakdowns were frequent and were usually fixed by the doctor. Punctures were regular, and two or even three each trip were to be expected. Roads were improved year by year, and in 1953 a bridge was built across the Atlantic to Bernera, an island of some 400 souls, which meant that an exposed stretch of strongly tidal sea could be crossed without danger of being washed away.

Gates to control sheep were everywhere, and one local round trip of six miles meant opening and shutting 16 gates. There is only one gate in the empy prairie of Talisker, and only one in these days of handfull of miles, in the early days there were only eight in the whole practice. Even when these were often shared, and no medical confidences could be divulged when up to five other subscribers might be listening

one. We will be able to discuss solutions, in terms of more crèche and nursery facilities and better opportunities for part time work (and paid leave for fathers, a desire recently identified by the Equal Opportunities Commission), only when we have freed ourselves from the idea that mothers should be the sole carers for their children and acknowledge how difficult and exhausting it can be to be with a small child all day and every day.

Pam: When he's home on holiday my husband says to me "How on earth can you stick this day in, day out?"

This comment, I suspect, applies as much to doctors as to other members of the public. My awareness started with the birth of my own first child: my wife is one of the women interviewed here who reported a particularly bad experience. I have only become sensitive to the problems of other mothers since then, and wonder how I would have felt if my wife had adapted more easily to the role of mother or had been able, like some women, to continue her job part time. Invaluable as this experience was for my understanding, I am not supporting the idea, which is popular in some feminist circles, that doctors have to experience events to help others through them. The learning is always limited to a sample of one, and there is the constant danger of projecting the experience onto others. No doubt some people reading this article will already have decided that it illustrates this very point, by making motherhood seem much more difficult and depressing than it really is. Whether the joyous, carefree motherhood of popular belief is a reality for most mothers is impossible to say. I hope that I have shown that for some it is a myth. The aim of this article is not to replace one stereotyped image with another but to encourage others to recognise how unhelpful any stereotype can be. This seems to be an duty in people's lives which evokes widely differing responses. Our event is to be sensitive to the variety of our patients' needs and help them sort out whatever adaption suits them best.

I thank the women who agreed to be interviewed.

References

- 1 Oakley A. *From here to maternity. Becoming a mother*. Harmondsworth: Penguin Books, 1979.
- 2 Bell C, McKee L, Priestley K. *Fathers, childbirth and work*. Manchester: Equal Opportunities Commission, 1983.

In. Moreover, the more receivers lifted the fainter the message to the doctor became. Night calls in winter were uncommon, partly because relatives would have to cycle miles to the nearest phone to summon help.

Medical expectations were lower in terms of diagnosis and treatment. For instance, home deliveries were the norm. The first consultant obstetrician was appointed to the local hospital in Stornoway in 1936. The average birth rate a year in the practice from 1937 to 1946 was 16. Over the past few years the rate had fallen to two or three a year. The district nurse attended all confinement cases, sending for the doctor near delivery: the nurses themselves covered enormous distances by bicycle, the first Ford Popular cars being supplied in 1948 with the coming of the National Health Service. Dispensing has been obvious changes—away from formidable Wauchope's and the aromatic smells of a dozen popular mixtures for "heart", "lung", or "indigestion" to the more prosaic packages of tablets with scientific names but, thankfully, more portable contents. Social changes have been profound: there has been a revolution in support for a fringe population of crofting folk in a declining economy in terms of social security benefits and pensions, of home helps, and of sheltered housing units. In these days of cutbacks it is helpful to remember how much has been gained and maintained since 1945.—C HAV, general practitioner, Mallaig, Isle of Lewis.