on whether a full course of triple and polio vaccination had been given. When the pertussis vaccine had not been given any contraindication recorded by the health visitor was noted. Ambiguous statements on the vaccination cards were checked against the health visitor's records and the medical records when necessary. Children who did not receive pertussis vaccination who had no stated contraindications were assumed not to have had the vaccination by parental request. For the years 1980 to 1982 additional information was collected for the children from the notification of birth certificates concerning the occupation of the parents.

Results

A total of 584 vaccination cards were examined, which represented 93% of the children in the practice age-sex register who had been born between 1978 and 1982. The data concerning contraindications to pertussis were analysed into three categories in accordance with Department of Health guidelines.¹

Absolute contraindications—(i) history of severe local or general reaction to preceding vaccination; (ii) history of cerebral irritation or damage in the neonatal period; (iii) history of fits or convulsions in the child.

Relative contraindications—(a) Parents or siblings with a history of idiopathic epilepsy; (b) children with developmental delay thought to be due to a neurological defect or with neurological disease.

Inaccurate contraindications—This group includes children who were considered ineligible for pertussis vaccination according to too outdated criteria or to inaccurate interpretations of these criteria.

A fourth group of children consisted of those whose parents refused the vaccination in the practice age-sex register. This group included two adopted children whose family history was unknown and the medical opinion was not to vaccinate against pertussis. These two children were excluded from the analysis.

Table I shows the uptake of pertussis vaccination and the reasons for non-vaccination. The uptake of pertussis was 63% over the five years, but this figure conceals a rise from 51% in 1978 to 64% in 1982. There was also a concomitant decrease in the number of parents who refused vaccination, from 28% to 7% over the same period, a mean of 13% in 1981 and 1982. Table II gives data concerning the social class of the families. This did not differ appreciably from the distribution by social class in England and Wales.

TABLE I—Uptake of pertussis vaccination and reasons for not vaccinating children in one general practice between 1978 and 1982

Year	Received pertussis vaccine	Absolute contraindications		Relative contraindications		Inaccurate contraindications		Parents refused
	No.	No.	%	No.	%	No.	%	
1978-82 (n=582)	371	63	28	5	10	2	40	7
	133	23						

TABLE II—Distribution of practice population by social class compared with distribution in England and Wales

Social class	I		II		III		IV		V	
	No.	%	No.	%	No.	%	No.	%	No.	%
Practice (n=582)	20	8	71	28	10	7	108	42	31	12
England and Wales	55	18.5	120	37.5	160	50	85	26.5	3	0.9

Table III shows the breakdown of the various groups of contraindications. Reaction to a previous dose of vaccine was the commonest reason for not completing the full course of vaccination. The commonest inaccurate contraindication was epilepsy in a second degree relative. It should be noted, however, that Department of Health guidelines on this point have been changed by the new guidelines for the study. Until May 1982 the guidelines stated that epilepsy in any family member could be taken as a contraindication.

Discussion

The data collected in this study raise several issues relevant to the debate on pertussis vaccination. In this practice the rate of pertussis immunisation has climbed steadily since 1978 and now comprises almost 90% of the population. This has been achieved through a combined initiative of the health visitors and general practitioners, who were more and more concerned at the high incidence of pertussis locally. This is advantageous not only in terms of the protection thus offered to individuals but also by so raising the level of herd immunity in the community the vulnerable non-vaccinated infants aged 3 months or under run a much lower risk of contracting the disease.

The incidence of absolute contraindications to pertussis vaccination was 5%, but for two thirds of these children the

TABLE III—Contraindications to receiving pertussis vaccination during 1978-82

Type of contraindication	No.
Absolute (n=28)	
Severe reaction to first or second dose	19
Fit in child	4
Relative (n=10)	
First degree relative with idiopathic epilepsy	10
Inaccurate (n=39)	
Epilepsy in first degree relative	18
Reaction to previous dose of vaccine	14
Severe allergy	4
Neonatal or birth respiratory difficulties without permanent sequelae	1
Child had had pertussis	1
Parents adopted—family history unknown	1
Normal incidence	1
Parents refused vaccination	1
Neurological illness	1
Gastro-oesophageal reflux	1

reason was a "severe reaction" to the first or second triple vaccination. This indicates that in a given population only 3.5% (1.5% absolute and 2% relative contraindications) should be advised not to start pertussis immunisation. Thus from the results of this study we estimate that in a given population 7% of children will either be advised not to start the course of pertussis immunisation or will fail to complete the course owing to reaction to the vaccine. A further 13% of parents (in 1981 and 1982) continued to choose not to have the vaccination, although the number is decreasing.

Thus a target of 80% uptake of pertussis immunisation seems to be achievable. Questions still remain, of course, as to how a "severe" reaction should be defined, and by whom this should be done. At present the decision seems to lie with the health visitors. There might be some value in clarifying the definition of "adverse reaction" by joint consultation between the doctors and health visitors. This is particularly important in view of the high incidence of minor reactions after routine immunisations—up to 15% using the double vaccine.

Perinatal difficulties also cause routine immunisation. Hull has illustrated the differences of opinion among professionals on several of these issues.² In the interpretation of contraindications further difficulties arise with respect to the precise definition of fits and neurological problems in the immediate family and with the need to exclude from consideration, for instance, febrile convulsions and metabolically caused fits. Allergic tendencies in the child have been taken as a contraindication and are still cited as such by the manufacturers of the vaccine. Moreover, manufacturers state "family history of epilepsy" as a contraindication, without distinguishing between first or second degree relatives, or between idiopathic epilepsy and epilepsy due to other causes. We welcome the news that one company (Wellcome) is to bring its package inserts into line with Department of Health recommendations for epilepsy. A recent study has questioned whether even the established criteria of neurological disability or developmental delay should in fact be contraindications to vaccination.³ It is interesting that in the United States the only

Eather: The sleeplessness, and the colic and the panic of that, and being left alone at home to cope and feeling totally trapped... and I started to get extremely distressed. I was so fragile he was and how easy it would be to hurt him. Frightening that I should feel that violence, and be able to rationalise it... but feeling very scared and upset by it.

Linda: I have never been so tired in all my life. I can remember being like a walking zombie, and I couldn't relax. She was a very small baby and I was so scared that something was wrong with her because of the bad pregnancy and the bad birth... and because you're tired (the worries) build up to enormous proportions.

None of this is extraordinary. The existence of mild or transient postnatal depression is common and well known to both the medical profession and lay public. The more subtle, and long term, disruption caused by the arrival of a first child, however, may still be largely unrecognised. As Ann Oakley has pointed out, whereas in a previous generation women gave up work on marriage, now they do so in large numbers only when they become mothers.⁴ Usually it is retirement or redundancy that makes doctors consider the effects of not having a job, but all of these effects were experienced by this group of mothers. Some regretted the loss of financial independence. All of them lost the loss of everyday social contact with other adults was keenly felt.

Alice: I was thinking about that the other day, with Christmas I was just thinking if I was at work we'd all be going down to the pub.

Ruth: It was an awful blow leaving work. That came as a real shock to me because I never realised I'd miss (the people) so much.

Many, but by no means all, mothers already had friends living nearby and unrelated to work, so that the isolation was minimised. For two, however, the isolation was made more worse by a move of house shortly after the birth.

Sarah: It was a horrible feeling—when you are used to having some support group and suddenly find yourself completely on your own with a baby.

Ellen: When we moved here I just stayed in the house all day, so I went nuts... one of the things I couldn't cope with, ever, about being at home is that you try and do something constructive, or you try and do some work. After you've been doing it for about two hours you think, well I'll sit down and have a cup of coffee now or a break, and there isn't anyone to talk to. I'm sure that's shared by loads of housewives.

Not all of these mothers regretted the loss of the job itself and the change of role it entailed. Some were only too happy to give up jobs which brought them no financial reward, and for some becoming a mother brought a distinct improvement in their enjoyment, and perhaps their status.

Margaret: I like being at home, seeing all my friends with their children.

For others, however, the loss of a career to which they had previously made a substantial commitment made adjustment to motherhood particularly difficult.

Alice (ex laboratory technician): You just lose your identity, don't you? I mean you're just the wife, and you fill in a form and what's your occupation? housewife.

Linda (ex geophysicist): I felt trapped six months later when the realisation grew that I was not going to get back to my job. That triggered a depression was that the company came back to me and said your maternity leave ends in a month... I had to say I can't go back, I'm still feeding the child.

Pam (ex assistant manager of a building society office): I object to my status being taken away. If I ever go back to work I've got that status to get back, which will take a long time.

Possibly the most insidious effect, and the hardest to explain, was the feeling that they had somehow lost their independence.

Alice: This thing was so dependent on me, that's what I couldn't bear; I couldn't get away from her.

Pam: I feel we never used to go out before we had children, but don't you miss it when you can't go out.

Joan: We all want a little bit of independence. I could have it, that's the thing that annoys me. My husband plays golf on a Saturday morning and he says "If you want to go and do something, you can do it, I don't mind." But you know more people and more things going on when you worked, and I say "I don't know what to do."

Pam: (What stops me going out) is the tiredness, and the misgivings isn't there. You feel your responsibility is still with your child.

The point is made not because it is impossible for mothers of young children to go out in the evening on their own. But it has become more difficult to do so more because of looking after their children all day makes them too tired, or because of the powerful tie they feel to their children.

What can general practitioners do?

Can general practitioners do anything to help first time mothers? Most doctors and patients would agree that recognising and treating severe or psychotic postnatal depression is a doctor's responsibility. There remains, however, considerable doubt, both inside and outside the medical profession, whether doctors should become concerned in matters that fall outside the traditional medical model of diagnosing and treating disease. Few of this group of mothers had discussed their problems with health visitors or doctors, perhaps because they felt professional people had little to offer.

Esther: I want to go out to have it medicalised. I didn't need to go to some medical jerk.

Linda: I'm not going to have Valium... that's an admission of failure.

Whatever their intentions, it seems unlikely that doctors will have any opportunity to help women to adjust to motherhood if they are perceived as concerned only with treating disease. Whether you agree with the Royal College of General Practitioners that general practitioners should undertake more responsibility for promoting health, in this area at least health visitors might claim to be more suited to the task.

There are, however, two ways in which either of these groups might offer practical help. On becoming a mother a woman is faced with the demands of a dependent infant while simultaneously adjusting to the loss of status and independence that comes with giving up a job. While the medical profession, and difficult is the gap between a woman's individual experience and the conventional view of motherhood as a happy and fulfilled state, embodying what many mothers feel is expected of them by the rest of society. The idea of a loving mother who has plenty of time to satisfy all her child's needs, as well as keep a home spotlessly clean and produce hot meals at regular intervals. (Note that it entails constant sacrifice of the woman to the needs of others.) Some mothers manage to achieve this and get a lot of fulfillment from succeeding, but for many others it is an impossibly remote goal to aim at, and their

1584

stated absolute contraindication to pertussis vaccination is the occurrence of convulsions, encephalitis, local neurological signs, or collapse after the first or second dose.¹

Clarification of the issues surrounding contraindications is important for parents and health visitors alike. This requires local and national initiatives, with doctors and health visitors discussing together the specific details and difficulties concerning definition of contraindications. It is also important to discuss the criteria for determining whether a child has a severe enough reaction to vaccination to warrant stopping the course of immunisation. To facilitate closer monitoring community immunisation records might be reorganised so that uptake rates are readily available not only for health districts but for units such as nurseries, schools, clinics, or health visitor lists. Such measures should take us closer to the target of 80% uptake of pertussis immunisation.

We thank the health visitors and general practitioners of Calverton practice, particularly Dr Tom Venables for support and encouragement, and June Summers for preparing the text.

Life Changes

Birth of the first child

M D JEWELL

The rewards are immeasurable really. She's a super little kid. We couldn't have hoped for anything better. Jean

It's been disastrous. My husband thinks my whole personality has changed. I've lost a sense of myself. Alice.

Unlike other major events that influence our lives, becoming parents is now for most of us a matter of choice. It is an event of which one is always warned and for which some preparation might be made. Yet interviewing women who have had their first babies in the last few years about their experiences has reminded me of some of the sacrifices entailed in parenthood. I talked to 10 women, whose first children are now aged between 22 months and 5 years. Their ages at delivery were between 25 and 30 years. There is no pretence that they constitute anything other than a selected group: because of my interest I chose to interview some whom I knew and had such experiences, as well as some with happier memories. They were predominantly middle class (this partly reflects the composition of my practice), and in particular all were working full time before delivery. The time of interview was two were working full time, three part time, and five were not working at all, although only one of those now working had returned to work within a year of the birth.

All these mothers had found many rewards in having children. None of them regrets the decision to become a mother, but some of them had taken a long time to adapt to and enjoy motherhood. Arriving home with a small baby demanding full

References

- 1 Joint Committee on Vaccination and Immunisation. *Whooping cough vaccination—a review of the evidence on whooping cough vaccination*. London: HMSO, 1977.
- 2 Kullen Ramfoll M, Schwartzman JS, Wilson J. Neurological complications of pertussis infection. *Arch Dis Child* 1974;49:46-9.
- 3 Billingham JS. Whooping cough—the public health viewpoint. *Health trends* 1983;18:71-3.
- 4 Miller DL, Ross EM, Aldeslade R, Bellman MH, Rawson NSB. Pertussis immunisation and serious acute neurological illness in children. *Br Med J* 1981;282:1595-8.
- 5 Joint Committee on Vaccination and Immunisation for the Secretary of State for Social Services. *Immunisation against infectious diseases*. London: HMSO, 1982.
- 6 Miller CL, Pollock TM, Clever ADE. Whooping cough vaccination: an assessment. *Lancet* 1974;ii:510-3.
- 7 Hull D. Interpretation of contraindications to whooping cough vaccination. *Br Med J* 1983;287:1231-3.
- 8 Miles RH, Hoakings GP. Pertussis: should we immunise neurologically disabled and developmentally delayed children? *Br Med J* 1983;287:318-9.
- 9 American Academy of Pediatrics. *Report of the committee on infectious diseases*. 19th ed. Evanston, Illinois: AAP, 1982:198-202.

(Accepted 14 February 1984)

time care is an event for which some women were unprepared. The words "surprise" and "shock" were frequently used to describe their reactions:

Sarah: I think a lot of people like us, who have their kids later, are just not prepared for it.

Ellen: I remember thinking I was going to leave (the baby) in the hospital. I was very surprised to be bringing it home because I just hadn't thought of myself with a child.

Conventional wisdom ascribes part of this problem to the predominance of small families: women who have contact with babies before they have their own find initial handling easier. One woman felt this was true for her:

Pam: (The baby) never got me down. I put it down to because I had come from a big family and I always knew how to handle children.

Similarly, there was support for the belief that the initial period of adjustment may be made more difficult by the sense of failure from an abnormal delivery:

Sarah: Immediately he was born everything collapsed because he was in intensive care. First reaction is total panic... there's something wrong with the baby and they won't tell me. A lot of women who have caesareans feel a phenomenal sense of failure at not having given birth naturally. I know you're not supposed to do it but I think quite a lot of people do.

Loss of sleep was mentioned as a problem in the early stages, but effects ranged from mere tiredness to a more profound mental disturbance.

1586

shortcomings only induce a sense of failure and guilt. Under these circumstances doctors may be able to help by clearly distancing themselves from the ideal, by helping mothers to understand that they are not alone in having such problems, and by giving them permission to express negative feelings to their children without guilt. What some mothers need is to be encouraged to think of their own needs, to understand, as one put it, that "mother and child" means "mother" as well as "child". For instance, many described the occasional need to do something without their children, and there is no reason why doctors shouldn't encourage mothers (and fathers) to arrange it. This sort of attitude may be adopted, I think, without either detracting from the sense of success enjoyed by "perfect mothers" or encouraging child neglect.

If these women were unwilling to share their worries with professionals, several accounts underlined the value of good close support: from a mother living nearby, paradoxically from an unemployed husband being constantly at home, or from self help groups, which in one case had been set up by a health visitor.

Esther: The best therapy was other women saying "I'm glad you said that" or "Yes, I remember feeling like that too," and what is awful is that some women feel that they cannot open about it for fear of the medical reaction.

Jean: The group provided me with other mothers with children of a similar age who lived near me that I could discuss problems with without going to a more specialised service.

It has been argued that informal contact with other mothers is the main value of well baby clinics, but formal postnatal groups could provide such help more reliably: perhaps general practitioners could ensure that they were offered to every first time mother in their care. Helpful as such groups may be, it may be dangerous to assume that women will always be able to share problems when all they have in common is motherhood.

Linda: The trouble with going to things like National Childbirth Trust classes is you meet the perfect mums there and I used to come home feeling more inadequate. They were obviously very content with motherhood and childrearing.

If such meagre suggestions seem to offer cold comfort to mothers facing major disruption to their lives that is because the problem is a political and social one rather than a medical one.

Changes

I recently had cause to ponder the changes that have taken place in my practice on the Isle of Lewis over the past 40 years or so, although I have been here for only a quarter of that time. The people on the island—the Macdonalds, the MacLeans, the MacIntoshes, MacInnans—have remained almost unchanged for centuries, but communication on the island and patients' medical expectations have changed dramatically.

My predecessor's first car was an early Morris, fitted with its engine and suspension against the hills and potholes of the unmetalled roads stretching 35 miles from end to end of the practice. Breakdowns were frequent and were usually fixed by the doctor. Punctures were regular, and two or even three each trip were to be expected. Roads were improved year by year, and in 1953 a bridge was built across the Atlantic to Bernera, an island of some 400 souls, which meant that an exposed stretch of strongly tidal sea could be crossed without danger to the passengers.

Gates to control sheep were everywhere, and one local round trip of six miles meant opening and shutting 16 gates. There is only one gate in the empy prairie between Talness and Talness. In these days of handsets, in the early days there were only eight in the whole practice. Even when these were often shared, and no medical confidences could be divulged when up to five other subscribers might be listening

one. We will be able to discuss solutions, in terms of more crèche and nursery facilities and better opportunities for part time work (and paid leave for fathers, a desire recently identified by the Equal Opportunities Commission), only when we have freed ourselves from the idea that mothers should be the sole carers for their children and acknowledge how difficult and exhausting it can be to be with a small child all day and every day.

Pam: When he's home on holiday my husband says to me "How on earth can you stick this day in, day out?"

This comment, I suspect, applies as much to doctors as to other members of the public. My awareness started with the birth of my own first child: my wife is one of the women interviewed here who reported a particularly bad experience. I have only become sensitive to the problems of other mothers since then, and wonder how I would have felt if my wife had adapted more easily to the role of mother or had been able, like some women, to continue her job part time. Invaluable as this experience was for my understanding, I am not supporting the idea, which is popular in some feminist circles, that doctors have to experience events to help others through them. The learning is always limited to a sample of one, and there is the constant danger of projecting one's experience onto others. No doubt some people reading this article will already have decided that it illustrates this very point, by making motherhood seem much more difficult and depressing than it really is. Whether the joyous, carefree motherhood of popular belief is a reality for most mothers is impossible to say. I hope that I have shown that for some it is a myth. The aim of this article is not to replace one stereotyped image with another but to encourage others to recognise how unhelpful any stereotype can be. This seems to be an duty in people's lives which evokes widely differing responses. Our event is to be sensitive to the variety of our patients' needs and help them sort out whatever adaption suits them best.

I thank the women who agreed to be interviewed.

References

- 1 Oakley A. *From here to maternity: becoming a mother*. Harmondsworth: Penguin Books, 1979.
- 2 Bell C, McKee L, Priestley K. *Fathers, childbirth and work*. Manchester: Equal Opportunities Commission, 1983.

In. Moreover, the more receivers lifted the fainter the message to the doctor became. Night calls in winter were uncommon, partly because relatives would have to cycle miles to the nearest phone to summon help.

Medical expectations were lower in terms of diagnosis and treatment. For instance, home deliveries were the norm. The first consultant obstetrician was appointed to the local hospital in Stornoway in 1936. The average birth rate a year in the practice from 1937 to 1946 was 16. Over the past few years the rate had fallen to two or three a year. The district nurse attended all confinement cases, sending for the doctor near delivery: the nurses themselves covered enormous distances by bicycle, the first Ford Popular cars being supplied in 1948 with the coming of the National Health Service. Dispensing has been obvious changes—away from formidable Wauchope's and the aromatic smells of a dozen popular mixtures for "heart", "lung", or "indigestion" to the more prosaic packages of tablets with scientific names but, thankfully, more portable contents. Social changes have been profound: there has been a revolution in support for a fringe population of crofting folk in a declining economy in terms of social security benefits and pensions, of home helps, and of sheltered housing units. In these days of cutbacks it is helpful to remember how much has been gained and maintained since 1945.—C HAV, general practitioner, Mallaig, Isle of Lewis.