the method appropriate; another 10 successes were reported but with a warning—in one instance CPAP stopped the obstructive episodes and restored breathing which was regular but so inadequate that hypoxaemia became dangerously worse.⁹ The patient was exceptional in having chronic respiratory disease, and such patients are quite different from those with the obstructive sleep apnoea syndrome.⁴ Nevertheless this one report suggests that no patient should be put on a CPAP regimen unless his response to it has been monitored with continuous measurement of oxygen saturation.

Another group using CPAP for obstructive sleep apnoea syndrome has now confirmed by computed tomography the assumption that the benefit is due to widening of the upper airway and claimed 100% success in 16 patients.¹⁰ Others, however, have found a similar improvement in patients with the obstructive sleep apnoea syndrome when they used only expiratory positive airway pressure, arguing that the improvement with either method might be by way of increasing functional residual capacity.¹¹

There seems no danger attached to nasal CPAP (with the exception quoted).⁹ Patients mostly tolerate it well and can sleep with it; it is not excessively complicated or expensive; and, indeed, there are good reasons to consider its use for recalcitrant snorers. Though they may not complain of daytime symptoms the enhanced quality of their sleep might improve their well being—as well as the peace and quiet of the home. With this treatment Berry and Block reduced the mean number of snores in their patients from 1000 plus to 23 a night,¹ with similar dramatic falls in the number of apnoeic or hypopnoeic events and the number and severity of desaturations.

It should not be too difficult to arrange a snore recorder for microcomputer analysis, so that the number of snores a night could be counted: the number appears to be some guide to the incidence of apnoea or hypopnoea and thus to the risk of the obstructive sleep apnoea syndrome and the need for treatment.

Until there are more centres in Britain with an interest in sleep disorders and equipped for "polysomnography," it is unfortunately difficult to know just who might be eager to apply themselves to the problem of providing nasal CPAP. It could well improve the quality of life for those who in the night, like Dionysius, make "much ado to take a breath."¹²

SHEILA JENNETT

Reader in Physiology,
Institute of Physiology,
University of Glasgow,
Glasgow G12 8QQ

¹ Berry RB, Block AJ. Positive nasal airway pressure eliminates snoring as well as obstructive sleep apnea. *Chest* 1984;85:15-20.

² Cherniack NS. Respiratory dysrhythmias during sleep. *N Engl J Med* 1981;305:325-30.

³ Guilleminault C, Dement WC. *Sleep apnea syndromes*. New York: Alan R Liss Inc, 1978:11. (Kroc Foundation Series.)

⁴ Catterall JR, Douglas NJ, Calverley PM, *et al*. Transient hypoxemia during sleep in chronic obstructive pulmonary disease is not a sleep apnea syndrome. *Am Rev Respir Dis* 1983;128:24-9.

⁵ Mauer KW, Staabs BA, Olsen KD. Upper airway obstruction and disordered nocturnal breathing in children. *Mayo Clin Proc* 1983;58:349-53.

⁶ Guilleminault C, Simmons FB, Motta J, *et al*. Obstructive sleep apnea syndrome and tracheostomy. Long-term follow-up experience. *Arch Intern Med* 1981;141:985-8.

⁷ Sullivan CE, Issa FG, Berthon-Jones M, Eves L. Reversal of obstructive sleep apnoea by continuous positive airway pressure applied through the nares. *Lancet* 1981;i:862-5.

⁸ Sullivan CE, Berthon-Jones M, Issa FG. Nocturnal nasal-airway pressure for sleep apnea. *N Engl J Med* 1983;309:112.

⁹ Krieger J, Weitzenblum E, Monassier JP, Stoeckel C, Kurtz D. Dangerous hypoxaemia during continuous positive airway pressure treatment of obstructive sleep apnoea. *Lancet* 1983;ii:1429-30.

¹⁰ Rapoport DM, Garay SM, Goldring RM. Nasal CPAP in obstructive sleep apnea: mechanisms of action. *Bull Eur Physiopathol Respir* 1983;19:616-20.

¹¹ Mahadevia AK, Onal E, Lopata M. Effects of expiratory positive airway pressure on sleep-induced respiratory abnormalities in patients with hypersomnia-sleep apnea syndrome. *Am Rev Respir Dis* 1983;128:708-11.

¹² Krieger MH. Sleep apnea. From the needles of Dionysius to continuous positive airway pressure. *Arch Intern Med* 1983;143:2301-3.

A fair trial?

Randomised clinical trials have become accepted as the standard procedure for comparing treatments, particularly for evaluating a new treatment against either an existing treatment or placebo. The many publications on this subject are continuing to grow in numbers but some aspects of trials are still not handled well. One important example is the assessment of the comparability of the patients in different treatment groups. Until recently this has received surprisingly little attention,¹ but an important paper by Lavori *et al* has considered the problem in some depth.²

In a clinical trial, randomisation ensures that the allocation of treatments to patients is left purely to chance. There is, however, no guarantee that the patients in the different treatment groups will be similar with respect to characteristics such as age, severity of disease, history, and so on. Despite randomisation, quite probably patients with a particular characteristic will not be equally common within each group. Imbalance of such baseline variables is clearly undesirable, as the essence of a controlled trial is the comparison of groups differing only as regards their treatment. Such imbalance really matters, however, only for variables with prognostic importance.

The usual and eminently sensible procedure is to begin the analysis of a trial by comparing the treatment groups with respect to several baseline variables. The fact that this is so common implies that most research workers realise that randomisation does not guarantee that the groups will be comparable. Unfortunately, the next step is often to perform significance tests, comparing these baseline characteristics in the different groups. Although carrying out such tests is not wrong, they are nearly always misinterpreted, perhaps seriously.

Presumably the intention behind performing significance tests to compare baseline variables is to see whether an observed difference is important. What such tests actually assess, however, is the probability that the observed difference (or a greater one) could have occurred by chance when in reality there was no difference. A moment's thought will show that when comparing baseline characteristics we know that the null hypothesis is true—because the treatment groups were produced by randomisation. If the randomisation was performed properly a difference of any sort between the groups before treatment must necessarily be due to chance. Tests of significance are thus testing only the fairness of the randomisation. We would expect one test in 20 to give a significant result with $p < 0.05$, and this is true however many patients are studied. We should not conclude from the lack of statistical significance that imbalance in a baseline variable did not affect the outcome of the trial; the comparability of treatment groups cannot be assessed in this way.

In very large trials this common error of interpretation is unlikely to be serious, but in smaller trials some quite substantial differences may occur by chance. Unless these differences are shown to be statistically significant they are nearly always ignored—as a result of the faulty reasoning described.² An extreme example of this was in a clinical trial of patients with acute myocardial infarction where the prevalence of previous infarction in the two groups was 7% and 30%.³ Even when differences are found to be statistically significant their possible importance is still often ignored.^{4,5} Armitage has suggested that major baseline differences will be rare.⁶ This is certainly true for large trials, but far too many clinical trials are performed on very small numbers of

patients, and in these the risk of imbalance is much greater.

How should imbalance be dealt with? The crucial question is whether the variable in question is related to the outcome being studied. If, for example, one treatment group is on average five years older than the other this will be of varying importance for outcomes such as survival time, reduction in blood pressure, or response to treatment for eczema. In the first instance both clinical knowledge and simple statistical analysis may be used to identify potentially important imbalances, and various statistical approaches can be used to investigate further. A more advanced statistical analysis, using modelling techniques, may be needed, in which case, as Lavori *et al* suggest, expert statistical help is recommended.² Only one of the 47 trials they reviewed had adopted this approach; they thought that over half could have benefited in this way. If there is a strongly prognostic variable there are good grounds for using the modelling approach even when perfect balance is achieved.^{2,6}

In many cases some variables of prognostic importance can be identified when a clinical trial is being planned. Various techniques exist for ensuring balance between groups for important variables, the best known being stratified randomisation.^{2,7} Such methods may be especially beneficial in small trials or where a particular variable is known to have a powerful influence on outcome.

DOUGLAS G ALTMAN

Statistician,
Clinical Research Centre,
Harrow, Middlesex HA1 3UJ

¹ Rothman K. Epidemiologic methods in clinical trials. *Cancer* 1977;39:1771-5.

² Lavori PW, Louis TA, Bailar JC III, Polansky M. Designs for experiments—parallel comparisons of treatment. *N Engl J Med* 1983;309:1291-9.

³ Flaherty JT, Becker LC, Bulkley BH, *et al*. A randomised prospective trial of intravenous nitroglycerin in patients with acute myocardial infarction. *Circulation* 1983;68:576-88.

⁴ Holman RR, Steenson J, Darling P, Reeves WG, Turner RC. Human ultralente insulin. *Br Med J* 1984;288:665-8.

⁵ Halliday HL, McClure G, Reid MMcC, Lappin TRJ, Meban C, Thomas PS. Controlled trial of artificial surfactant to prevent respiratory distress syndrome. *Lancet* 1984;i:476-8.

⁶ Armitage P. Importance of prognostic factors in the analysis of data from clinical trials. *Controlled Clinical Trials* 1981;1:347-53.

⁷ Pocock SJ. *Clinical trials. A practical approach*. Chichester: Wiley, 1983.

Ultrasound brain scanning in the newborn

Tired of reassuring mothers that the soft spot on top of baby's head is harmless and will close, neonatologists are now exploiting it as the window to the brain. Good quality coronal and sagittal images of the brain may be obtained using a real time ultrasound probe positioned over the anterior fontanelle.¹ Both periventricular haemorrhage and cystic degeneration within haemorrhagic or infarcted areas may be diagnosed reliably and ventricular size accurately measured.^{2,3}

Research studies using daily imaging have shown that 40-50% of preterm babies who weigh less than 1500 g suffer periventricular haemorrhage in the first three days of life.^{4,5} The haemorrhage (often silent) originates in the subependymal region; it may be confined here or it may extend by varying degrees into the ventricular system. A few infants suffer large haemorrhages with intracerebral extension. Some ventricular dilatation, presumably associated with obstruction to the flow of cerebrospinal fluid, is seen in 30-50% of babies with periventricular haemorrhage.⁶ Between 10% and 20% of these babies develop progressive enlargement of the ventricles and enlargement of the head requiring a cerebrospinal fluid shunt procedure.

Daily scanning has also identified the clinical antecedents of periventricular haemorrhage in preterm babies—notably respiratory distress syndrome, hypoxaemia, hypercapnoea, acidaemia, mechanical ventilation, and pneumothoraces.^{5,7} Perhaps its most important research application is in evaluating preventive measures against periventricular haemorrhage, such as phenobarbitone,⁸ ethamsylate,⁹ and vitamin E.¹⁰

Should ultrasound imaging of the brain now be introduced for routine use in special care baby units? The dynamic nature of periventricular haemorrhage and ventricular dilatation limits the value of a single scan performed during the first week of life. Daily scanning requires not only a portable scanner to examine ill babies by the cot side but a dedicated and experienced ultrasonographer. No specific treatment is available for periventricular haemorrhage so that the diagnosis by ultrasound of a silent periventricular haemorrhage cannot really influence a baby's care. When there are non-specific signs consistent with periventricular haemorrhage—such as recurrent apnoea, seizures, or circulatory failure—confirmation by ultrasound of periventricular haemorrhage should not deter the vigilant neonatologist from seeking a treatable coexisting disorder such as sepsis or hypoglycaemia which might itself be responsible for the baby's deterioration. Whether early recognition of progressive ventricular dilatation and treatment by repeated lumbar puncture reduce the incidence of clinical hydrocephalus or neurological sequelae is uncertain—and is currently the subject of a randomised multicentre trial (convenor, A Whitelaw, Hammersmith Hospital, London).

When brain scanning was first introduced there were hopes that it would help to resolve the ethical dilemma of whether to withdraw ventilatory support from critically ill, neurologically unresponsive babies with clinical evidence of hypoxic-ischaemic or haemorrhagic encephalopathy. The initial optimism must now be tempered with caution. Most parents prefer such babies to die a dignified death in their arms rather than in an incubator surrounded by the trappings of intensive care; it would be a pity if that humane approach was precluded when the brain scan failed to show gross abnormalities. Paediatricians who argue that the reward of having a more complete picture of their patients' condition is itself justification for daily brain scanning must be warned that a price has to be paid for routinely "looking" at babies' brains. Firstly, unless nursing and medical staff are properly counselled, the confident, positive, and optimistic ambience of the special care baby unit might be replaced by endemic despondency with whispers of "brain damage." Secondly, what should parents be told when, for example, a small, silent periventricular haemorrhage is observed by routine scanning? No matter how the doctor may try to convey the good prognosis there is no euphemism for brain haemorrhage. "Just a little leak of blood" or "a tiny vessel has leaked" might do if it were some other part of the baby's body—but not his brain.

Recently, Stewart and her colleagues at University College Hospital, London, have defined the probability of neuro-developmental disorders in preterm babies at 18 months based on the neonatal brain scan findings.¹¹ A group with a low risk (8%) for such disorders was clearly distinguished from a group with a high risk (83%). A single scan performed when a preterm baby is discharged from the special care baby unit might prove to be the most cost effective use of ultrasound brain imaging.

Finally, and most important, knowledge derived from the