PRACTICE OBSERVED

Practice Research

Computer controlled repeat prescribing used to analyse drug management

F DIFFORD

Abstract

In a practice that has been using a computer routinely for more than two years to print, monitor, and audit repeat prescriptions a program was written to analyse combinations of drug treatments. A method is described in which for given diagnoses the computer can identify the appropriate drugs and add them together in a systematic way to produce detailed printouts of the drug management in the practice. A technique by which the computer user can analyse them further on a pharmacological basis is shown.

Introduction

More than 200 general practices are now using computers for repeat prescribing, but there has been little progress in developing software for such systems.¹ Our practice, with five partners and a list of 9700 patients, has been using a computer for repeat prescribing and other purposes for more than two years and our system has been described in part.² We had accumulated considerable data on morbidity and repeat prescribing but had no method of directly analysing treatment with combinations of drugs on the computer. We wanted to develop a program that would show our drug management for various diagnoses for which repeat prescriptions are normally given, with particular reference to our treatment of hypertension.

The Surgery, 328 Wells Road, Knowle, Bristol BS4 2QJ
F DIFFORD, MA, MRCP, general practitioner

Method

I wrote a program, closely following the method that would be adopted if the task were done manually. The practice uses a system in the office that runs on a modified TRS-80 Model III computer with hard disk. Extracting information from the computer was simplified because in our system each drug is coded, and this includes different strengths and preparations of the same drug under different codes. Similarly, diagnoses are coded in a manner that is closely similar to the classification of the Royal College of General Practitioners. Thus there is no free text to search and all the information is contained in clearly defined parts of the problem summary and prescription files.

Firstly, the computer requests input of the code of the diagnosis—for example, hypertension—to be analysed and makes a preliminary search of the prescription records of patients with that diagnosis to identify all possible drugs being used in treatment. Since some of these drugs may be prescribed for indications other than the relevant diagnosis, each drug is then displayed in turn in order of frequency of use and the user (the doctor) asked if it is to be included in the analysis. This produces a list that is more easily obtained and more accurate than one the user could construct himself, as he would inevitably overlook some items.

The computer then carries out a second search, adds together the relevant drugs for each patient into drug combinations, calculates how often each one is used, and prints out the results. Up to this stage the program takes under 20 minutes, including the time that the user issues instructions, but the analysis may be further refined by renaming each drug according to pharmacological concepts. By using control keys the user can do this with little typing as it is essential that similar new names are absolutely identical where appropriate so that the computer can recognise them for addition. The computer then recalculates the frequency with which these pharmacological combinations are used and prints out the results. The user has the option of continually redefining and printing this list to obtain the most appropriate breakdown of drug treatment profiles.

Results

The first analysis of the output for hypertension (not shown) gave 63 different regimens of 31 hypertensive drugs for 254 patients given

repeat prescribing. These figures are inflated because they include the same drug in different strengths and preparations and both brand and generic names presented separately. The analysis is valid, however, because the practice is dealing with this range of preparations and dosages and thus the potential for confusing patients, doctors, and ancillary staff is great. Figure 1 shows the printout which results from renaming drugs to eliminate the variation of strength and brand name—for example, Isordil La and Propranolol 80 mg into

FIG 1—Computer printout of drugs prescribed for hypertension

FIG 2—Computer printout of redefined drugs prescribed for hypertension

Propranolol. It still includes 50 drug combinations. The combinations are listed alphabetically in the form of a histogram and the display is limited by the character set of the computer and the line length of the dot matrix printer. Figure 2 shows a printout after the drugs were redefined according to our views on pharmacological classification—for example, Nadolol and Propranolol into Beta 1 + 2 blockers that is, non-selective. Figures 3-5 show simpler printouts for asthma and epilepsy, and for diabetes treated with oral hypoglycaemic drugs.

the last two of which are produced more quickly because of the small numbers. Each of these has also been simplified by renaming in the program the drug presentations in the practice formulary.

Discussion

Looking at the analysis for hypertension (fig 1)—the prescribing habits of five partners plus those inherited from other general practitioners and outpatient departments—it is easy to identify favoured drug treatments, those that are disappearing, and some in which the pharmacological actions are similar. Such variety surprised us because we had already started to reduce the range of drugs prescribed.³ If we use one drug from

FIG 3—Computer printout of drugs prescribed for asthma

FIG 4—Computer printout of drugs prescribed for epilepsy

five categories—for example, diuretic, adrenergic blocker, centrally acting drug, vasodilator, and angiotensin converting enzyme inhibitor—alone or in combinations of two or three, using simple arithmetic the possible number of combinations totals 25, and for each alternative drug that we add to each category another 11 combinations become possible. The result would undoubtedly be the same for many practices similar to ours, but the computer is totally revealing and comprehensive.

It might be argued that there are too many drug combinations being used here, and that their number does not represent a

true variation in pharmacological properties of benefit to the patient. In trying to determine the fundamental attributes of a drug each doctor has his own concepts, such as whether the membrane stabilising effect of β blockers is clinically important, whether indapamide has a hypotensive effect below a diuretic dose, and where to classify prazosin, but the program will allow for such individuality. Thus figure 2, although still complex, begins to make sense of profiles for the treatment of hypertension and points to further possible rationalisation. We easily reached agreement on preferred hypotensive drugs in the light of this information and can critically examine the control and side effects of different treatment combinations.

The detection and treatment of hypertension in general practice is gaining momentum but in reality there are difficulties in doing it well.⁴ Confusing and multiple drug regimens will not help, and the practice nurse and trainer will make little sense of figure 1. The complexity of these profiles ought not to be added to by introducing a new preparation without critically evaluating its place in improving management. An unnecessarily complicated method of prescribing can lead to poor shared understanding for both patient and doctor of treatment and problems of compliance. It may also divert energy from detecting new patients with hypertension who have a good chance of benefiting from treatment.

Figure 1 shows that only 254 of the 290 hypertensive patients have repeat prescription files, and of these, 19 have no identified hypertensive drug prescribed, although they are on other repeat drugs. Notes of all 290 have been examined to confirm that hypertension was definitely detected at some stage, and our diagnostic index list must in any case be incomplete. Thus using records of repeat prescribing to establish a list for audit is questionable. The discrepancy is explained by new patients with hypertension who have not been stabilised to the stage of receiving repeat prescriptions, by patients who attend or are visited regularly who have other major problems, and by patients whose hypertension no longer requires treatment because of the onset of ischaemic heart disease or a cerebrovascular accident. A small group of patients seems to compare

mentalise their view of drugs into those they obtain from fairly frequent consultations—for example, hypotensives—and those they obtain by repeat prescription and almost regard as "non-drugs," such as psychotropic drugs. Changes in treatment will not show up in records of repeat prescribing until after reestablishment. Thus all of the treatment profiles currently used will not be included on the computer but the computer will probably produce an accurate overall picture. The computer output, however, alters from morning to afternoon as it computes its prescribing data when the program is run. Thus it is difficult to track the figures in this article with the latest printouts sent for publication as ancillary staff make alterations to the data base.

Operating this kind of program would be more difficult and take longer if we had used free text instead of codes. Such considerations and warnings about potential drug interactions weigh heavily in favour of drug formulary based computer systems in general practice. It seems important to be able to produce reports quickly when needed, especially when, as in this program, user intervention is required. Anomalies, such as illogical combinations on the same patient prescription file (figs 1 and 2), are also brought to light and any patient on a particular drug regimen is easily identified so that errors may be corrected. It is possible to carry out a continuous manual audit by frequently running programs. It is desirable that ancillary staff initiate such searches as well. The time taken for the computer to produce reports such as described in this paper is a fraction of that taken by clerical staff. Programs must be written to analyse adequately the data that is being collected.

References

1. Turner S. Writing for micro. *Practice Computing* 1984 Mar; 2.
2. Telling JP, Davies RB, Difford F, Forster JE, Reading CA. Developing a computer controlled repeat prescribing system. *Br Med J* 1984; 289: 1786-2.
3. Coates J. Hypertension in general practice: what is to be done? [Editorial]. *Br Med J* 1984; 289: 880-1.

(Accepted 11 July 1984)

GPs and Their Staff

Absence from work

NORMAN ELLIS

The general practitioner with a small staff may face great difficulty when handling prolonged or frequent absence by an employee. There are several types of absence from work: certificated or uncertificated sick leave; unauthorised absences, including lateness; and authorised absences (apart from holidays, absence for public duties, antenatal care, maternity leave, and for attending a training course). It may seem presumptuous to advise doctors on how to deal with sickness absence among their staff, but we are all familiar with the

proverbial cobbler's children's shoes. Members of any profession or occupation may unknowingly forget or neglect to apply their special knowledge and skills to their own circumstances.

It may be helpful to explain the difference between authorised and unauthorised absences. Each employee will decide for himself the grounds on which absence may be authorised. It is important that you lay down clear rules and that your staff are familiar with these. Your staff should know the parameters within which absence is permissible and you should apply these fairly and consistently. For example, different employees will apply different rules to the circumstances when absence may be authorised for a bereavement. Undoubtedly, you will have your views about how much leave is appropriate and in what circumstances it should be allowed and will decide when leave may be authorised for this reason; but it is vital to decide on a policy and apply it consistently.

British Medical Association, London WC1H 9JR
NORMAN ELLIS, MA, FRD, under secretary

In any practice the direct and indirect costs of a high level of absence may be considerable: there may include the cost of providing sick pay, paying additional overtime to other staff, employing temporary staff, a reduction in the standards of your services to patients, disruption to working arrangements, and a lowering of morale, together with increased dissatisfaction among staff. Thus it is worth developing and maintaining a procedure to monitor and control absence from work.

Recognising patterns of absence

There has been considerable research into patterns of absence. Some trends have been found and a summary of these may alert you to the types of absence to watch for in your own practice: (i) young people tend to take more frequent and shorter periods of sick leave, and older people longer but less frequent periods; (ii) unauthorised absence may be common among new employees; (iii) established employees will have learnt the standards of your practice and know how to follow these; (iv) unfortunately, for the general practitioner the incidence of absence from work is a far greater problem in large and impersonal organisations but when it occurs in a small organisation the effects may be disproportionately disruptive.

It is difficult to deal with absenteeism unless you can establish the facts of each case and have kept adequate records. If all absences are recorded you will be in a better position to identify problem areas and also ensure that any action that you take is fair and reasonable. The principle of consistency may be embodied in the procedure (however informal or simple it may be) that you choose to apply when investigating and dealing with any case of frequent or prolonged absence.

Dealing with long term sickness

Absence because of long term illness is almost always a most difficult problem to handle, especially in a small organisation. If one of your staff is away sick for a long time, or is unable to come to work regularly and consistently because of chronic ill health, you are faced with the unwelcome task of keeping the job open to offer security to your sick employee, and thus (it is hoped) help recovery, and having to maintain the standard of service in the surgery. Your predicament is even greater if the sick employee has had long service with you and occupies a key position in the daily running of the surgery. Doctors probably find this dilemma even more intractable than most employers. Not only do they have the good fortune of being blessed with particularly loyal and long serving staff, but their professional calling instils empathy and compassion for chronic ill health.

There is obviously no easy answer to this problem, and the decision that you finally reach will depend on the circumstances of your own practice. It may be helpful, however, to consider the following questions if only to ensure that you have assessed the available options:

- (1) Just how much difficulty is being caused by this employee's absence? How long can you manage without a replacement? Can you fill the post satisfactorily temporarily? (Do not forget to ensure that your temporary replacement is aware that the job is temporary.) Can the work be shared or reorganised to accommodate a long term absence? Are your other staff willing to accept an additional workload or responsibility for a prolonged and possibly indefinite period?
- (2) Have you obtained a prognosis from your employee's general practitioner? (You may wish to obtain prior consent to approach your employee's general practitioner by providing for this in the contract of employment.) Have you considered seeking an independent medical opinion? Is a full recovery likely or will the employee be unlikely ever to return to the old job? (Particular difficulties may arise if one of the partners is the employee's general practitioner. An obvious conflict of interest may arise and

may make your decision more difficult to reach. Although as the employee's own general practitioner, you should have a good knowledge of the prognosis, it may be advisable to obtain an independent medical opinion before you reach a final decision. In circumstances where this is practicable, adjacent practices have been known to agree an informal arrangement whereby staff are encouraged to register in the neighbouring practice even if they were on one of the partner's lists when they originally joined the practice staff. But such an arrangement is not always feasible, especially in rural areas.

(3) Could your employee return to work if special arrangements were made—for example, could he be given with transport, the work reorganised, or the job redesigned?

(4) Are you likely to have any alternative, lighter, or less stressful work available? Would a part time job be feasible? Could a full time job be shared?

(5) The age and length of service of your sick employee will be vital considerations. Would early retirement be a mutually acceptable solution, perhaps with an enhanced pension or an ex gratia payment?

(6) Can you help the employee to find more suitable employment elsewhere in your locality? A general practitioner may well be placed to give such help.

Whatever conclusion you reach after taking these and any other factors into account, it is most important that your employee is made aware of the position. On medical and humanitarian grounds there may be good reasons for leaving the matter in the air, implying that you will accept the employee back when he has recovered. But this deliberate "indiscretion" must eventually come to an end when you feel that the job can no longer be kept open. At this stage you must inform the sick employee without delay.

Dealing with short term sickness

There are several steps that may be taken to control the level of sickness absence: (i) your practice manager can have an informal talk with your employee to see if any special help is required; (ii) you may wish to obtain an independent medical opinion on any employee who has frequent sickness absences; (iii) try to ensure that you have a proper procedure on the provision of medical certificates and that all your staff understand and follow this policy—for example, notification of sickness should be made as early as possible on the first day of sickness; (iv) where necessary the employee may need to be told that their level of sickness absence is putting their job at risk.

In previous articles on the statutory sick pay scheme I have mentioned the possibilities of operating a system of "self certification" for sickness absence of up to one week. The requirement to give a formal written statement of the reasons for his or her absence can help to concentrate an employee's mind on this important matter.

Frequent absence may indicate general ill health which requires medical investigation and attention, and if it continues it may suggest that the employee is unable to continue to do the job. Obviously, any general practitioner will encourage his staff to seek medical attention. Furthermore, you may need to explore whether there are any personal or domestic difficulties, or problems with the job itself. Particular attention needs to be paid to any pattern in frequent short absences on grounds of sickness; this may indicate that sickness is being used as a cover for other reasons, and may emphasise the need for good supervision of staff.

Dealing with unauthorised absences, including lateness

Unauthorised absence, the occasional day off work, when sickness is often given as a reason or excuse, is generally known as "absenteeism." Lateness and poor time keeping often

reflect the same syndrome. All such absences may be disruptive, particularly because they occur without any prior warning, although ironically enough they may become so patterned that others can predict when they will occur. "She always takes a day off sick before a bank holiday weekend."

Absenteeism and lateness often point to problems arising from the job itself; your employee may be reluctant to come to work because he or she is unhappy in the job, or simply doesn't like work. These problems need to be closely investigated. Is the supervision of your staff adequate? Are working relationships satisfactory? Is the job properly organised in terms of the quantity and quality of the work? Above all, such absenteeism must never be allowed to go unnoticed; you or your practice manager should inquire into the matter without any delay.

Other actions to be taken include: (i) keep a careful check of the individual's record of absenteeism, time keeping, etc.; (ii) must that the absent employee (or a member of the family or friend) should telephone the surgery before a specified time on the first day of absence (say 11.00 am); (iii) obtain some indication of how long the employee is likely to be away, which will help you cope with the absence; but you may wish to require your employee to ring before a specified time on each day of absence; (iv) ensure that the employee has an informal talk with your practice manager on the day immediately after the period of absence; (v) watch closely any absences that are next to holidays and look for other obvious patterns; (vi) look at the working environment—could this be contributing to absenteeism?

Continued absences of this type may lead to disciplinary action, which eventually ends in dismissal. The disciplinary procedures that you should follow are outlined in another article.²

Life Changes

Divorce

C F DONOVAN

There are a few variables in divorce that those affected inevitably experience this transition in different ways. Some of these differences are illustrated in the following accounts written by two of my patients. The first is by a father of three teenage children. The second is by a mother of three children aged 9, 13, and 17, who had been married for 17 years.

Personal account (father)

Even though my separation from my wife had been preceded by long years of widening alienation, caused by a continuing relationship with another woman and culminating in separate bed, cessation of social life, and constant talk of divorce and departure, the psychological impact of the actual event was unexpectedly severe. I supposed as best as I could the pain of telling the children that I wouldn't be living with them any

Temple Fortune Health Centre, 23 Temple Fortune Lane, London NW11 7TE
C F DONOVAN, MR, DRCO, general practitioner and part time senior lecturer, Royal Free Hospital.

Fair, firm, and sympathetic approach

Most general practitioners will rely on their practice managers to deal effectively with absence. He or she will need to be fair, firm, and sympathetic when handling the many varied and often difficult problems that lead to absences from work. To ensure fairness your procedures should be applied with consistency and your employees should be aware of these. To be firm in your approach to absence, you must never hesitate to act if you have good grounds for believing that sickness is being used as an excuse for indefensible absences. Sympathy is particularly necessary when dealing with long term sickness and special domestic or travelling difficulties. Much goodwill may be generated by giving staff authorised time off to meet occasional special needs—for example, the illness of a child or dependent relative, a family bereavement, or specific religious observances of minority groups.

I am most grateful to Dr Ewen Beamanwell and Mr Peter Syson and his colleagues of ACAS for their comments on this paper. I am, of course, responsible for any omissions or errors.

This article can provide only a general guide to the provisions of existing legislation. It does not offer a definitive statement of law.

References

1. N. Looking after your new employee. *Br Med J* 1984;289:417-8.
2. N. Statutory sick pay: it will affect you. *Br Med J* 1983;286:1819-20.
3. N. Statutory sick pay: how it works. *Br Med J* 1983;286:114-6.
4. N. Statutory sick pay: keeping records. *Br Med J* 1983;286:149-7.
5. N. Dismissal: fair or unfair? *Br Med J* (in press).

more—although suppression made the pain no less acute. What I had not expected was a wave of tenderness for my wife which, very briefly, restored between us a confidence and rapport that had been long lost.

In hindsight this was clearly mutual shock; the same kind of reaction that follows a long awaited death. Soon my wife's mood became more bitter, and more depressed, as I realised that, firstly, the triangle in which I had long existed had merely been perpetuated in a different form, and secondly, the move I had made was irrevocable. In the initial state of shock I had thought it possible that I would return and had even written to my wife about my strong desire to do so.

As things settled down my presumably mild depression settled in. The symptoms were early waking, reduced libido, and above all, apathy, not in my work, which was wholly unaffected by the traumatic events, but in my private life. Going out to a theatre or a film, for example, seemed difficult or impossible, like a heavy unladen household chore that you keep putting off. This was only part of a general inability to build a new relationship—I had joined the woman in question—which showed itself in several little attempts to escape.

The relationship with my children, fortunately, caused few direct problems, although I found meeting for "restaurant meals" as poor a substitute for living with them as I had expected. Where their welfare was concerned I had no difficulty in

combining well with their mother. That was virtually the only area of accord and agreement, however. The pressure caused by her continuing alienation between dependence and bitter recrimination has certainly borne heavily on me, so has the pressure from my partner to force a total breach with my wife.

Personal account (mother)

Even though we had not been very happy together for a few years I now realise that when my husband left me for another younger woman I went into a state of shock that lasted for months. It came as a complete surprise, not only to me and the children, but to all friends and relatives. With hindsight, I feel that I was far too well behaved and controlled and should have put up a greater fight—he probably wouldn't have stayed, but I made it easier for him by being reasonable and calm while he was around and only breaking down in the middle of the night in the privacy of my room. I could only think of a day at a time; the future meant nothing; I was absolutely numb.

The children were devastated. They adored him, and he had not only betrayed me, but he had betrayed them as well—he had left! They felt insecure, frightened, and guilty and did not know whom they should blame, even though I was honest and told them our breakdown in marriage was only one sided and that we should have both tried harder. Their absolute despair and misery nearly broke me; they cried night after night for months. They were ashamed. They couldn't talk to me, their father, or even their close friends. They wanted to hide like animals and lick their wounds. After the first couple of weeks their father persuaded them to see him and "Mary" frequently and regularly. They withdrew from him to hide their pain. He rarely saw their real distress. I think it would have been better if he had given them longer to adjust. They know that he loves them, but they are no longer the most important people—they come second. It takes time to come to terms with that. I have recently crucified "Mary," but two years later the children are still distant and disinterested in her. I think he expected them to accept her too soon.

It was lucky we both decided that we had to do the best for the children. He had been left money by his father, so I was able to stay in the family home and he has free access. Our standard of living has dropped and I am often worried about money, but my husband earns a good salary and, though not generous, he won't let us starve. I have found friends who have loved and supported us through the worst moments. Friends are essential but it is very difficult to be truly honest with them because they don't understand. They were and probably are still friends with both parties and won't and don't want to take sides. There should be a self help group of newly separated single parents in the locality, for these are the only people who "really understand" what it means to have to take every decision and to carry total responsibility 24 hours a day 52 weeks a year.

Doctor's comment

Today, almost one in three marriages breaks up. The United Kingdom has the highest divorce rate in the EEC, yet there is surprisingly little discussion among general practitioners about how we can help people like my two patients and their children.

In this article we look at three stages in the process of divorce where general practitioners may be able to help: (i) after the break up; (ii) before the break up; and (iii) preventing divorce.

AFTER THE BREAK UP

Both writers mention the word shock. It is in this state that those whose marriage is breaking up have to deal with the

problems of custody, finance, legal wrangles, and "finding themselves." During this period of turmoil and loss their general practitioner can give them support and possibly some insight into the bereavement they are experiencing. I agree with my second contributor who wrote along with her contribution: "your general practitioner will listen, help, and prescribe, although I am not sure that medication is beneficial because eventually you will have to come to terms with the situation yourself."

In both the above break ups the father left the family. Although for them there was the consolation of a new relationship, like many others they experienced "the persistence of attachment" to those they had left behind. They have told me how they loved their children and were concerned for their future health. Here a general practitioner can act as a "go-between" without breaking confidences. He can assist both parties at a time of bitterness and hurt to cooperate over the interests of their children. Of course the business of meeting for restaurant meals is no substitute for living with your children, but it is far better than losing contact, as half of fathers do a year after their divorce. Some American divorcees find a neutral place where the visiting parent can meet their children in a more natural way—for example, at a grandmother's house.

It is the discord between parents that upsets children. After divorce this may become extreme, and children are often left confused and depressed. Some are not concerned for their father's going to happen to them, others feel responsible, a few protective. I have heard children say "poor daddy, he almost had a breakdown and I didn't know how to help him." I try to get mummy to talk, but she will not help herself. They are likely to be present at the surgery with psychosomatic or behavioural problems. In one of the above families the academic performance of the youngest at school has deteriorated seriously.

Divorce often leads to financial difficulties. The families in our two examples were more fortunate in this respect than many. The conclusion of a report based on a national representative sample, conducted by Eliza Perry of the National Children's Bureau, and published as *Growing up in One Parent Family*, was that children might be disadvantaged less by the emotional consequences of divorce than by the financial hardship. Interviewing over 2000 divorced persons, they found that women's income dropped by half and men's dropped by a tenth in the same year that "intact" families increased their income by over a third, also about 40% of maintenance orders are in arrears. The effect of this financial plight is to force an increasing percentage of women to go out to work, often with inadequate resources to leave their children in the hands of a responsible person to enable them to get away. Today many have difficulty in finding employment.

A general practitioner, along with health visitors and social workers, should ensure that these families are receiving the full financial support that they are entitled to. This is costly to the state. In 1981, 392 000 one parent families were receiving regular weekly payments in 1979 it was 322 000. On average each family got £42.05 a week—a total for the state of £612m. The expenditure in 1980-1 on 100 000 children in care in England and Wales provided by local authorities was £284m. One can

not help wondering if we spent more on attempting to keep families together we might save both children and the state from unnecessary hurt.

BEFORE THE BREAK UP

Our two contributors were asked to write about the break up of their marriages after the event. The general practitioner may often be concerned in the run-up to a divorce. This may be full of tears and fears or suppressed problems. During this time many patients turn to the general practitioner. Often it is the wife who comes to the surgery. Her husband may refuse to act or may use the wife's help-seeking as a scapegoat, especially if

Divorce figures for 1960-82. Reprinted with the permission of the National Marriage Guidance Council from their Annual Report 1982-3.

there is a sexual or alcohol problem. (Some suggest that one in 20 marriages face a homosexual problem, and one in 10 some form of mental illness.) Commonly, the problems are presented as a physical illness or depression. It is the general practitioner's difficult task to identify the real problems behind such a "ticket." Dr Dommon suggests that general practitioners should be especially alert for such presentations during the first years of marriage, after the bustle of the wedding and the first 20 years when the children have left home.

Sometimes the discord in the marriage is all too clearly presented, as in their first example. Here violent and acrimonious scenes often lead to a total break-up. How should the general practitioner react? Perhaps the minimum that he can do is to sit down with both parties and give them time to discuss the situation as they see it with a third party present. It may then become clear where the problems lie and the general practitioner can suggest sources of help. If the marriage is clearly "dead" and the parties hopelessly alienated then the general practitioner can help them to seek suitable legal help and proceed with a relatively amicable divorce.

Some patients do not know that if married for more than three years and if there are no children or property they can simply file in a form (obtainable from the Citizen's Advice Bureau), send it off with a fee of a few pounds to the District Registrar, and get divorced. Many couples do not know about the fixed fee interview, where couples can get legal advice for a half hour for approximately £10, nor that solicitors can take on divorce cases through a "green" scheme, which allows them to deal with an income under £75 and the solicitor has to apply to the Law Society for legal aid. Often solicitors lead couples into bitter confrontation. Other solicitors (those in the Solicitors and Social Workers' Association) are "reconciling" couples in decreasing confrontation and encouraging clients to work together for the benefit of their children.

divorce before they are 16 years old. A growing number of these children are emotionally injured by the break up of their family.

The general practitioner should be alert to the need to identify those who are most at risk. He can support some of these patients and lead others to suitable sources of help. He can fight for more to be done by society to assist marriages to survive and to improve parenting skills. But in reality a general practitioner is limited by lack of time and the many other tasks expected of him. There is a danger that those who write articles such as this unwittingly add to the impossible expectations placed on family doctors. Some of our young, well motivated, vocational trainees may try and attain these impossible standards in too many spheres of general practice. The irony is that in the process they can place such a burden on their own personal relationships that it may lead to their own divorce. We should not forget that doctors have a higher rate of marital problems than the rest of the population. Perhaps a general practitioner's primary task is to do all he can to avoid this transition himself.

References

1. Commission of the European Communities. *Sex research annual report on the activities of the European Communities*. Luxembourg: Commission of the European Communities, 1981. (European Books, London.)
2. Wallerstein JS, Berkley J. *Surviving the break-up*. California Divorcees of Divorce (UK Marriage Research Centre). National Marriage Guidance Council Statistics, 1980.
3. Dommon J. *Marital pathology*. London: Darton, Longman and Todd, 1980.
4. Citizens Advice Bureau. *Divorce and family matters*. London: Citizens Advice Bureau, 1981.

Further reading

1. Chartered Institute of Public Finance and Accounting. *Financial, general and saving statistics*. Annual report. London: CIPFA, 1982.
2. Department of Health and Social Security. *Social security statistics*. London: HMSO, 1982.
3. Kanner L. Marriage, divorce and family patterns. In: *Families in focus*. London: Study commission on the family, 1981.

ONE HUNDRED YEARS AGO

The final arrangements in connection with the medical equipment of Lord Wolseley's expedition to Egypt have been made, and an agreement has been reached between the British and Egyptian Governments that throughout has been taken to insure their completeness and efficiency. In respect to the choice of medical officers, the Expedition is not less fortunate. Surgeon-General O'Neil, who will probably take the command of the advance corps, Surgeon-Major Corbett, who will join the Expedition, and Surgeon-Major R. Waters, Sanitary Officer, who has had great experience in the Ashantee war of 1874 under Surgeon-General Gordon, as well as other efficient medical officers, will insure confidence among those who are acquainted with their past history. The advance corps, with a strength of 40 officers and 164 non-commissioned officers and men, many of them wearing the medals for Egypt and South Africa, sailed on Thursday last from Woolwich Arsenal in the steamship *Argonaut*, which will remain at Alexandria, and as soon as she has disposed of her freight of doctors, dispensers, and nurses, will probably be converted into a floating hospital. No one who has served less than eighteen months in the Medical Department has been selected. The despatch of a hospital contingent so large in proportion to the other reinforcements is understood to imply an intention on the part of Lord Wolseley to establish hospitals at various points in rear of his army, upon a system which has been for some time under consideration. Experience in Egypt has shown that the majority of the hospital cases are of a light character, a few days' treatment generally enabling the men to re-join their regiments; but the more serious cases, a place being carefully equipped, must be slow in moving through the crowded roads in the rear of a general action. (British Medical Journal 1884;ii:625.)

Specialised help may be needed if there is an alcoholic, gambling, violence, or sexual problem. Where couples are clearly not communicating, or where there is a history of bitterness, or acting as "passive victims" of the situation, the general practitioner may decide that counselling is an option that they might consider. Sometimes the general practitioner refers to the health visitor, social worker, or private counsellor, but most often he may advise a marriage guidance counsellor. If the general practitioner works with a local counsellor and knows what his local organisations have to offer he is more likely to make a "positive referral"—that is, he makes it clear to his patients that he is not rejecting them but rather offering them more than he can provide in "time" and "counselling skills." The general practitioner may, on the other hand, find that the couple do not wish this sort of help but only want further information, so that they can deal with the situation themselves. Here the health visitor, social services, or local Citizens Advice Bureau may be of help. The last is a fund of information—everything from grounds for divorce to rights of custody and custody, from conciliation services to injunctions to control violent spouses. They can also provide a list of local voluntary organisations, which may be of help.

PREVENTION

Over the past four years the number of divorces has increased relatively little, but during the seven preceding years it doubled (figure). Second marriages are leading to an increasing number of divorces. Every year an increasing percentage of children under 16 have experienced their parents going through this transition. 1980: 24%, 1981: 25%, 1982: 27%. Prevention should be considered as important as help with the problems caused by the marriage break up. Is there anything a general practitioner can do in this sphere? Not a great deal.

We can encourage more widespread and responsible contraception by discussing the services available with teenagers when we feel it is appropriate. Unwanted pregnancies leading to early marriage clearly should be discouraged. Those who marry under the age of 20 have a higher risk of divorce than those who wait until they are more mature. General practitioners can also urge couples to consider waiting for a year when they get married and settle down in their own relationship before embarking on their first pregnancy. Unfortunately, most of this advice is unlikely to be heeded. This is especially true for those who themselves did not receive all they needed from their parents during their childhood, and who may be searching for what they lack by entering early marriage, or for those rebelling against their parents or wishing to get away from home and finding marriage or pregnancy the only way of doing so. This brings us full circle, for perhaps the best form of prevention of divorce is to be found in increased efforts to keep families together during the years that children are growing through the complicated process of growing up, not only by providing them with two "good parents" but also by ensuring that they become good parents. These attempts at prevention can be successful if society feels that the family is worth fighting for. Perhaps of the general practitioner's most important tasks is to attempt to convince those in authority that more needs to be done to support marriages and to educate parents into how to manage the personal strains that every family generates.

Conclusions

With almost 150 000 new divorces a year, this transition has been described as a greater social problem than unemployment. Marriage break up is a painful process for all those concerned. Divorced parents have a high incidence of heart diseases and psychosomatic problems than the rest of the population. One in five children will experience their parents'

Some helpful organisations

Brook Advisory Centres
See local telephone directory or write to:
Brook Advisory Centre,
153A Euston Road,
London SE17.	Families Need Fathers,
97c Shakespeare Walk,
London N16 8TB.
Tel: 01-951 8932
(Advice on birth control and relationships)
Marriage Guidance Councils
See local telephone directory or write to:
National Marriage Guidance Council,
Herbert Grey College,
Little Chum Street,
Rugby CV21 3AP.
Tel: Rugby 73241	Mothers Apart from Their Children (MATCH),
64 Delaware Mansions,
Delaware Road,
London W9.
National Council of Clubs for Divorced and Separated,
13 High Street,
Little Chum Street,
Cambridge CB2 5BS.
Tel: 0223 246661
Gingerbread,
35 Wellington Street,
Farnham,
Tel: 01-240 0953
(Self-help organisation for one-parent families)	Bristol Courts Family Conciliation Service,
59 Broad Street,
Commercial Rooms, Box 101,
Bristol BS1 2EJ.
Tel: 0272 201813
National Council for One Parent Families,
255 Kentish Town Road,
London NW5 2LX.
Tel: 01-257 1361	British National Step-Family Association,
329 Hills Road,
Cambridge CB2 2QT.
Tel: 0223 246661

Solicitors Benevolent Association, incorporating The London Solicitors and Families Association:
30 Lonsdale Chambers,
27 Chancery Lane,
London WC2
Tel: 01-405 2840