PRACTICE OBSERVED

Practice Research

Influence of patient characteristics on test ordering in general practice

ROBERT M HARTLEY, JOHN R CHARLTON, CONRAD M HARRIS, BRIAN JARMAN

Abstract

Information regarding all consultations was collected in seven general practices for one year. From these data we report on the use of laboratory tests and its association with patient characteristics—including social class, age, sex, and diagnosis—and with which doctor was consulted. Most of the requests were for technically simple tests of low cost. There was a noticeable variation in the use of tests with regard to all patient characteristics. Diagnosis, identity of doctor, age of patient, and social class were each shown by multivariate techniques to be independently related to use of tests. Whereas fewer tests were used per consultation for social classes III-V compared with other social classes, more were used per patient per year for these same groups, reflecting in part the higher consultation rates of social classes III-V. Variation in diagnosis fully accounted for the greater test ordering for women.

Nearly two thirds of all tests were ordered for 10% of all patients who consulted and 7% of all registered patients. The results of our analysis suggest that this

concentration is determined primarily by those patient characteristics most indicative of medical need and by which doctor is providing care.

Introduction

Laboratory testing is an important resource for medical practice. Though comparable figures are not available for the United Kingdom, it accounts for about a quarter of the cost of ambulatory care in the United States.¹ Little attention, however, has been given to the relationship between the use of laboratory resources and the clinical and demographic characteristics of patients for whom the tests are ordered. The results of studies of the variation in test ordering among doctors have shown that consideration of who the doctor is, independent of the characteristics of patients he or she is, is important.²⁻⁴

The characteristics of patients, particularly in light of their persistent associations with morbidity and mortality,⁵ would seem to be as important for understanding the distribution of laboratory and health care resources. The results of several studies have shown that social classes III-V are referred more frequently per individual for specialty evaluation and together with women and elderly people make greater use of general practitioner services.⁶⁻¹¹ Whether these observations hold true for the use of laboratory testing is unknown.

We undertook the following study to examine the use of laboratory resources and its relations to the clinical and demographic characteristics of patients in a population of 50 000 registered patients served by 22 general practitioners.

Methods

Twenty two general practitioners and their trainees in Greater London recorded data on every consultation in 1980. The consultations in which a National Health Service patient was seen either by a

Department of Community Medicine, St Thomas's Hospital Medical School, London

ROBERT M HARTLEY, MD, MSc, Millbank Memorial Fund scholar

JOHN R CHARLTON, MSc, lecturer in medical statistics

Department of General Practice, St Mary's Hospital Medical School, London

CONRAD M HARRIS, MD, FRCP, senior lecturer

BRIAN JARMAN, MSc, MRCP, professor

Correspondence and reprint requests to: Dr R M Hartley, Division of General Medicine, Department of Medicine, Brigham and Women's Hospital, Boston MA 02115 USA.

principal or by a trainee were analysed; this included 87% of all consultations recorded during 1980. The consultations that were excluded were: (a) visits by private patients; (b) visits to locum doctors; (c) visits to one principal who had incomplete laboratory reporting.

At each surgery consultation the doctor used a printed form on which was entered a confidential patient identification number, the age, sex, and social class of each patient seen, and the number and type of laboratory or referral requests that had been made. Typical combinations of tests—for example, electrolytes, liver enzymes—were entered as single requests. The only tests that were done on the premises were urine dipsticks, which were not recorded. The International Classification of Diseases (9th revision) was used to record patient diagnosis. Social class was defined according to the Registrar General's categories I, II, III manual and non-manual, IV, and V. Women who worked more than 20 hours a week were classified by their own occupations. Social classes I and II were combined for analysis and a category "social class not recorded" was added. For approximately 11% of the consultations no social class was entered. We therefore repeated regression analyses using only a group of doctors with near complete (>97%) recording of data on social class; the results were essentially unchanged, as reported below.

For their patient diagnoses were grouped, with some modifications, according to chapter headings from the International Classification of Diseases. Organ specific symptoms from chapter XVI (Symptoms, signs and ill defined conditions: 781-782, 785-789) were incorporated into the pertinent preceding chapter. Chapters XI (Complications of pregnancy, childbirth, and the puerperium), XIV-XVI (Constitutional anomalies, certain conditions originating in the perinatal period) were combined together; they constituted less than 10% of all consultations. A separate category was formed to include all visits of healthy and pregnant patients whose attendance was routine and not prompted by a specific complaint (Health maintenance: V01-V39, V60-V82).

ANALYSIS OF DATA

We used two tailed *t* tests to make simple comparisons of differences in the use of tests among groups. Linear modelling, using a least squares model, was used to examine the independent effects of patient characteristics on the number of tests ordered per consultation. We used the same linear modelling package (GLIM) to compare test use in the different levels of each independent variable.¹² Because of the large number of cells in the model three modelling analyses were necessary. The variables in the first were: sex, age, diagnosis, and doctor. Age was divided into four groups (<5, 5-14, 15-39, >39 years) that reflected test use; diagnoses were grouped as described above. Sex had no independent predictive value, and the second and third models examined men and women separately. They had as variables social class, age, diagnosis, and doctor. The number of tests ordered per patient a year was used as a dependent variable in linear modelling because diagnosis often varied from consultation to consultation as did, to a lesser extent, the patient's doctor. A *t* test for linear trend, adapted for data expressed in patient years, was used to test the difference among social classes in the amount of testing per year.¹³

Accuracy of coding was greater than 97% for test ordering in a 10% sample of forms from consultations. Detailed comparison in one practice of computer outputs with surgery records showed test ordering to be more than 97% accurate. Trends related to patient factors were compared with the information available from the second national study of morbidity in general practice and are presented below.¹⁴

Results

The mean number of patients on the doctors' lists was 2263 and the mean number of consultations per person a year was 2.7.

PATTERNS OF GENERAL TEST ORDERING

Laboratory resources were used for small proportions of both the populations at risk and the group of patients who consulted. Two thirds of all tests requested during the year were ordered for less than 7% of the population at risk and for about 10% of all the patients

who consulted. Eighty one per cent of these requests were for single investigations. Haematology accounted for 22% of all tests, chemical pathology 15%, radiology 20%, bacteriology 29%, and others (including pregnancy tests and cytology) 14%. Eighty eight per cent were "simple" investigations—that is, blood counts, routine chemical pathology tests (such as electrolytes and urea concentrations), pregnancy tests, plain x ray examinations, or bacterial cultures.

ORDERING BY AGE AND SEX

The numbers of tests ordered per consultation (mean (SE)) for women and men were 0.13 (0.002) and 0.10 (0.002) respectively; 0.34 and 0.19 tests were ordered per person (registered on list) a year for women and men respectively. Figure 1 and table 1 show that laboratory use per consultation varied with both age and sex. Children

FIG 1—Number of tests ordered per consultation.

TABLE 1—Numbers of tests, consultations, and registered patients according to age and sex

Age (years)	Women	Men
0-4	172	150
5-14	345	266
15-39	2153	1489
40-49	1716	1405
50-59	862	625
60-69	682	518
70-79	682	518
Total	9494	7397

were particularly unlikely to be investigated and some drop in the rate of investigation was also noted in older age groups. The highly significant difference between sexes in tests ordered per consultation was primarily due to differences during the reproductive years. Figure 2 shows a somewhat different pattern for the number of tests ordered per person a year. The peak during the reproductive years remained evident for women and the difference between the sexes was magnified. Services used for men, after climbing slowly in early life, accelerated in middle age, but only in the seventh decade did it approach the level for women.

Figures 1 and 2 also show data derived from the national study.¹⁴ In that report only the occurrence of an investigation was recorded, not the number. The distinction thereby introduced into a comparison with our data would be small, however, as noted above for our study, only one test was ordered in over 80% of those consultations in which any investigations were requested. The age and sex related trends in the national study are closely similar to those we report, though the absolute numbers of tests ordered differ.

FIG 2—Number of tests ordered per year for each registered person.

ORDERING BY DIAGNOSIS

Table 2 shows a more than tenfold variation in test use among diagnostic categories. Those diseases most heavily investigated are often not responsible for a large proportion of the overall burden on the laboratory; comparison of blood and genitourinary disorders is one of several interesting examples.

TABLE 2—Laboratory use according to diagnosis

Diagnostic category	Proportion of all consultations (%)	Proportion of all tests (%)	Tests per consultation
1 Infectious diseases	0.5	0.2	0.04
2 Neoplasms	1.8	0.9	0.20
3 Diseases of haemopoiesis	0.6	0.2	0.04
4 Disease of blood and blood forming organs	0.6	0.2	0.04
5 Mental disorders	7.2	3.0	0.09
6 Nervous system diseases	1.1	0.8	0.03
7 Respiratory system diseases	19.1	13.7	0.09
8 Digestive system diseases	2.4	2.0	0.17
9 Genitourinary system diseases	9.9	21.8	0.36
10 Skin and subcutaneous tissue diseases	7.0	6.0	0.07
11 Musculoskeletal and connective tissue diseases	7.5	9.9	0.19
12 Injuries	4.0	2.2	0.08
13 Health maintenance	11.2	8.2	0.06
14 Symptomatic	11.2	8.2	0.06
15 All ill defined conditions, certain personal conditions	8.0	10.4	0.16

INDEPENDENT EFFECTS OF PATIENT CHARACTERISTICS AND OF DOCTOR CONSULTED

We used linear modelling to investigate the independent relation of patient characteristics with the number of tests ordered per consultation, while controlling for the effects of the other characteristics. Table 3 shows that diagnosis accounts for the greatest part of the explainable variation among patients in test use. The identity of the general practitioner and age of the patient also contribute appreciably. Sex does not, the substantial difference between the sexes being fully accounted for by the other patient variables, most notably diagnosis.

To determine whether some doctors referred patients for specialty

TABLE 3—Variation in test use accounted for by patient characteristics

Variable	Proportion of variation due to regression	p value
1 Diagnostic category	67.4	<0.001
2 Patient age	8.4	<0.001
3 Patient sex	0.4	>0.05
4 Doctor	1.0	<0.001
5 Social class (women)	<1.0	>0.05
6 Social class (men)	<1.0	>0.05

*Explained by model (see Methods).

opinion rather than test them, we compared doctors' test ordering with their rate of referral for specialty opinion. The two were directly associated ($r=0.55$, $p<0.01$), suggesting no such substitution was occurring.

ORDERING BY SOCIAL CLASS

After controlling for the effects due to patient's age, diagnosis, and doctor, appreciably fewer tests per consultation were ordered for women in classes III, IV, and V and for men in class V than for patients of the same sex in classes I and II combined (table 1V). The same trends were found when analysis was limited to the patients of those doctors with near complete (>97%) recording of social class. The amount of variance explained by social class, however, was less than 1% for both men and women (table 3).

The last column of table IV gives the number of tests ordered per patient per year for each social class. A highly significant trend of

TABLE IV—Test use according to social class

Social class	Tests per consultation*				Tests per patient per year (all doctors)
	All doctors	Selected doctors†	Men	Women	
I & II	0.119	0.143	0.110	0.152	0.264
III	0.080	0.124	0.084	0.109	0.304
IV	0.115	0.086	0.086	0.087	0.237
V	0.124	0.144	0.071	0.103	0.292
Not recorded	0.124	0.144	0.071	0.103	0.292

*Adjusted for age, diagnosis, and doctor.

†Doctors with complete or near complete (>97%) recording of social class.

‡*t* test ($p<0.01$).

§*p* compared to social class I and II of same sex.

§*p* < 0.01.

increasing test use from social classes I and II to V was found. The reverse of the social class trend seen for test per consultation indicates that though patients of social classes III-V were more tests ordered at any particular consultation, they have more tests ordered for them over a period of time.

Discussion

When tests are ordered in general practice they are most commonly simple, inexpensive, and requested singly. In most consultations no tests were requested at all. This pattern, but not necessarily the actual number of tests, is largely unchanged from that of 20 years ago.¹⁻⁴ The effects of the availability of new facilities and technologies seem to be limited to the increase in the relative proportion of requests for routine chemical pathological tests. Hitchens and Lowe, among others, reported in 1966 that requests for chemical pathological investigations constituted about 5% of general practitioners' requests (radiology excluded); in our sample that proportion would be about 18%.¹

We used multiple regression techniques to examine individually the association of each patient characteristic with test use while the effects of other characteristics were controlled. Only diagnosis, identity of doctor, and the patient's age remained independently important. The pronounced difference in the patterns observed for men and women was fully explained by the differences in their diagnoses; once adjustments were made for the effects due to diagnosis no difference in test use remained.

We also showed an association between test use per consultation and social class. Though these findings were statistically significant, their importance is limited because they accounted for only a small amount of the difference in test use among patients. The pattern of medical care in social classes III-V, however, seems to differ from that in social classes I and II: the former see their doctors more often and have more tests ordered for them over a period of a year, but at any particular consultation fewer tests are ordered. Cummins *et al* noted a

not been shown to be useful.¹⁵ Secondly, of medical and non-medical characteristics of patients, the use of laboratory resources seems to be primarily related to those indicating medical need. Diagnosis and age, a proxy for disease severity, are related more to test use than are sex or social class. Other non-medical factors, such as a patient's education or ethnicity, might influence test use and deserve to be studied, though they are unlikely to be more important than the ones that we analysed.

Initially, most laboratory testing is requested for a small proportion of patients. This concentration of resources is primarily determined both by the nature of the patient's illness and by who is taking care of it.

We thank Drs Antoniou, Brent, Ciechan, Coffman, Cohen, Constantoudis, Crowther, Elder, Evans, Jacobs, Law, Lacey, Lister, Magill, Newman, Palmer, O'Brien, Strang, Wills, Wills, and Wills.

We also thank Dr Arnold Epstein, and Professors Frey and Schroeder for reviewing earlier drafts of the manuscript, and Kristin Mortimer for editorial assistance.

References

1. Schwartz AA. Changes in the use of ancillary services for "common" illness. In: Altman SH, Broder R, eds. *Medical technology: the impact on health care*. Washington DC: Department of Health, Education and Welfare, 1979:36-56.
2. Hitchens RAN, Lowe CB. Laboratory services in general practice. *Med Care* 1966;4:142-9.
3. Green AL. General practitioners and open access pathology services. *J R Coll Gen Pract* 1973;23:116-20.
4. Forster CB. The use of hospital laboratory facilities by general practitioners. *Br Med J* 1976;3:434-8.
5. Rose H, Abul-Soud B. *Diagnosis, pathology, and pathology*. London: Bell, 1972.
6. Scheraga SA, Kessler R, Gumpel J, et al. The use of laboratory tests and pharmaceuticals among physicians and effect of cost paid on subsequent use. *JAMA* 1973;229:967-73.
7. Charlton JRH, Hartley RM, Silver R, Holland WB. Geographical variation in mortality from conditions amenable to medical intervention in England and Wales. *Lancet* 1983;ii:993-6.
8. Cummins NO, Jarman B, White PM. Do general practices have different referral thresholds? *Br Med J* 1981;282:11-4.
9. Hicks J. *Primary health care*. London: HMSO, 1978.
10. Office of Population Censuses and Surveys. *General Practice and Population*. Series C15 No 8. London: HMSO, 1980.
11. Office of Population Censuses and Surveys. *General Practice and Population*. Series C15 No 9. London: HMSO, 1980.
12. Moenich DC, Kassir JP. The effect of an appointment system on demand for medical care. *Am J Epidemiol* 1974;100:443-50.
13. Murrell DC, Gage HC, Robinson NA. Referral to hospital by general practitioners. *Br Med J* 1980;281:147-50.
14. Nelder JA, Wedderburn RWM. Generalized linear models. *J R Statist Soc Ser A* 1979;142:37-64.
15. Robinson KJ, Boice JD. Epidemiologic analysis with a programmable calculator. *Biometrics* 1970;26:1-10.
16. Main KA, Kelly PM. Laboratory and radiological investigations in general practice. *Br Med J* 1983;287:1188-90.
17. South-East London Screening Study Group. *Screening for breast cancer in the South-East London Screening Study*. London: HMSO, 1983.

(Accepted 17 May 1984)

Diary of Urban Marks: 1880-1949

On my return from the war I intimated to the local medical committee my intention to resign. Their reply to this was to present me with two more diaries and a soup tureen suitably inscribed at the next meeting. Then I was begged to keep my position as they felt no one else could properly fill it. There was a good deal more sob said poured out and at the end I was forced to withdraw the resignation. Two or three years later, on the resignation of Dr Begg, I became chairman of the panel committee, a position which I was holding in 1925 when I was elected chairman of the Swansea Branch of the British Medical Association after being the secretary for a number of years.