Appropriate Technology

Orthopaedic aids at low cost

KO DE RUÏTER, OTTO LELIEVELD

Western orthopaedic aids are seldom suitable for Third World countries. They are often made from costly materials and require complex technology for their manufacture. In countries where resources are limited alternative aids must be made, and here we describe some of the aids and techniques that are in use in the physiotherapy department of a district hospital, leprosarium, and school for the handicapped in Zambia.

Orthopaedic aids need to be simple and durable, and in our view it is important that items are made locally, using raw materials and skills that are readily available. This obviates the problem of having to get foreign currency (which is difficult) to import the items of equipment. Home produced aids are also less expensive, and easy to repair, and their production creates local employment.

Crutches and walking aids

We make four types of crutches, three of wood and one of metal. The axilla crutch is made of wood, with a piece of car tyre at the bottom for protection. The top is made of cotton wool, foam, or another soft material and covered with leather (fig 1(a)). This type of crutch is suitable for small children. The two other types of axilla crutch shown in figure 1 have a better handgrip and are suitable for adults. Elbow crutches may also be made out of metal rods with a piece of rubber tubing to pad the handgrip (fig 1(d)). Welding equipment is needed to make this sort of crutch, but it has the advantage of being stronger than the wooden crutch and avoids putting pressure on the axilla. If welding equipment is available other items—for example, walking aids (fig 1(e)) and adjustable parallel bars—may be made. (Fixed parallel bars may also be made of bamboo and various other woods.)

Simple prosthetic above and below knee walking aids may be made out of plaster of Paris, wood, and metal strips (fig 2). Extra layers of plaster of Paris are used to connect the metal strips on to the plaster case. The kneeling prosthesis is attached to the leg by leather straps.

Sandals

Thick soled sandals are needed to protect the feet of patients with leprosy who have lost normal sensation. Different types of shoe may be made to suit the individual, but they are all designed to prevent damage from sharp stones and to provide a soft top layer to prevent pressure sores developing. The materials that we use include:

- (1) used car tyres for a firm sole,
- (2) tropical or microcellular rubber as a soft protective insole,

FIG 1—Crutches and walking aids.

FIG 2—Prosthetic walking aids.

- (3) leather straps (or straps made of the inner tube of a car),
- (4) buckles and rivets (when rivets are not available the buckles may be stitched on),
- (5) glue,

- (6) nylon or cotton threads to stitch the straps on to the tyre.

The sandals are made as shown in figure 3. In brief, the patient puts his foot on a tropical or microcellular rubber sheet and the foot size is drawn and cut out. Leather straps are passed through slits cut in the microcellular rubber and stitched on firmly. The sole of microcellular rubber is then glued on to a broad piece of car tyre, and once the glue is dry the shape of the sandal is cut out of the tyre. Patients with normally shaped feet can usually be fitted from a stock of standard shoes of varying sizes, but for those who have lost toes or have other deformities each shoe must be custom made to ensure a good fit. (Nails should never be used for they may cause pressure sores.) This type of sandal is very cheap to produce and usually lasts for about a year.

Department of Physiotherapy, Kasama, Zambia

KO DE RUÏTER
OTTO LELIEVELD

Correspondence to: Mr Ko de Ruïter, Casilla 3718, Cochobamba, Bolivia.

FIG 3—Simply made sandals.

Calipers

Above and below knee calipers are used to control movement of the leg. They may be adapted to prevent motion, to limit joint motion to the normal range or a portion of the normal range, to correct or prevent the development of deformity, and, finally, to compensate for muscle weakness. We have found that splints and calipers are especially useful in treating patients with acute poliomyelitis, postpolio paralysis, cerebral palsy, and various orthopaedic conditions such as clubfoot and knock knees. The above knee caliper (fig 4) may be made with simple materials and tools and is cheap, strong, and easy to repair. It is based on the caliper that was designed by Dr Huckstep in Uganda in the 1960s.

Production of a below knee caliper is similar to that of an above knee caliper, the only difference being that the ring is horizontal under the knee. Both types of caliper should be moulded around the ankle if the patient has a fixed varus or valgus deformity. A piece of leather placed anteriorly on the knee is used to keep the knee in 0° of extension or in as much extension as is possible. Two square pieces of leather are cut in sizes ranging from 6×6 cm to 13×13 cm and sewn together. One piece is cut from soft leather and the second from stronger leather

with a round hole cut in the middle to prevent too much pressure on the patella. A posterior knee leather is used to support an unstable knee and is useful in genu recurvatum. The knee is put in 0° of extension when the caliper is applied and measurements are then taken. The strap should be 5–7 cm broad. With the same materials a knock knee strap may be made to prevent or correct a valgus deformity of the knee. A square piece of leather, about

FIG 5—Additional shoe support.

10×10 cm, is used with two straps on top with a buckle on one side. This side of the strap should be long enough to avoid pressure on the skin. An ankle T strap, placed laterally or medially, may be used to prevent or correct a varus or valgus position of the ankle. Finally, a leather ankle strap is necessary to hold both ends of the caliper into the shoe.

A caliper may be fitted into a normal shoe with a rubber heel by drilling two holes in the sides of the heel of the shoe. If the patient cannot afford a shoe a clog is made out of wood with some leather strappings on top. Sometimes a normal shoe does not give enough support and it will be necessary to adapt the shoe into a boot. This may be done by cutting out a piece of leather and stitching it on to the shoe (fig 5). A drop foot may be controlled by a back check stop: a metal pipe in front of a bar is bent and flattened at the ends, which stand vertically. The caliper

FIG 4—Construction of calipers.

goes into the pipe, and the ends of the flattened bar permit dorsiflexion but not plantar flexion. Patients with polio may have shortening of the affected legs and need a shoe raise, which may be made out of layers of tropical sheets glued together with a final layer of car tyre.

The items described above have all been made and put to practical use in a district hospital. Furthermore, I have found that these simple, durable aids serve their purpose equally as well as their western counterparts.

Recommended reading

Simple Orthopaedic Aids by Chris Dartnell. This covers how to set up a workshop and gives full details and simple drawings of the production of orthopaedic aids using simple materials. It is available from the Leonard Cheshire Foundation International, Leonard Cheshire House, 26-29 Maunsel Street, London SW1P 2QN, price £2.50 plus p and p.

Poliomyelitis, a guide for developing countries including appliances and rehabilitation for the disabled by Dr Huckstep. Churchill Livingstone.

Footwear manual on leprosy. Available from the London leprosy mission (contact Jean Watson).

Clinical Algorithms

Generalised pruritus

ROBERT H CHAMPION

I can still remember the misery of itching with a combined attack of chickenpox and German measles and after 36 hours of sleepless torment the itching coming through the euphoria of 30 mg morphine. Many would hardly believe how itching can dominate life. It is often the symptom that brings patients with skin disorders to their doctors. Almost half of all patients with rashes complain of itching at some time, even those suffering from disorders like psoriasis or pityriasis rosea that are said not to itch. We are all taught that generalised itching without primary lesions (and distinguishing primary lesions from the secondary ones produced by scratching is not always easy) should ring an alarm bell in our minds. The first thing to consider when a patient complains of generalised itching without any rash is whether a medically trained eye can detect some subtle change not noticed by the patient. This too may be harder than it sounds. Most doctors must have gratified a colleague by missing what later emerged as typical scabies. The commonest type of minimal rash to present with itching is the drying out of the skin, usually called asteatotic eczema but also sometimes called senile pruritus, a term that means different things to different people. Jumping to the diagnosis of senile pruritus because the patient is old is almost as hazardous as diagnosing menopausal pruritus because the patient is 50 or psychogenic pruritus because the patient admits to some stress.

Having established that generalised itching is not associated with any detectable change in the skin, it is next essential to exclude a systemic cause. This may be a well established and obvious disorder such as chronic renal failure or obstructive jaundice. Often, however, patients with Hodgkin's disease, polycythaemia, or iron deficiency present with generalised itching. The role of zinc deficiency as a cause of itching cannot yet be evaluated. Diabetes is often included in lists of such causes, but the association is relatively rare. Diabetes may, however, often present as anogenital pruritus and candidiasis.

A simple battery of tests can easily be devised to exclude most systemic causes of itching (see algorithm). About half the patients with generalised itching without primary skin lesions eventually get better without any other diagnosis than idiopathic generalised pruritus. Prolonged observation may, however, be needed to exclude, for example, Hodgkin's disease or biliary cirrhosis.

Treatment

Treatment can be fairly simple if there is a diagnosable and removable cutaneous or systemic cause. All too often such common diagnoses as eczema defy treatment and still require symptomatic treatment for itching. Topical corticosteroids can be effective when appropriately applied to inflammatory skin disorders. The result tends to be disappointing when they are applied to itching skin without any lesions. If there is no gross rash a topical emollient such as aqueous cream may provide as good relief as anything. Calamine lotion tends to be too drying if used for more than a day or two. Active ingredients applied to control itching tend to be disappointing. Crotamiton (Eurax) may be useful. Topical local anaesthetics and antihistamines are best avoided.

There are no specific drugs that will do for itch what opiates and analgesics will do for pain, though antihistamines are often prescribed with this aspiration. The newer, peripherally acting antihistamines such as terfenadine and astemizole are valuable for disorders such as urticaria, in which histamine may play a part, but can be expected to be little more than placebos when a central effect is needed. A centrally acting, more sedative antihistamine such as trimeprazine or promethazine hydrochloride (Phenergan) may then be more successful. H_2 antagonists such as cimetidine have been used with success in small numbers of patients with itching due to several systemic causes but have often failed in other trials. So far the indications for cimetidine in the general management of itching have not been adequately assessed. It is certainly not a panacea.

Another fairly simple empirical remedy for many types of generalised itching with normal skin is ultraviolet radiation. The mode of action is not certain.

Some patients, perhaps surprisingly, need advice about not getting too hot, not having very hot baths, wearing comfortable loose clothing, avoiding harsh clothes next to the skin, and taking care with alcohol. Itching is, moreover, a symptom that may be alleviated by any measure that encourages the patient to relax. Thus various therapeutic possibilities are raised, from placebos to tranquillisers, hypnotherapy, acupuncture, and many others. The use of these must depend on the experience and inclinations of the practitioner.

Further reading

Beare JM. Generalised pruritus: a study of 43 cases. *Clin Exp Dermatol* 1976;1:343-52.
Champion RH. Pruritus. *Medicine International* 1983;1:252.
Cunliffe WJ. The skin and the nervous system. In: Rook AJ, Wilkinson DS, Ebling FJ, eds. *Textbook of dermatology*. Oxford: Blackwell Scientific, 1978:1993-2008.
Herdson JJ. Itching: the pathophysiology of pruritus. *Int J Dermatol* 1975;14:465-84.