in which the incidence of the effect would be measured so that a careful benefit-risk assessment could be formed, taking account of information on other similar drugs. Resources giving rapid and accurate assessments on these lines would allow better decisions about products by both companies and regulatory authorities.

Perhaps the most important conclusion of the Minster Lovell seminar was that an automated national data base linking prescriptions filled with hospital diagnoses in Britain would be extremely valuable in allowing the rapid study of drug effects and providing quantitative data. This would lessen public anxiety and restore much needed confidence. The Prescription Pricing Authority and the National Health Service hospital system provide an opportunity for such a system of drug surveillance based on computerised linkage of prescriptions with hospital diagnoses. A start has been made in the Dundee record linkage study³ and the cimetidine postmarketing surveillance study.⁴ Confidentiality should not be allowed to be an insuperable problem, and the Minster Lovell seminar also concluded that technical and organisational barriers could be overcome—provided that enough individuals were encouraged to make a commitment to research in pharmacoepidemiology.

D H LAWSON

Consultant Physician,
Royal Infirmary,
Glasgow G4 0SF

- 1 Turner P. Long-term assessment of drug safety and efficacy. *J R Soc Med* 1984;77:93-4.
- 2 Rawlins MD. Postmarketing surveillance of adverse reactions to drugs. *Br Med J* 1984;288:879-80.
- 3 Walker SR, Goldberg A, eds. *Monitoring for adverse drug reactions*. Lancaster: MTP Press, 1984.
- 4 Colin-Jones DG, Langman MJS, Lawson DH, Vessey MP. Postmarketing surveillance of the safety of cimetidine: 12 month mortality report. *Br Med J* 1983;286:1713-6.
- 5 Inman WHW. Postmarketing surveillance of adverse drug reactions in general practice. II. Prescription monitoring at the University of Southampton. *Br Med J* 1981;282:1216-7.
- 6 Mann JI, Vessey MP, Thorogood M, Doll R. Myocardial infarction in young women with special reference to oral contraceptive practice. *Br Med J* 1975;ii:241-4.
- 7 Lawson DH, Henry DA, Lowe JM, et al. Severe hypokalemia in hospitalised patients. *Arch Intern Med* 1979;139:978-80.
- 8 Skegg DCG, Doll R. Frequency of eye complaints and rashes amongst patients receiving practolol and propranolol. *Lancet* 1977;ii:475-8.
- 9 Jick H, Watkins RN, Hunter JR, et al. Replacement estrogens and endometrial cancer. *N Engl J Med* 1979;300:218-22.
- 10 Jick H. Drugs—remarkably non-toxic. *N Engl J Med* 1974;291:824-8.
- 11 Inman WHW. Risks in medical intervention: balancing therapeutic risks and benefits. Oxford: University of Oxford, 1984. (Wolfson lecture, 1984.)

antibody plays a part in this vulnerability to infection but is by no means the whole explanation. Indeed, the picture is further confused by the occurrence of arthritis,⁴ meningitis,⁵ septicaemia,⁶ and pneumonia^{7,8} due to *H influenzae* in older children and in adults.

Septic arthritis usually affects the larger joints such as the knee, ankle, and hip, and occasionally those of the arm.⁹ The features are those of an acute pyoarthrosis with pain (which may be referred), limitation of movement, swelling, and variable erythema. Fever is usual, sweating common, and rigors an occasional accompaniment. The younger child may be unwell in a non-specific way and fractious. Reluctance to use a limb is an important clue. An infected purulent exudate at arthrocentesis confirms the diagnosis. Osteomyelitis of the adjacent bone is rare, as is spread to other joints. Rest, analgesics, antibiotics, and judicious arthrocenteses summarise the management.

H influenzae may cause a cellulitis that usually affects the cheek, occasionally the periorbital area, and may follow a coryzal prodrome.^{10,11} The area is raised, warm, and tender and may be purplish rather than red. Fever, sweating, and generalised toxicity are common. The anatomical distribution and purplish colour provide a clue to the nature of this infection, which must be differentiated from streptococcal pyoderma, trauma, and the slapped cheek syndrome, otherwise known as erythema infectiosum or fifth disease.^{12,13}

The true incidence of pneumonia caused by encapsulated strains of *H influenzae* is unknown, since in the absence of bacteraemia this infection can rarely be diagnosed with confidence by sputum culture. None the less, *H influenzae* is well described as an occasional cause of pneumonia in adults,^{7,8,14,15} and accounted for 3% of infections in a recent study of adult pneumonia in Britain, while 2% of the bacteraemic infections in the Finnish study complicated pneumonia.¹ Pneumonia in early childhood may be caused more frequently than has been thought by *H influenzae* according to recent antigen detection studies.¹⁶

The heart is rarely the target for *H influenzae* infection; pyogenic pericarditis may occur in isolation or complicate respiratory infection, meningitis, or epiglottitis,¹⁷ and endocarditis occurs uncommonly and has no distinguishing features.¹⁸ Other less common infections include conjunctivitis, enophthalmitis, subcutaneous abscess, peritonitis, glossitis, thyroiditis, laryngotracheobronchitis without epiglottitis, and biliary tract infections.

Laboratory diagnosis of *H influenzae* infections is not difficult, although its pleomorphic Gram negative staining characteristics may make it difficult to visualise against a background of pus. Type b capsular polysaccharide may be identified with potent antiserum. Rapid methods for detecting this capsular polysaccharide are gaining in popularity and may have a bearing on the early management of these infections. Various body fluids such as serum, cerebrospinal fluid, and urine have been tested for type b capsular antigen by counterimmunoelectrophoresis, coagglutination, and latex agglutination.^{16,19}

Treatment of *H influenzae* infections has been made more difficult with the increasing resistance to ampicillin. A recent survey has shown that some 6.2% of isolates in Britain were resistant, though there was considerable geographical variation.²⁰ Chloramphenicol has well known undesirable properties, and resistance among *H influenzae*, though still uncommon, is well recognised.^{21,22} Drugs such as the injectable cephalosporins seem likely to be used more frequently. Among those that are highly active against

Uncommon haemophilus infections

Haemophilus influenzae is a familiar cause of meningitis and epiglottitis—two diseases which present dramatically and, untreated, threaten survival. Otitis media, sinusitis, and infective exacerbations of chronic bronchitis are also frequently caused by *H influenzae*, though non-encapsulated strains predominate in chest infections. What is less well known is the range of other infections that may be caused by this organism. A recent review from Finland has shown that in 17% of patients with bacteraemia with *H influenzae* the infection was in sites other than the meninges and epiglottitis.¹ The joints, bones, eyes, skin, parotids,² the genitourinary tract,³ and the heart valves need to be added to the list. Invasion of the blood stream is generally secondary to infection in these sites, but it may occur in isolation.

Infections caused by encapsulated *H influenzae* occur predominantly in early childhood. Septic arthritis, cellulitis, and bacteraemia tend to affect children under 2 years of age, while epiglottitis and pneumonia are more frequent in those between 2 and 5. The nadir of passively acquired maternal

H influenzae, including β -lactamase producing strains, are cefuroxime, cefotaxime, latamoxef, and ceftazidime. Experience with cefuroxime in treating *H influenzae* meningitis suggests that it should be a suitable alternative to treat such infections.²³

Case clustering studies and epidemiological surveys have underlined the communicability of *H influenzae* infection.^{24,25} In the United States whenever meningitis occurs chemoprophylaxis with rifampicin is now recommended for the index patient, and households with children below 4 years of age, as well as for staff and children in day centres.²⁶ The evidence that this is necessary for infections other than meningitis is lacking, though secondary cases of *H influenzae* infection may complicate non-meningitic primary infections.²⁷ Prevention through immunisation remains an attractive hope, with several vaccines as candidates, though this leap into the realms of practical preventive medicine still appears distant.²⁸⁻³⁰

R G FINCH

Consultant and Senior Lecturer in Microbial Diseases,
City Hospital and University of Nottingham,
Nottingham NG5 1PB

- 1 Peltola H, Virtanen M. Systemic *Haemophilus influenzae* infection in Finland. *Clin Pediatr* 1982;23:275-80.
- 2 Fainstein V, Musher DM, Young EJ. Acute bilateral suppurative parotitis due to *Haemophilus influenzae*. *Arch Intern Med* 1979;139:712-3.
- 3 Albritton WL, Hammond GW, Ronald AR. Bacteremic *Haemophilus influenzae* genitourinary tract infections in adults. *Arch Intern Med* 1978;138:1819-21.
- 4 Raff MJ, Dennaher CL. *Haemophilus influenzae* septic arthritis in adults. *J Bone Joint Surg* 1984;56A:408-12.
- 5 Eykyn SJ, Thomas RD, Phillips I. *Haemophilus influenzae* meningitis in adults. *Br Med J* 1974;ii:463-5.

- 6 Farrand RJ. *Haemophilus influenzae* type b septicaemia. *Br Med J* 1969;i:150-2.
- 7 Levin DC, Schwarz MI, Matthay RA, LaForce FM. Bacteremic *Haemophilus influenzae* pneumonia in adults. *Am J Med* 1977;62:219-24.
- 8 Macfarlane JT, Finch RG, Ward MJ, Macrae AD. Hospital study of adult community-acquired pneumonia. *Lancet* 1982;ii:255-8.
- 9 Fink CW, Dick VQ, Howard J, Nelson JD. Infections of bones and joints in children. *Arthritis Rheum* 1977;20(suppl):578-83.
- 10 Rasmussen JE. *Haemophilus influenzae* cellulitis. *Br J Dermatol* 1973;88:547-50.
- 11 Pedlar SJ, Hawkey PM. Cellulitis in children caused by *Haemophilus influenzae* type b. *Journal of Infection* 1983;6:269-72.
- 12 Cramp HE, Armstrong BDJ. Erythema infectiosum: an outbreak of "slapped cheek" disease in north Devon. *Br Med J* 1976;ii:885-6.
- 13 Mortimer PP. The 80th year of fifth disease. *Br Med J* 1984;289:338-9.
- 14 Wallace RJ, Musher DM, Martin RR. *Haemophilus influenzae* pneumonia in adults. *Am J Med* 1978;64:87-92.
- 15 Everett ED, Rahm AE, Adaniya R, Stevens DL, McNitt TR. *Haemophilus influenzae* pneumonia in adults. *JAMA* 1977;238:319-21.
- 16 Turner RB, Hayden FG, Hendley JO. Counterimmunoelectrophoresis of urine for the diagnosis of bacterial pneumonia in pediatric out-patients. *Pediatrics* 1983;71:780-3.
- 17 Echeverria P, Smith EWP, Ingram D, Sade RM, Gardner P. *H influenzae* b pericarditis in children. *Pediatrics* 1975;56:808-18.
- 18 Bayliss R, Clarke C, Oakley CM, Somerville W, Whitfield AGW, Young SEJ. The microbiology and pathogenesis of infective endocarditis. *Br Heart J* 1983;50:513-9.
- 19 Tompkins DG. Comparison of Phadebact coagglutination tests with counterimmunoelectrophoresis for the detection of bacterial antigens in cerebrospinal fluid. *J Clin Pathol* 1983;36:819-22.
- 20 Philpott-Howard J, Williams JD. Increase in antibiotic resistance in *Haemophilus influenzae* in the United Kingdom since 1977: report of a study group. *Br Med J* 1982;284:1597-9.
- 21 Ward JI, Tsai TF, Felice GA, et al. Prevalence of ampicillin- and chloramphenicol-resistant strains of *Haemophilus influenzae* causing meningitis and bacteraemia: national survey of hospital laboratories. *J Infect Dis* 1978;138:421-4.
- 22 Macmahon P, Sills J, Hall E, Fitzgerald T. *Haemophilus influenzae* type B resistant to both chloramphenicol and ampicillin in Britain. *Br Med J* 1982;284:1229-30.
- 23 Swedish Study Group. Cefuroxime versus ampicillin and chloramphenicol for the treatment of bacterial meningitis. *Lancet* 1982;ii:295-9.
- 24 Ward JI, Fraser DW, Baraff LJ, Plikaytis BD. *Haemophilus influenzae* meningitis. A national study of secondary spread in household contacts. *N Engl J Med* 1979;301:122-6.
- 25 Glode MP, Daum RS, Goldman DA, Leclaire J, Smith A. *Haemophilus influenzae* type b meningitis: a contagious disease of children. *Br Med J* 1980;280:899-901.
- 26 Anonymous. Prevention of secondary cases of *Haemophilus influenzae* type b disease. *MMWR* 1982;31:672-80.
- 27 Fillice GA, Andrews JS Jr, Hudgins MP, Fraser DW. Spread of *Haemophilus influenzae*. *Am J Dis Child* 1978;132:757-9.
- 28 Schneerson R, Robbins JB. Induction of serum *Haemophilus influenzae* type b capsular polysaccharide antibodies in adult volunteers fed cross-reacting *Escherichia coli* 075:K100:H5. *N Engl J Med* 1975;292:1093-6.
- 29 Peltola H, Katthy H, Sinoven A, Makela H. *Haemophilus influenzae* type b capsular polysaccharide vaccine in children: a double-blind field study of 100,000 vaccinees 3 months to 5 years of age in Finland. *Pediatrics* 1977;60:730-7.
- 30 King SD, Wynter H, Ramal A, et al. Safety and immunogenicity of a new *Haemophilus influenzae* type b vaccine in infants under one year of age. *Lancet* 1981;ii:705-9.

Shredding of rejected manuscripts

Every year the *BMJ* receives for publication well over 5000 articles, of which over 4500 are rejected. As other journals have found, the expense in handling these is considerable: five years ago Arnold Relman estimated that the *New England Journal of Medicine* spent about \$50 per paper in registering the manuscript, sending it out to two referees, and returning it to the author if it was rejected for publication, and our own calculations suggest a similar figure. Other journals, particularly in the United States, have sought to transfer this considerable burden on to the authors' shoulders: they have introduced handling charges for all manuscripts (currently around \$50 per paper for several journals), or they make page charges for those articles accepted for publication. Such charges are another drain on already sparse research funds, and we do not believe that they should be applied save in great financial difficulties. Nevertheless, we can no longer go on handling manuscripts in the traditional way, and hence from 1 January 1985 we intend not to return those rejected for publication but to destroy them.

This new scheme will work as follows. An acknowledgment will be sent on the receipt of each manuscript, reminding authors that this will not be returned if it is rejected for publication. To deal with any inquiries manuscripts that are rejected will be held for three months before being shredded and complete confidentiality will be ensured. We will still require three copies of each manuscript, to allow one to be sent to each referee and for the editor to retain one for reference. For those articles that are rejected a

letter will be sent to the author advising him of this decision and also stating that the manuscript will be destroyed.

Some might see this as yet another manifestation of editorial "uppitiness." We would resist such a claim strongly: journals such as our own thrive on publishing original research work, and we welcome articles for publication. Nevertheless, the present system is throwing a heavy strain on the office, and tending to concentrate on processing the rejected article rather than improving the accepted one. If such a scheme has the effect of diverting articles that should properly have been sent to specialist journals in the first place then so much the better, but we hope that authors will accept it in a constructive way. Of 30 advisers and referees consulted, only one was against the scheme, on the grounds that it might be more trouble than it was worth, but, given that we intend to introduce it initially on a pilot basis, we should readily be able to see whether he was right or not. Now that most people have access to photocopying machines, and often to word processors, the scheme should cause relatively little difficulty, and we will make exceptions—for example, for manuscripts coming from the Third World or when original artwork is enclosed—though we would much prefer initially to receive three copies of manuscripts and illustrations prepared by photocopier or word processor.

Given all these safeguards, then, we hope that the scheme will help to obviate a lot of difficulties.

STEPHEN LOCK
Editor, *British Medical Journal*