contraception in teenagers. The prevention of pregnancies possible in this high risk group, however, would amply justify the increased time and effort. If those of us working with this group are not prepared to offer contraceptive advice and help when it is needed, then we are indeed failing our teenagers.

Thanks are due to Mrs G M Buick, Fareham family planning clinic secretary, for her help with the records and Dr M Laxton for allowing access to Southampton family planning clinic records.

References

- 1 Jamieson L, Bury JK, McGlew T. Young persons' contraceptive clinic 1975 to 1980: patterns of use. *Health Bull (Edinb)* 1983;41:68-77.
- 2 Chief Medical Officer, DHSS. Sexually transmitted diseases; extract from annual report for 1982. *Br J Vener Dis* 1984;60:199-205.

- 3 Office of Population Censuses and Surveys. Legal abortions 1983. *OPCS monitor*. London: HMSO, 1984. (AB 84/6.)
- 4 Office of Population Censuses and Surveys. Live births during 1983 by mother's age, legitimacy and birth order. *OPCS monitor*. London: HMSO, 1984. (FMI 84/4.)
- 5 Guillebaud J. Minimising the hazards of contraception in the under-16s. *Modern Medicine* 1980;25:38-40.
- 6 Lowry TP. First coitus. *British Journal of Sexual Medicine* 1982;9:31-3.
- 7 Harrison N, Bury JK. A young people's advisory service. *Health Bull (Edinb)* 1982;4:133-9.
- 8 Charnie M, Eisman S, Forrest JD, et al. Factors affecting adolescents' use of family planning clinics. *Fam Plann Perspect* 1982;14:126-7,131-9.
- 9 Jones JB, Namerow PB, Philliber S. Adolescents' use of a hospital based contraceptive program. *Fam Plann Perspect* 1982;14:224-5,229-31.
- 10 Zabin LS, Kantner JF, Zelnik M. The risk of adolescent pregnancy in the first months of intercourse. *Fam Plann Perspect* 1979;11:215.
- 11 Zabin LS, Clark SD. Why they delay—a study of teenage family planning clinic patients. *Fam Plann Perspect* 1981;13:205-7,211-7.
- 12 Zelnik M, Shah FK. First intercourse among young Americans. *Fam Plann Perspect* 1983;15:64-70.
- 13 Zelnik M, Young JK. Sex education and its association with teenage sexual activity, pregnancy and contraceptive use. *Fam Plann Perspect* 1982;14:117-9,123-6.
- 14 Forrest JD, Hermalin AI, Henshaw SK. The impact of family planning clinic programs on adolescent pregnancy. *Fam Plann Perspect* 1981;13:109-16.

(Accepted 19 October 1984)

Clinical Algorithm

Hypercalcaemia

M J JAMIESON

Four key questions are central to the assessment of the apparently hypercalcaemic patient. As assessment evolves they arise in the following order: Is this true hypercalcaemia? Is urgent treatment required? What is the cause? What treatment, if any, is required electively?

In deciding whether a single estimation of serum calcium concentration reflects true sustained hypercalcaemia, three factors should be considered: firstly, a normal person may have a high serum calcium concentration on a given occasion (compared with his laboratory's reference range) solely as a chance occurrence; secondly, measurements of serum calcium concentration generally include a protein bound fraction of 40-45%, of which about 90% is bound to albumin. To assess whether measured serum calcium includes a physiologically appropriate free calcium concentration a correction must be made for bound calcium. Estimates of average calcium-albumin binding derived from regression equations are widely used, and popular correction factors are 0.025 mmol (1 mg) calcium/g albumin and 0.1 mmol (4 mg) calcium/6 g albumin. In either case an appropriate amount is subtracted from or added to total calcium according to the amount by which the serum albumin concentration exceeds or falls short of a normal concentration (usually around 40 g/l). For example, using the first of these correction factors, a patient whose serum calcium concentration is 2.80 mmol/l (11.2 mg/100 ml) and serum albumin concentration is 50 g/dl has a corrected serum calcium concentration of 2.55 mmol/l (10.2 mg/100 ml)— $2.80 - ((50 - 40) \times 0.025)$. A patient whose serum calcium concentration is 2.00 mmol/l (8 mg/100 ml) and serum albumin concentration is 30 g/dl has a corrected calcium concentration of 2.25 mmol/l (9 mg/100 ml)— $2.00 + ((40 - 30) \times 0.025)$.

The application of average correction values is somewhat empirical as, strictly, they should be applied only to members of the populations from which they are derived and, more particularly, as individual variation in calcium-albumin binding appears to be considerable. Most studies, nevertheless, have been able to assign a diagnosis in most cases without defining individual regression coefficients. Thirdly, sampling variables should be excluded; serum calcium concentrations in samples taken standing, after venous occlusion, and postprandially may exceed the supine, tourniquet free, and fasting values by as much as 0.25 mmol/l (1 mg/100 ml) in each case.

TABLE 1—Causes of hypercalcaemia. Figures in parentheses represent typical incidences in hospital series

Type of cause	Cause
Common	Malignancy (55%)
Uncommon (less than 10%)	Hyperparathyroidism (35%)
	Renal dialysis and transplantation
	Thyroiditis
	Sarcoidosis
Rare (less than 1%)	Vitamin D excess
	Milk-alkali syndrome
	Thiazide diuretics
	Lithium
	Familial hypocalcaemic hypercalcaemia
	Immobilisation (eg in Paget's disease)
	Addison's disease
	Acromegaly
	Phaeochromocytoma
	Cystic breast disease
	Coccidioidomycosis
	Vipoma (vasoactive intestinal polypeptide tumour)

Causes

Table 1 gives a list of causes. In an unselected population the commonest cause will probably be primary hyperparathyroidism; in hospital series malignancy is commoner. Together these account for about 90% of cases. Other causes are individually uncommon and when present may coexist with hyperparathyroidism.

Department of Therapeutics and Clinical Pharmacology, Royal Infirmary, Aberdeen AB9 2ZB

M J JAMIESON, MRCP, lecturer

Correspondence to: Division of Clinical Pharmacology, University of Texas Health Service Center, San Antonio, Texas, 78284, USA.

Presentation

The commonest presentation is as a coincidental finding on biochemical screening. Non-specific symptoms including anorexia, weight loss, and fatigue are common, particularly when there is underlying malignancy. Symptoms more usually associated with hypercalcaemia include constipation, bone pain, thirst and polyuria, and mental changes. These do not discriminate between malignancy and hyperparathyroidism and do not correlate well with serum calcium concentration. Renal calculi are present in about one fifth of cases of hyperparathyroidism but are rare in association with malignancy. Hypertension, peptic ulcer, and acute pancreatitis are occasional presentations. Acute hypercalcaemia (with dehydration and confusion) tends to develop in older patients and to indicate malignancy. Malignancy becomes increasingly unlikely with longer histories. A positive family history may be obtained in hyperparathyroidism, familial hypocalciuric hypercalcaemia, and in the uncommon syndromes of multiple endocrine neoplasia.

Signs of hypercalcaemia are few; ectopic subcutaneous, corneal, arterial, and cardiac calcification are typically seen (or heard) in association with chronic renal failure and after phosphate treatment.

In practice, most cases of malignancy are apparent by the time hypercalcaemia has developed. It is unusual for occult malignancy to present as hypercalcaemia in an otherwise fit patient. Bronchogenic and breast carcinomas account for about 45% of cases, the remainder largely comprising tumours of the renal tract, ears, nose, and throat, female genital tract, and oesophagus and multiple myeloma. More than three quarters of these patients will have detectable metastases at the time of presentation (with the exception of those with urothelial tumours, about a third of whom will have metastases).

The acute milk-alkali syndrome, as described in the 1920s, is now very rare. A rather more insidious type of this syndrome may occur, particularly in elderly patients, and can easily be missed unless a history of excessive intake of calcium and alkali is specifically sought.

Investigation

History, examination, and baseline investigations as outlined in the algorithm will detect most cases of malignant disease and will identify most patients with unusual causes of hypercalcaemia. Simple analysis of the biochemical findings at this stage gives useful guidance but does not discriminate between hyperparathyroidism and other diagnoses. For example, more than three quarters of patients with serum calcium concentration over 4.0 mmol/l (16 mg/100 ml) will have malignancy; a serum phosphate concentration higher than the upper normal limit or an alkaline phosphatase activity greater than twice the upper limit of normal makes primary hyperparathyroidism unlikely. The ratio of serum chloride to phosphate appears to be useful in classifying patients into parathyroid and non-parathyroid groups (a ratio of less than 33, for example, makes primary hyperparathyroidism unlikely) as is discriminant function analysis using, for example, calcium, chloride, and parathyroid hormone concentrations and alkaline phosphatase activities.

Assay of immunoreactive parathyroid hormone is potentially the simplest discriminatory test, and indeed appreciably raised parathyroid hormone concentrations confirm hyperparathyroidism while undetectable concentrations are inconsistent with this diagnosis. Normal or moderately increased concentrations (which in the face of hypercalcaemia would appear to imply parathyroid autonomy) may be found, however, in various conditions including malignancies and familial hypocalciuric hypercalcaemia. These may be associated with ectopic or inappropriate production of parathyroid hormone but do not respond well to parathyroid surgery. Parathyroid hormone assay is best restricted to two situations: (1) when history and baseline investigations suggest hyperparathyroidism; (2) when these have given no clues to a diagnosis.

Most patients at this stage will ultimately prove to have hyperparathyroidism, in which case it is legitimate to arrange the following supportive investigations while awaiting the results of tests of parathyroid hormone concentrations.

(1) X ray examination of hands (looking in particular for subperiosteal erosions) is likely to be normal (especially if the alkaline phosphatase activity is normal) but may provide useful evidence of parathyroid bone disease. Its value in screening for causes of hypercalcaemia is limited; skull radiographs are unnecessary.

(2) Intravenous urography may occasionally detect an otherwise occult hypernephroma but should be restricted to showing stones in symptomatic patients and to detecting nephrolithiasis and nephrocalcinosis in patients with established hyperparathyroidism. Its use at this stage is, however, probably justified.

(3) Hypercalciuria is of limited diagnostic value, although extremely high values (for example, >12.5 mmol (500 mg)/24 h) may be found in sarcoidosis, vitamin D intoxication, and some malignancies. It has, however, prognostic value in hyperparathyroidism and may therefore influence treatment (see management). Relatively low values of urinary calcium excretion suggest familial hypocalciuric hypercalcaemia, but as a screen for this uncommon disorder urinary calcium measurement is probably best reserved for those at higher risk—that is, patients with a history of failed parathyroidectomy and patients with first degree relatives known to have familial hypocalciuric hypercalcaemia. As familial hypocalciuric hypercalcaemia may mimic hyperparathyroidism it is also reasonable to assess urinary calcium excretion when hyperparathyroidism appears to be the diagnosis, and here, as an overlap exists in urinary calcium excretion between familial hypocalciuric hypercalcaemia and hyperparathyroidism, the ratio of calcium clearance to creatinine clearance (which discriminates well) should be expressed.

After parathyroid hormone assay a small group of patients will still be left without a diagnosis. Most of these will, again, ultimately prove to have hyperparathyroidism, with occult malignancy the main differential diagnosis. The options are to review and repeat earlier examination and investigations, including parathyroid hormone assay (in the hope that time will aid diagnosis) or to proceed with investigations (including, potentially, parathyroid surgery) aimed at establishing or excluding hyperparathyroidism. Investigations that may be considered include steroid suppression and calcium infusion tests. Neither of these, unfortunately, reliably discriminates between hyperparathyroid and non-parathyroid hypercalcaemia or between individual non-hyperparathyroid causes. Nevertheless, lack of significant suppression of serum calcium concentration by hydrocortisone and of urinary phosphate excretion by intravenous calcium may be useful evidence in favour of hyperparathyroidism.

Serum and urinary protein electrophoresis should be carried out along with serum thyroxine estimation. Suspicion of hyperthyroidism despite thyroxine concentrations within the normal range will be allayed by normal serum triiodothyronine concentrations and if necessary by normal thyroid stimulating hormone response to thyrotrophin releasing hormone. Investigations aimed at diagnosing rare causes (such as the other endocrine disorders listed) when these are not clinically apparent are unlikely to give positive results. Isotopic bone scanning (which is more sensitive than radiological skeletal survey in detecting early bony deposits) is reasonably indicated now if no other diagnosis is apparent. Parathyroid localising studies (including grey scale ultrasonography, computed tomography, arteriography, and selective venous sampling) should be reserved for cases of known hyperparathyroidism requiring surgery; the invasive studies are best restricted to cases where neck exploration has been unsuccessful and reoperation is being considered.

A group of patients will remain in which the diagnosis is "hypercalcaemia, cause unknown" or "hypercalcaemia, probably hyperparathyroidism." Parathyroid surgery may be considered in these patients; the decision to carry out neck exploration will be influenced primarily by how strongly hyperparathyroidism is suspected. Other factors that will influence this decision in

Hypercalcaemia

established or strongly suspected hyperparathyroidism are considered below.

Management

The algorithm outlines management. Indications for surgery of hyperparathyroidism are not absolute but include bone disease, appreciable hypercalciuria, impaired renal function, recurrent renal stones, peptic ulceration, hypertension, pancreatitis, psychosis, and persistently raised serum calcium concentration (>3.0 mmol/l (12 mg/100 ml) being a typical, somewhat arbitrary, figure). Other factors that may be considered are that surgery usually provides a definite tissue diagnosis, is curative in about 90%, simplifies follow up (and reduces the risks to patients lost to follow up), and, in experienced hands, carries a low risk to life and of postoperative complications. (In one series of 500 consecutive cases, for example, the hospital mortality was 0.2%, incidence of vocal cord paralysis 0.2%, and incidence of prolonged hypoparathyroidism 2.0%.)

The course of untreated mild hyperparathyroidism appears to be similar to that following successful parathyroid surgery; there does not appear to be any adverse effect on renal function or blood pressure, and it is reasonable to manage patients aged over 60 without evidence of hypertension, renal impairment, renal calculi, or bone disease conservatively. Younger patients should be referred for surgical advice. The use of oral phosphate carries a risk of metastatic calcification and should be restricted to patients who would benefit from but are unfit for surgery (or in whom surgery has failed to correct hypercalcaemia). Cyclical oestrogen treatment may be considered for postmenopausal women in the same category.

Management of established hypercalcaemia in association with chronic renal failure and dialysis should be undertaken by or in conjunction with a nephrologist. Reversal of phosphate retention is central and vitamin D sterols (such as alfalcidol) may be useful. Parathyroid surgery may be necessary.

The principles of management of acute hypercalcaemia and of symptomatic hypercalcaemia of malignancy are similar; rehydra-

tion is of prime importance, and specific drug treatment is usually reserved for those who remain hypercalcaemic and symptomatic, or both, despite adequate fluid replacement. The popularity of the loop diuretics varies from centre to centre and overall appears to be declining. None of the second line drugs is ideal; of these, a combination of calcitonin and corticosteroids appears currently to be the treatment of choice, although the new diphosphonates, particularly disodium aminohydroxypropylidene diphosphonate (APD), are promising in this respect. Table II lists alternatives, their effectiveness, and disadvantages.

TABLE II—Drugs used in acute hypercalcaemia

Drug	Comments
Mithramycin	Effective, but rebound hypercalcaemia, thrombocytopenia, and renal liver damage
Phosphate	Effective, but (when taken by mouth) diarrhoea and (when given intravenously) metastatic calcification
Corticosteroids	Neither particularly effective alone; possibly treatment of choice in combination
Calcitonin	Ineffective
Indomethacin	Disodium etidronate (EHDP) ineffective (osteomalacia with long term treatment)
Diphosphonates	Disodium aminohydroxypropylidenediphosphonate (APD) effective

Peritoneal dialysis or haemodialysis may be necessary when renal or cardiac failure complicates large volume fluid replacement and as a last resort when the above measures have failed. Their use should be considered in the light of the prognosis of the underlying disease.

References

- Goodwin FJ. Symptomless abnormalities: hypercalcaemia. *Br J Hosp Med* 1982;28:54-8.
- Fiskin RA, Heath DA, Somers S, Bold AM. Hypercalcaemia in hospital patients, clinical and diagnostic aspects. *Lancet* 1981;ii:202-7.
- Anonymous. Diagnosis and treatment of primary hyperparathyroidism. *Lancet* 1980;ii:1339-40.
- Wong ET, Freier EF. The differential diagnosis of hypercalcaemia. *JAMA* 1982;247:75-80.
- Heath DA, Marx SJ, eds. *Clinical endocrinology 2. Calcium disorders*. London: Butterworth, 1982.

A fit man of 80 suffers from severe burning pains in the knees under the patella, worse on walking but also present when resting. Radiographs of the knees show minor changes but those of the back show much intervertebral joint derangement of lumbar vertebrae. He also has paraesthesia of the ankles. Arterial function is good and the erythrocyte sedimentation rate and blood examination are normal. What treatment is advised?

In a man of 80 pains in the knees on walking, whatever the description, are probably due to osteoarthritis. In patellofemoral osteoarthritis the x ray changes are often minimal, and the diagnosis is usually clinched by reproducing the pain on patellofemoral compression. Standard treatments include weight reduction, avoidance of precipitating causes, non-steroid anti-inflammatory drugs, and courses of local heat and static quadriceps exercises with the knees in slight flexion. A possible alternative diagnosis might be localised compression neuropathy. The best known of these in the leg affects the anterior branch of the lateral femoral cutaneous nerve of the thigh as it pierces the femoral fascia and gives rise to "meralgia paraesthetica." A similar lesion may affect the anterior femoral cutaneous nerve of the thigh, also as it pierces the fascia. Either may be provoked by local pressure, such as the lower end of a corset, surgical scars, or adipose tissue. The characteristic symptom is burning paraesthesiae, sometimes numbness, and a distribution distinguishable from a lumbar dermatome.

The infrapatellar branch of the saphenous nerve might also be affected as it pierces the fascia, and this would lead to similar symptoms. Sunderland mentions lacerations and surgical incisions damaging this nerve,¹ but repeated local pressure on the inner thighs might also cause the problem. Diagnosis is strengthened by finding a local pressure point that aggravates the symptoms and by obtaining relief from a local anaesthetic/steroid injection into the trigger point. Presumably decompression of the nerve as it emerges through the fascia would be curative. The presence of paraesthesiae of the ankles, however, suggests a generalised sensory neuropathy, in which case the ankle reflexes might be absent and sensory nerve conduction tests

abnormal. "Degenerative" changes in x ray films of the lumbar spine and knees are so common at this age as not to be diagnostic. Nevertheless, reproduction of the symptoms by mechanical stress of the lumbar spine or knees, or relief by local anaesthetic (epidural or intra-articular) injections would lend weight to these sites of origin.—JOHN A MATHEWS, consultant physician (rheumatology), London.

1 Sunderland S. *Nerves and nerve injuries*. 2nd edn. Edinburgh, London and New York: Churchill Livingstone, 1978.

A healthy man in his 70s suffers from osteoarthritis of the ankle relieved by 1800 mg aspirin twice daily. Is this dose safe if continued over an indefinite period?

Aspirin has now largely been replaced in the United Kingdom by the non-steroidal drugs, which are far less toxic to the gastric mucosa and have a far higher compliance (acceptance) record with patients than aspirin. In a 70 year old man with osteoarthritis there is unlikely to be a major inflammatory component. Theoretically, analgesics might be preferred. If, as the question implies, the patient finds aspirin the best choice, then so be it.—G R V HUGHES, consultant physician and reader in rheumatology, London.

Would a daily intake of 200 g of chocolate mints (which consist mainly of icing sugar) be harmful to the cardiovascular system of a man in his 60s?

There is no convincing evidence that dietary sugar is important in the development of coronary artery disease, nor is there any scientific evidence that dietary sugar exhausts the pancreatic islets and predisposes to obesity. Nevertheless, 200 g of icing sugar does provide 700 calories and this will therefore predispose to obesity. As Hippocrates stated 2400 years ago, people who are very fat are apt to die earlier than those who are slimmer.—C W H HAVARD, consultant physician and endocrinologist, London.