Plastic and Reconstructive Surgery

D M DAVIES

SKIN COVER

Partial thickness grafts

A skin graft is a segment of dermis and epidermis that has been completely separated from its blood supply and is transplanted to another area of the body. The first successful skin graft was performed in France in 1869 by Jacques Reverdan. In this country partial thickness grafts (split thickness or Thiersch's grafts) are harvested as large sheets from either the arm, leg, or buttocks using a blade with an adjustable protective roller at the leading edge, which determines the thickness of the skin. As epidermal remnants are left behind in the donor site the skin heals within 10 to 14 days, depending on the thickness of the graft, usually with only minimal pigment change. If a large area of skin loss needs to be covered the partial thickness graft can be meshed using a machine that permits an expansion in the graft rather like a string vest. Skin grafts can be stored in a domestic fridge if kept moist and used up to three weeks later.

The survival of a skin graft depends on the recipient site being sufficiently vascular to support the metabolic needs of the graft and free from gross infection. A successful "take" also requires immobilisation and the prevention of seromas and haematomas, which may lift the graft off the vascular bed. Gross infection mobilises the fibrinolytic system and breaks down the fibrin adhesion between the graft and the recipient bed. Partial thickness grafts may be applied to a wound directly at the time of surgery and the immobilisation and prevention of seroma obtained by the application of a bolus pressure dressing using proflavine wool or thick foam. Grafts may also be applied 24-28 hours after surgery and left exposed for frequent wound toilet and debridement without any tie-over dressing. The main disadvantages are the cosmetic mismatch between the graft and surrounding normal skin in colour, quality, and contour and the tendency of the wound to contract. So called pinch grafts are not favoured by plastic surgeons principally because of the appalling stippled donor site that is produced in harvesting these grafts.

Full thickness grafts

Full thickness (or Wolfe grafts) were first described by a Scottish ophthalmic surgeon John Wolfe, who in 1875 successfully reconstructed a lower eyelid by removing a full thickness graft from the forearm. In this procedure the whole of the epidermis and dermis is transferred; sometimes it may include fat, hair, and sebaceous glands. The advantages of such a graft are the improved cosmetic appearance, the smaller likelihood of contracture, and the possible transfer in certain cases of hair. These grafts are often used for facial reconstruction and can be harvested from behind or in front of the ears, from the upper eyelids or from the supraclavicular region. Similarly, grafts for resurfacing the hand may be harvested from either the groin or the instep of the foot.

Composite graft from ear

The disadvantages are that the donor site does not regenerate and therefore has to be primarily closed or itself grafted with a partial thickness skin graft, thus limiting the size of the graft. The recipient bed must possess good vasculature and permit adequate immobilisation; these grafts consequently are often applied using a tie-over bolus dressing.

Skin grafts are initially more susceptible to actinic damage and should therefore be protected for six months. Thin partial thickness skin grafts do not have sebaceous glands and therefore tend to be dry and benefit from the application of lanolin cream. After the initial stage of graft contracture there is a secondary phase of growth in the graft, which parallels the rate of growth of the rest of the body. Scars, however, do not follow this secondary phase, and in children additional surgical procedures may therefore be required where skin grafts have been previously applied, particularly on the hand during adolescence.

Skin flaps

Skin and its subcutaneous tissue can be moved from one part of the body to another provided the vascular pedicle is maintained between it and the body for nourishment.

RANDOM SKIN FLAPS

Most local flaps are raised as random skin flaps and their design is dictated by experience. Their geometric design is only rough, and there are well defined limits to which the length of a flap can be raised despite increasing its width, which limits the safe transfer of skin. The skin in these flaps survives on blood vessels from the subdermal skin plexus, which provides a blood supply many times in excess of the metabolic requirements of the skin. This is because the skin has an important physiological role in regulating temperature. Once raised these flaps may be moved to an adjacent area of skin loss by rotation or transposition.

Rotation of random pattern flap

The design of a random flap obviously presents certain disadvantages in reconstruction. There are two ways of increasing the ratio of length to breadth. Firstly, a flap may be "delayed"; the flap is raised and transferred in more than one stage to insure its safety. At the first stage a longer flap than would normally survive is incised and partially raised. Ischaemia is thought to be a stimulus to increasing the blood supply to the dermis, possibly by an effect on the capillaries, the delay preventing the shunting of blood away from the dermis. At seven to 10 days a second delay or final raising and transfer of the flap is undertaken. Delaying the skin flap permits roughly a 60-100% increase in survival of flap length. At present despite recent reports flaps cannot be manipulated pharmacologically and the exact physiological basis of delay is not fully understood.

Secondly, much larger random flaps may be raised by joining two random flaps together, forming a bridge of skin that can be tubed with two pedicles, one at each end. The tube pedicle was introduced in 1916 simultaneously by Sir Harold Gilles in England and Vladimir Filatov, an ophthalmic surgeon, in Russia. At subsequent stages the flap can be divided at one end and inset on to the wrist. At a later stage the other end can be divided, allowing the flap to be transported on the wrist and inset on a distant site on the body. It is, however, a multistage procedure and therefore susceptible to many complications.

AXIAL PATTERN FLAPS

With the extended use of the previously described random flaps it became evident that on various areas of the body much longer flaps could be raised successfully without delay. McGregor and his colleagues studied such a flap in the groin and were able to raise a flap that was at least four times as long as its base. This was due to there being a constant anatomical blood vessel in the pedicle, in this case the superficial circumflex iliac artery and accompanying veins. It is now known that skin on the anterior chest wall may be raised as a long flap based on perforating branches from the internal mammary artery—the deltopectoral flap. Similarly, the forehead flap, which may be used to resurface areas of the face and oral cavity, is based on the anterior branch of the superficial temporal artery.

MUSCLE AND MYOCUTANEOUS FLAPS

Having established the anatomical basis of axial pattern flaps, plastic surgeons searched around the body for similar flaps. It became apparent that large areas of skin were not supplied on an axial basis but were supplied by perforating vessels coming through from underlying muscles. Muscles usually receive their blood supply via a pedicle at one end, and the other end of the muscle may therefore be detached and the muscle isolated on its pedicle with or without the overlying skin. This paddle of muscle can then be rotated through an arc, usually of 360 degrees. If muscle is moved alone skin grafts can be applied to its surface. On the other hand, muscles are often moved with the overlying skin as a myocutaneous flap. The commonly used myocutaneous flaps are pectoralis major used in head and neck reconstruction, latissimus dorsi in breast reconstruction, gluteus maximus in closure of sacral pressure sores, and gastrocnemius in reconstruction around the knee.

FASCIOCUTANEOUS FLAPS

More recently it has become evident that there are further areas of skin that are supplied by blood vessels issuing between muscles and supplying a rich vascular plexus that lies just above the deep fascia. This in turn supplies the subdermal plexus of the skin and is the basis of the fasciocutaneous flap that, in certain areas of the body, permits long flaps to be raised for local transposition. Such flaps are important for supplying skin cover to the lower leg and can be raised on the medial or lateral aspects of the shin and transposed over the tibia. Similarly, flaps may be raised from the upper arm or lateral aspect of the thorax for skin cover of the axilla.

FREE FLAPS

Despite the wealth of flaps that have become available to us there are certain areas of the body that remain extremely difficult to cover, in particular the scalp and the lower leg when there is extensive injury. With the advent of microsurgery, axial pattern flaps, surviving as they do on a single artery and accompanying vein, may be detached from their normal blood supply at the pedicle and the artery and vein anastomosed to a local undamaged artery and vein adjacent to the area of skin loss; provided the anastomoses remain patent these flaps will survive. Muscle and myocutaneous flaps may be similarly transferred as may the more recently described compound flaps such as the deep circumflex iliac artery flap that supplies the skin of the groin, adjacent muscle, and the bone of the iliac crest. This can be moved as a compound flap and used particularly in jaw reconstruction and reconstruction of severe injuries of the tibia.

Development of free flaps has permitted the transfer in one stage of tissue to a hostile environment where routine flaps would normally not survive because of the poor vasculature or because of infection. Such procedures, which are now becoming far more routine, although they undoubtedly take more operating time and require the surgical technique of microsurgery, permit a safer and more sophisticated reconstruction than was formerly possible and a greatly reduced stay in hospital compared with the original tube pedicles, which could take at least six months to move into the position in which they were required and had a 50% failure rate.

Mr Dai Davies, FRCS, is consultant plastic surgeon, West London Plastic Surgery Centre, West Middlesex University Hospital, Isleworth, and consultant plastic surgeon and honorary senior lecturer, Royal Postgraduate Medical School, Hammersmith Hospital, London W12 0HS.

Needs and Opportunities in Rehabilitation

Rehabilitation after stroke—2: Language and memory training and the requirements of rehabilitation services

DAPHNE GLOAG

"On the one side of the city was an area marked 'thoughts.' On the other side of the city was an area marked 'speech.' Down the middle was an unscaleable brick wall." So wrote Lord Smith of Marlow of one type of aphasia, a "Berlin Wall" that he himself did eventually scale (Chest, Heart and Stroke Association's magazine *Hope*, Spring 1983).

Communication problems and speech therapy

Patients who have been without speech often say later how upset they were at being talked about as though they could not understand.¹ Few staff, even doctors, are trained to communicate with aphasic patients and Mulley lists the simple do's and don'ts that make such a difference—including talking to the patient and avoiding tests and questions that may diminish him.¹ He urges the referral of all patients with language problems to a speech therapist as soon as possible for a precise diagnosis; different forms of aphasia, which may include poor comprehension, will need different forms of management, and these in turn will differ from the approaches needed for dysarthria and speech dyspraxia—and many patients have more than one type of disturbance.¹ Besides providing a diagnosis and if possible some form of speech stimulation, the speech therapist is important for general counselling of patients and family, for teaching the use of some communication aid if appropriate (usually in cases of dysarthria), and for advising staff. The broad elements of speech therapy are discussed by Code and

Müller² and alternative methods of communication by Rowley.³ Publications and other materials for those giving therapy are produced by, for example, the Winslow Press, Buckingham, and Action for Dysphasic Adults (ADA).

Develop communication by any means, said a consultant running a rehabilitation centre: any form of communication makes a great difference to the family as well as the afflicted person. Even the sexual relationship is apparently vulnerable in the face of speech loss.¹ *Coping with Disability*,³ *Help Yourself*,⁴ and pamphlets by the College of Speech Therapists, ADA, and the Chest, Heart, and Stroke Association are useful. Even if the patient never says a word again speech therapy may help to sort out the "muddle in the head," as this consultant put it. Language and communication generally, and not just speech, are the point. In this respect many studies of speech therapy may have too narrow a perspective.

Nevertheless, speech therapists are not plentiful and therapy as distinct from diagnosis and advice must be justified by hard evidence. Are there improvements that are not due simply to spontaneous recovery and are sufficient to justify all that goes into it? Do different forms of speech therapy have different degrees of success, and is therapy more than general stimulation and support—in other words, are speech therapists essential or can other people provide similar "treatment"? Finally, is timing crucial: should speech therapy coincide with the period of natural recovery (chiefly the first three months⁵) or take over where nature leaves off; and is it never too late for it to help, as individual cases would suggest? Theories of recovery, methods of treatment, and evaluation are discussed by Miller.⁶

Among more than 20 studies of speech therapy after stroke reviewed in 1979 most appeared to show benefit.⁷ Miller, however, reviewing the evidence (for aphasia of different types and causes), argues that there is no convincing evidence so far that speech therapy has a *specific* effect on aphasia.⁶ A recent trial suggested that