- Secretariat, GPO Box 5067, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico 00936.)
- 18 Mayhew DR. Alcohol, age and risk of accident involvement. In: Proceedings of ninth international conference on alcohol, drugs and traffic safety, 1983. (Available from Dr S Kaye, T-83 Secretariat, GPO Box 5067, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico 00936.)
 - 19 Allsop RE. *Alcohol and road accidents: a discussion of The Grand Rapids Study*. Crowthorne: Transport and Road Research Laboratory, 1966. (TRRL report No 6.)
 - 20 Fine EW, Steer RA. Short-term spatial memory deficits in men arrested for driving while intoxicated. *Am J Psychiatry* 1979;136:594-7.
 - 21 Tartar R. An analysis of cognitive deficits in chronic alcoholics. *J Nerv Ment Dis* 1973;147:138-47.
 - 22 Tartar R, Parsons O. Conceptual shifting in chronic alcoholics. *J Abnorm Psychol* 1971;77:71-5.
 - 23 Robertson I. Does moderate drinking cause mental impairment [Editorial]? *Br Med J* 1984;289:711-2.
 - 24 Emerson VJ, Hollyhead R, Isaacs MD, Fuller NA, Hunt DJ. The measurement of breath alcohol. *J Forensic Sci Soc* 1980;20:1.
 - 25 Nightingale J. Problem drinking among drunk drivers. *Br Med J* 1983;286:1648-9.
 - 26 Spoerer E. Rehabilitation of drinking drivers in the Federal Republic of Germany. In: Proceedings of ninth international conference on alcohol, drugs and traffic safety, 1983. (Available from Dr S Kaye, T-83 Secretariat, GPO Box 5067, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico 00936.)
 - 27 Siegal HA. The intervention approach; drunk driver rehabilitation. In: Proceedings of ninth international conference on alcohol, drugs and traffic safety, 1983. (Available from Dr S Kaye, T-83 Secretariat, GPO Box 5067, Medical Sciences Campus, University of Puerto Rico, San Juan, Puerto Rico 00936.)

(Accepted 13 November 1984)

Paediatrics Among Ethnic Minorities

Asian Families II: Conditions that may be found in the children

JOHN BLACK

Genetically determined conditions

BLOOD DISEASES

β thalassaemia is widely distributed throughout the Indian sub-continent, particularly in the north, though it is not nearly so common as in the Mediterranean area. More cases of homozygous β thalassaemia (thalassaemia major) are now seen in Pakistani families in London than in Greek, Cypriot, and Italian communities (for further discussion on β thalassaemia see article No 5: Families from the Mediterranean and Aegean regions). Moslem parents have more difficulty in accepting the ideas of genetic advice, antenatal testing, and termination of pregnancy than have parents from Mediterranean areas.

Haemoglobin D disease may occur in combination with β thalassaemia in children from northern India or Pakistan, and haemoglobin E thalassaemia occurs in Bengalis from Bangladesh. Both of these combined blood disorders result in chronic haemolytic anaemia with splenomegaly, requiring the same sort of treatment as homozygous β thalassaemia.

Sickle cell disease is not common in those groups that have emigrated to Britain, though this possibility should be considered in families originating from the Nilgiri Hills in southern India. There are also small areas with a notable incidence of sickle cell trait in Maharashtra, Bihar, Uttar Pradesh and West Bengal; (for further discussion on sickle cell disease see article 6: Afro-Caribbean (West Indian) and African families).

Glucose-6-phosphate dehydrogenase deficiency is fairly common throughout the northern part of the Indian subcontinent. The possibility of such a deficiency should always be considered in cases of neonatal jaundice when blood groups are compatible and in any cases of severe and sudden haemolytic anaemia that develop during an acute illness or as a result of taking certain drugs (for further discussion see later articles). When practicable G6PD deficiency should be excluded before giving primaquine for malaria.

RECESSIVELY DETERMINED METABOLIC DISORDERS

There is little doubt that the high incidence of recessively determined metabolic disorders and some of the more unusual malformations in Moslem families from Pakistan is due to the number of marriages among first and second cousins.

Acquired diseases

IRON DEFICIENCY ANAEMIA

Iron deficiency anaemia is the commonest form of anaemia in Asian children and is of dietary origin. Maternal iron deficiency in pregnancy and prematurity and low birth weight are both important factors, as is prolonged milk feeding and the late introduction of solids. Infants who are exclusively breast fed for a year or 18 months develop severe iron deficiency anaemia. The use of artificial milks, fortified with small amounts of iron, reduces the severity of the anaemia but does not necessarily prevent it.

A vegetarian diet in Britain, as eaten by strict Hindu families, results in iron deficiency, but many Hindu parents in Britain give their children meat and eggs. The peak incidence of iron deficiency is at about 3 years, though for the reasons already discussed the condition may develop much earlier. On a vegetarian diet the best sources of iron are eggs (if these are part of the normal diet of the family), dried fruit, green vegetables, whole wheat chapatis, peas, beans, and other pulses (seeds of leguminous vegetables, which when split are known as "dahl"). If one member of a family is found to have iron deficiency anaemia the other members of the family should have haemoglobin concentrations measured and receive dietary advice from someone familiar with the diets of that particular community. Though not vegetarians, children from Bangladesh are also likely to have iron deficiency anaemia.² Many Asian families have now adopted a partly European diet.

Poor economic circumstances in any group may result in a diet deficient in iron.

A child with iron deficiency anaemia who has been living in Asia from the age of 1 or who has become anaemic after a visit to the home country should be suspected of having hookworm disease, particularly if his or her response to treatment with iron is poor (for further discussion on hookworm disease see below).

Epidemiological assessment of status of Malaria, June 1982. Reproduced from Communicable Disease Report 82/52 with permission of the editor, Communicable Disease Reports, PHLS Communicable Disease Surveillance Centre, London.

NUTRITIONAL RICKETS

The main factors causing rickets in Asian children are:

(a) lack of adequate exposure to sunlight: this affects preschool children and adolescent girls, particularly Moslems, who may be unwilling to uncover their arms and legs in public;

(b) vegetarian diets: the group most at risk are strict vegetarian Hindu families (often Gujaratis);

(c) the use of household cows' milk for infant feeding: this contains negligible amounts of vitamin D;

(d) maternal deficiency of vitamin D during pregnancy and lactation.

Clinically, nutritional rickets is seen in newborn infants, toddlers, preschool children, and adolescents.

The newborn infant—Fetal rickets may result in rickets in newborn infants, which may be recognised by craniotabes or hypocalcaemic fits, though neither is specific to rickets alone.

Infants, toddlers, and preschool children—From the age of 6 months severe rickets may develop in a child who is solely breast fed by a mother deficient in vitamin D or who is given cows' milk and unfortified weaning foods. The site of the deformities depends on the age at onset—for example, bowing of the forearm may develop in an infant who is able to crawl. Active rickets is not always recognised as a painful disease and can cause delay in, or discourage walking in a child who was previously walking quite well.

School age children—Active rickets in school children is rare as they usually have an adequate exposure to sunlight, but previously unrecognised deformities such as bow legs or genu valgum may be detected on entry to school.

Adolescent rickets—Girls are more likely to be affected than boys. Deformities are unlikely, but limb pains and backache are common presenting symptoms.

PREVENTION OF NUTRITIONAL RICKETS

Education—In 1981 the "Stop Rickets" campaign was started jointly by the Department of Health and Social Security and Save

the Children Fund to inform and educate Asian communities and health workers about the causes, prevention, and detection of nutritional rickets. In September 1984 the "Asian Mother and Baby" campaign was launched by the same organisations to improve attendance at antenatal clinics and to encourage breast feeding.

Nutrition—Each Asian group should be considered separately because their dietary customs, religious beliefs, and socioeconomic circumstances are different. Leaflets in the main Asian languages on antenatal care, nutrition, rickets, and breast feeding are available from clinics or the Health Education Council. Information about instruction material for health workers for both campaigns can be obtained from the Save the Children Fund. In practice the vitamin D content of the typical diet of most Asian families makes a very small contribution to their vitamin D intake. Mothers who are bottle feeding their babies should be advised to use one of the standard baby milks, which are all fortified with vitamin D. Baby milks should be continued, instead of cows' milk, up to the age of 1. Other foods that normally contain added vitamin D are baby cereals and breakfast foods. Most weaning foods containing fruit and vegetables used by Asian mothers (who are generally unwilling to use those containing meat) are not fortified, though those manufactured by Milupa contain added vitamin D.

Vitamin supplements—The standard "children's vitamin drops" supplied through child health clinics and welfare food distribution centres are cheap and available free to families claiming supplementary benefit or family income supplement for children up to the age of 5 (for further information see DHSS leaflet MV11 November, 1983). Five drops given once daily contain 7 µg vitamin D (280 iu), 200 µg vitamin A, and 20 mg vitamin C (the daily requirement of vitamin D is 7.5 µg up to the age of 1 and 10 µg from 1 onwards). These vitamin drops should be given from the age of 1 month and continued up to 5 years. Older children who are considered to be especially susceptible to rickets should also be given the vitamin drops. Children should not be receiving other preparations containing vitamin D in addition to the drops as there is a risk of overdose in such circumstances.

INTESTINAL INFECTIONS

Bacterial infections

Bacterial infections may be acquired while visiting Asia, from an infected person within the family, or from infected food. Newborn infants can be infected during delivery if the mother is a carrier of pathogenic bacteria.

Shigellosis—In infants shigellosis may be a septicaemic illness and in those aged under 5 the initial fever may be accompanied by a convulsion. The symptoms are a high temperature and sudden onset of severe diarrhoea with blood and mucus in the stool.

Salmonellosis (numerous strains of *Salmonella typhimurium*; also *S typhi* and *S paratyphi* A, B, and C)—*S typhimurium* causes profuse diarrhoea sometimes with streaks of blood and mucus. A septicaemic illness with splenomegaly occurs in older children and mimics tuberculous meningitis. Typhoid and paratyroid fever affect mainly schoolchildren; severe frontal headache is a common initial symptom followed by vomiting, abdominal pain, and diarrhoea. The rash is difficult to detect in dark skinned children. Infection of children in Britain from cases or carriers is not uncommon.

Protozoal infections

Giardiasis should be considered when there is chronic diarrhoea and colicky pain with pale, frothy stools and increased flatus, producing in its chronic form a condition like coeliac disease.

Amoebiasis causes diarrhoea and tenesmus with blood and mucus but is less acute than bacterial dysentery; occasionally severe intestinal haemorrhage with shock occurs. Amoebic and bacillary dysentery may occur together.

Worms

Roundworm infection (ascariasis) is extremely common in children born in the Indian subcontinent and may be acquired during a visit there. Although infants may be infected, children aged 1-5 are the most common sufferers. The diagnosis may be determined by the passage of one or two worms from the rectum or from the nose, which may be frightening for the parents. Other symptoms may be colicky abdominal pain or actual intestinal obstruction. In heavily infected children pre-existing malnutrition may be exacerbated. The life of the adult worm is about one year. Infection in Britain is unlikely. The diagnosis can easily be made by identifying the ova in the stool. Eosinophilia is unlikely to be observed.

Hookworm infection (ankylostomiasis) is as common as ascariasis infection, and the two often coexist. Infection usually occurs in toddlers, but infants are occasionally infected. No intestinal symptoms occur, but in heavily infected children hypochromic microcytic (iron deficiency) anaemia slowly develops. The diagnosis is made by identifying ova in the stools; a test for faecal occult blood is usually positive. Infection does not occur in Britain but if untreated infection with ankylostoma may persist for up to four years.

MALARIA

Malaria in children born in the United Kingdom is acquired during visits to the home country. A few older children entering the country for the first time develop a relapse of malaria acquired previously. Many parents do not realise that the area from which they came, previously free from malaria, may once again have become malarious and therefore do not take any precautions on their return to Asia. Infants are rarely given antimalarial drugs in the mistaken but widespread belief that these drugs are too toxic for them. Malaria is never acquired in Britain (with very rare exceptions of infected mosquitoes arriving by air) and cannot be transmitted to other people in the United Kingdom.

Malaria should be considered whenever fever develops suddenly in a child who has recently returned from Asia. In those aged under 5 a febrile convulsion may occur. In a first attack the spikes of fever

are often irregular and the spleen may not be palpable during the first two days after the onset of fever.

Infections acquired in the Indian subcontinent are invariably due either to *Plasmodium vivax* infection (benign tertian malaria), which produces a fairly mild disease or to *Plasmodium falciparum* infection (malignant tertian malaria), which can produce a severe and occasionally fatal illness. Falciparum malaria should be suspected when any of the following features are present.

(1) Convulsions or coma with a variable degree of fever, a meningitis like picture, hemiplegia, or confusional state may indicate cerebral malaria. This condition is an acute emergency and should be treated with parenteral chloroquine (or quinine if there is a possibility of chloroquine resistance, which is most likely in infection acquired in Bangladesh or eastern India), but not before thick and thin blood films have been examined. All hospitals should carry supplies of parenteral chloroquine and quinine. (2) Hyperpyrexia. (3) Fever, diarrhoea, vomiting, and shock. (4) Pallor, jaundice, and splenomegaly with fever; this condition is sometimes mistaken for hepatitis A.

Recurrences of benign tertian malaria may occur up to two years after exposure and plasmodium malaria up to six weeks or, exceptionally, up to one year after infection.

KALA AZAR (VISCERAL LEISHMANIASIS)

Kala azar is not uncommon in Bangladesh and should be considered in a chronically ill child with intermittent fever, lassitude, and a firm enlargement of the liver and spleen. The infection may have been acquired before the child came to Britain or on a visit to Asia. The incubation period is usually two to six months but may be as short as three to four weeks or as long as two years.

TUBERCULOSIS³

When the possibility of tuberculosis is considered the first thing to look for is a BCG scar, though this is not an absolute guarantee of protection. Primary tuberculosis should be considered in children who fail to gain weight or lose weight for no obvious reason. A strongly positive result from a Mantoux or Heaf test invariably indicates an active infection. Respiratory tuberculosis in Asian children sometimes takes atypical forms, and the adult type of infection with cavitation may occur. Tuberculosis should also be suspected in any chronic lung condition that fails to clear up within two to three weeks.

Less common forms of infection are abdominal tuberculosis, which may occur with intestinal haemorrhage, ascites or obstruction; or bone or joint infection. Miliary and meningeal infections are fairly common.

Traditional practices and medicine⁴

Traditionally in Asia a black eye make up (Surma, Kajil) is put round the eyes and inside the lids in babies and small children, partly for its cosmetic effect and partly because it is thought to protect the eyes against infection. Originally made from soot from oil lamps and vegetable dyes or antimony sulphide, an adulterated form of surma is now made from lead sulphide⁵; cumulatively, lead can be absorbed in sufficient amounts to cause lead poisoning.^{6,7} The importation into Britain of this lead containing preparation has been banned, but some of it may still be brought into the country through non-commercial channels. Suitable hypoallergenic make up preparations can be substituted quite safely. Kajil may also contain lead and is used in the same way as surma but is in the form of an ointment. Several medicines for infants also contain lead, which may be prescribed by practitioners of traditional Indian medicine (hakims for unani medicine and vaidas for the ayurvedic system).

A similar substance known as al kohl is used in Arab countries and may also contain lead.

Some common problems

Illiteracy is not uncommon,⁸ particularly in older women who have spent their childhood in the subcontinent. Leaflets and instructions in the appropriate language may therefore not meet the need, though it is unusual to find a family without at least one adolescent or school age member who is completely fluent in English. If there is a possibility of non-compliance because of language difficulties, however, and failure to take the prescribed drugs may have serious consequences, as in tuberculosis, it is essential to obtain the help of a competent interpreter. The use of children as interpreters should be discouraged, except for the most straightforward situations.

Apparent refusal to breast feed—Many Asian women believe that colostrum is harmful to the baby and may therefore be reluctant to breast feed their baby during the first two or three days. If this is the mother's belief it should be respected and it should not be assumed that she does not wish to breast feed at all.

Carriers of hepatitis B virus in pregnancy—From 3% to 10% of adults in the Indian subcontinent are potentially infective carriers of hepatitis B virus and can infect an infant at or after delivery. An infant infected at birth can become an asymptomatic carrier, who is infective for many years and may occasionally develop acute or chronic liver disease or hepatocellular carcinoma in adult life. It is therefore important to screen all pregnant Asian women at their first antenatal visit. As infection within the family is common women who have been born in Britain and have never left this country should be included in the screening programme. If the mother is found to be infective special precautions should be taken at delivery to prevent infection of the infant and medical and nursing staff. The infant should be given specific hepatitis B immunoglobulin and hepatitis B vaccine, according to the schedule described for Chinese and Vietnamese families. As the incidence of the carrier state is two to three times higher in women from south east Asia than in those from the Indian subcontinent the management of this problem will be discussed in greater detail in article 7: "Chinese and Vietnamese Families."

Neonatal jaundice—As G6PD deficiency is not uncommon in Asian families this possibility should always be considered when non-obstructive jaundice develops on the second or third day of life. Girls may be affected (for further discussion see article No 5: Families from the Mediterranean and Aegean), but boys are affected 40 times more often in most communities. In preterm infants or when other factors are contributing to the jaundice an exchange transfusion may be required.

Small stature, particularly in boys, is a common cause of parental anxiety among the Bengalis from Bangladesh. Generally, both parents are small but would like their children to be tall to cope with possible physical aggression. The presenting complaint is usually that "he is not growing." This may be true but is more likely to mean that the child is not as tall as the parents would like and is in fact growing at a normal rate but at or below the third centile for British children. Having excluded chronic disease and the more obvious endocrine disorders, the parents may accept that nothing can be done except to confirm a normal rate of growth by measuring the child's height again after six months; or a growth hormone stimulation test may be needed to add more convincing evidence.

Anaemia without splenomegaly—Iron deficiency is the commonest cause of moderate or severe anaemia and should be considered first; the causes of iron deficiency have already been discussed. In strictly vegetarian families folic acid or vitamin B₁₂ deficiency may occur and should be suspected initially from the appearance of the blood film. In these nutritional anaemias splenomegaly does not occur; if it is present an alternative diagnosis should be considered.

Anaemia with hepatosplenomegaly—In children who have never been to the Indian subcontinent or outside Europe the most likely diagnosis is one of the haemoglobinopathies. Homozygous β thalassaemia (thalassaemia major) is the most common condition, but haemoglobin D thalassaemia or haemoglobin E thalassaemia is also possible; sickle cell anaemia is very rare. (For discussion on geographical distribution of the haemoglobinopathies see article No 6: Afro-Caribbean (West Indian) and African families.) In children who have lived in or visited the subcontinent additional diagnoses

such as chronic malaria and kala azar (in India, West Bengal, Assam, and Bangladesh predominantly) may have to be considered.

Hepatitis—Chronic active hepatitis B infection or early cirrhosis is not uncommon in children presenting with chronic ill health and an enlarged liver. Splenomegaly is found only in advanced cirrhosis with portal hypertension in older children. Hepatitis A may develop in children who have recently returned from a visit to the Indian subcontinent; chronic or persistent liver disease is a rare sequel to this form of hepatitis.

I thank Dr Malcolm Arthurton, formerly consultant paediatrician, Bradford Children's Hospital, Dr Edwin Lobo, consultant paediatrician, Luton and Dunstable Hospital, Dr J S Gill, Department of Medicine, University of Birmingham, and Vanessa N Shaw, paediatric dietitian, Queen Elizabeth Hospital for Children, London.

References

- 1 Karan VK, Prasad SN, Prasad TB. Sickle cell disorder in aboriginal tribes in Chotangapur. *Indian Paediatr* 1978;15:287-91.
- 2 Harris RJ, Armstrong D, Ali R, Loynes A. Nutritional survey of Bangladeshi children aged under 5 years in the London borough of Tower Hamlets. *Arch Dis Child* 1983;58:428-32.
- 3 Darbyshire JH. We don't have tuberculosis in this country any more . . . do we? *Maternal and Child Health* 1978;8:176-82.
- 4 Aslam M, Healy MA. Present and future trends in the health care of British-Asian children. *Nursing Times* 1982;1353-4.
- 5 Green SDR, Lealman GT, Aslam M, Davis SS. Surma and blood lead concentrations. *Public Health* 1979;93:371-6.
- 6 Warley MA, Blackledge P, O'Gorman P. Lead poisoning from eye cosmetics. *Br Med J* 1968;i:117.
- 7 Fernando NP, Healy MA, Aslam M, Davis SS, Hussein A. Lead poisoning and traditional practices: consequences for world health. A study in Kuwait. *Public Health* 1981;95:250-60.
- 8 Aslam M, Davis SS, Fletcher R. Compliance in medication by Asian immigrants. *Nursing Times* 1979;931-2.

Further reading:

Arthurton M. Immigrant children and the day-to-day work of a paediatrician. *Arch Dis Child* 1972; 47:126-30.

(This article is the fourth in a series)

Are there any interactions between acetazolamide given for glaucoma and other diuretics given for heart failure? Do thiazide or loop diuretics affect pressure in the eye?

It is not considered necessary to give potassium supplements to patients taking long term acetazolamide when it is given in the absence of other thiazide diuretics. Nevertheless, additional potassium should be given to patients receiving acetazolamide as well as thiazide diuretics. There is no evidence that clinically useful lowering of intraocular pressure follows the use of thiazide diuretics. It should be remembered, however, that digitalis, which reduces the production of aqueous by the ciliary epithelium, will lower intraocular pressure.—R A HITCHINGS, consultant ophthalmic surgeon, London.

Chandler PA, Grant WM. *Glaucoma*. Philadelphia: Lee and Febiger, 1979:85.

How may the risk of osteomalacia be reduced in a woman of 67?

Given that the patient is not suffering from malabsorption or renal disease, osteomalacia should be preventable. An adequate intake of vitamin D should reduce the risk to nil. The relative importance of vitamin D intake due to exposure to sunlight and from the diet has not been finally settled. Quite minor exposure to sunlight may be sufficient to supplement a poor vitamin D intake in preventing clinical osteomalacia.¹ A combination of a good summer holiday with plenty of sunshine which produces a well tanned skin on the face and forearms and a regular intake of margarine, eggs, and oily fish such as tuna, mackerel, and herring should be sufficient to maintain an adequate intake throughout the year.—L KLENERMAN, consultant orthopaedic surgeon, London.

1 Hodkinson HM, Round P, Stanton BR, Morgan C. Sunlight, vitamin D, and osteomalacia in the elderly. *Lancet* 1973;ii:910-2.