PRACTICE OBSERVED

Practice Research

Study of medicine prescribing for elderly patients

CHARLES B FREER

Abstract

An analysis of drug prescribing over six months in a random sample of 146 elderly patients showed that 42% were receiving one or more medicines long term. Diuretics, analgesics, and non-steroidal anti-inflammatory drugs were most often prescribed, and variations in prescribing by age, sex, and consultations with the general practitioner were examined. Only 17% of the group were taking three or more medicines on repeat prescriptions, and 42% received no prescriptions during the study. These results have implications for the workload of supervising prescriptions but also underline the need for a balanced view of drug prescribing and medicine taking in elderly people.

Introduction

For elderly patients, more than for patients in other age groups, growth in the availability of drug treatments has been a double edged sword, and a recent report from the Royal College of Physicians has reminded doctors of the problems associated with inappropriate and excessive prescribing for older patients. Dunnell and Cartwright found that 92% of patients aged over 75 had taken at least one medicine in the two weeks before their survey, and up to half of all older patients were taking at least one medicine long term.¹ Similar results were found in two studies in general practice.^{2,3} The results of an unpublished study of long term drug prescribing for the elderly in a university department practice showed a similar order of repeat prescribing, but it also appeared that many older patients were receiving few or no prescription medicines.

University of Southampton
CHARLES B FREER, MA, MRCP, senior lecturer in primary medical care
Correspondence to Aldermore Health Centre, Southampton SO1 6ST

This paper reports the results of a study designed to provide detailed information on the distribution of all medicine prescribing for a random sample of older patients during a six month period.

Background

Aldermore Health Centre has a National Health Service practice of just over 8000 patients (10% are aged 65 and over) and is staffed by members of the primary medical care department of the university. Current research activities include the effect of a restricted drug list on the prescribing practices of the doctors. This is not a rapid list but provides recommendations on prescribing for common conditions based on pharmacological and economic factors.

Method and results

A 20% random sample of all patients aged 65 and over was obtained from the practice age-sex register, and details of the medicines prescribed and the number of repeat and single prescriptions were collected manually from their records for the six months April to September 1984 inclusive. The total number of contacts with the general practitioner in the surgery and home for each patient for the 11 months January to November 1984 was also noted. A repeat prescription was defined as the same medicine issued on two or more occasions as continuous treatment for a chronic problem; for example, two prescriptions for 100 mg of steroid cream in a patient with eczema were defined as repeat prescriptions, but two prescriptions for prednisolone during the six months for exacerbations of bronchial asthma were not.

Study group.—The 20% random sample produced 161 patients, of whom 12 had left the practice and three had died, leaving 146 patients. The female to male ratio was 1.7:1 (ages 65-74: 32 men, 38 women; age 75 and over: 23 men, 53 women) compared with 1.5:1 for all elderly patients in the practice. Patients in the older age band were overrepresented, with a ratio of 65.74 years to 75 years and over of 0.9 compared with 1.5 for the practice.

Number of prescribed medicines.—Sixty-two patients had received at least one repeat prescription, 61 no repeat prescriptions at all, and only 23 at least one single prescription but no repeat medicines over the six months (table 1). The 62 patients had received repeat prescriptions for 148 drugs, an average of 2.4 per patient, with only three taking five or more prescription medicines regularly (figure 1). Some patients on repeat prescription had also

reasonable to conclude from this information that elderly people have a disproportionately high intake of medicine compared with patients in other age groups.

The results have an important implication for the workload of supervising repeat prescribing in elderly patients—which is a reasonable expectation of adequate clinical care—because not all patients need to be supervised. For example, in this study only 25 patients were taking three or more medicines on repeat prescription, representing only 17% of all patients over the age of 65.

Elderly people are at particular risk from drug treatment, and acceptable standards of clinical care should include appropriate prescribing practices and the supervision of long term treatment. Justifiable concern, however, should be based on a balanced view of the problem of excessive and inappropriate prescribing for such patients, and examples that are extreme or apply to only a few patients must not be extrapolated to hide the large numbers of older patients who are neither frequent attenders nor on long term drug treatment. This is not only likely to overstate the problem of poor prescribing practices and medicine taking in elderly people but unwittingly reinforce negative stereotypes of elderly people.

I thank Diane Gluckin for help in collecting information from patients' records and Jean Gibson for typing the manuscript.

References

- Report of the Royal College of Physicians. *Medication for the elderly*. J R Coll Physicians Lond 1984;18:7-17.
- Stuart DG, Oyle J. *Need for supervision in the elderly receiving long-term prescribed medication*. Br Med J 1979;199:505-7.
- Talbot AJ. Repeat prescribing for elderly patients. Br Med J 1981;282:1672-5.
- Stewart J. The epidemiology of prescribing in an urban general practice. J R Coll Gen Pract 1980;30:191-6.
- Anderson KM. The use of repeatedly prescribed medicines. J R Coll Gen Pract 1984;34:409-11.
- Goorney M, Talbot AJ. Prescriptions of cardiovascular and antineoplastic drugs in elderly patients admitted to hospital. Lancet 1984;1:944-5.
- Stewart J. Prescribing and the physician in prescribing in general practice. J R Coll Gen Pract 1979;29:100-2.
- Martyn CR. Drug treatment in elderly patients. GP update. Br Med J 1982;285:1623-5.
- Bain M, Hunt J, Joyce D, Macdonald M, Woodhouse J. *Treatment of disease: a study of prescriptions in general practice*. London: Tavistock Publications, 1976.
- Frost CR. Old people. Lancet 1983;1:268-9.

Accepted 18 January 1985

Computerised repeat prescriptions: simple system

MALCOLM AYLETT

Abstract

In a small practice it is possible to develop a computer system for repeat prescriptions that requires little extra work and has other applications that may improve the quality of care of chronic conditions and promote the education of the general practitioner and his trainees. A program that can maintain a patient register, check illness and dispensing figures, prepare call up lists for inoculations, and search specific age groups for particular conditions, as well as provide a repeat prescription function, is described.

Introduction

Some early general practice repeat prescription systems offered little more than the facility to store details of continuing medication and print prescriptions either when required or in batches. Systems with a wider range of functions are sometimes installed with the intention of carrying out many other tasks but end up being used almost exclusively to print repeat prescriptions.

Modern systems are interactive. That is, details of any prescriptions printed are fed back into the patient's file and with a set of controls, warnings, and observations to staff and patients on the screen and the blank space of the FP10 (COMP), repeat prescribing is efficiently monitored. But these systems depend on consensus and tight discipline among doctors and their staff for any handwritten prescriptions will lead to a false indication of non-compliance. The flexibility that is available on a card system to give, for example, larger or smaller amounts, early or late prescriptions, or allow for many other common exigencies becomes impaired. A Sheffield practice reported last year that it found these snags almost insuperable. After several years' experience with a sophisticated

operation supported by the Medical Research Council, the practice doubted whether a computerised repeat prescription function was necessarily more efficient or more reliable than a manual system.

It is hard to improve on the accessibility of the information held, the ease of entry, and the "user friendliness" of the manual card system run by most well organised practices. Why not retain a card system that runs well and add a simple computerised back up that will print all the prescriptions (and labels if you dispense) and also keep data that enable additional monitoring of the prescribing plus an existing range of other applications? In a small card system that fulfils all the functions expected of a good record, plus facilities for enhancing care and considerably extending audit, education, and research.

The practice

A single-handed practice with a list of 1600 patients, dispensing for 900, in a village some 17 miles from the nearest community hospital and 40 miles from the district general hospital could be described as atypical. But with a normal incidence of chronic illness in a top heavy age structure, repeat prescribing is commensurate with that of a national average list. Exceptions are that fewer patients are on psychotropics (11 on daytime treatment and 28 taking hypnotics), and a larger number of patients are taking hypotensive drugs (94, which is 8% of those aged 40 to 65 years) as a result of screening.

Repeat prescribing is monitored by a card kept in the patients' records with letter codes for the items sanctioned, serial entries of dates and codes recording prescribing, and regularly updated months up to which a prescription is allowed without consultation. The necessity to revise frequently the "to be seen" month because of unscheduled consultations is a major problem with any system since these patients have a high incidence of intercurrent consultation and doctors are liable to omit making this change each time the patient is seen. Signing the prescription thus therefore be accompanied by checks on the repeat card, of the prescription itself, and not infrequently of the patient's record. Attention to these details,

Gloucester Surgery, Wooler, Northumberland NE1 6DN
MALCOLM AYLETT, MA, MRCP, general practitioner

TABLE 1—Number of patients who received repeat prescriptions, single prescriptions only, and no prescriptions from April to September 1984, by age and sex

	Repeat prescriptions	Single prescriptions	No prescriptions	Total
All men	16	5	11	32
All women	9	11	11	31
65-74 years	12	7	10	29
75 years and over	25	8	19	52
Total	62 (42%)	23 (16%)	61 (42%)	146 (100%)

Number of repeat prescriptions per patient by age (n=62)

TABLE 11—Drugs prescribed by repeat prescription

Type of drug	No of prescriptions
Diuretics	16
Analgesics	15
Non-steroidal anti-inflammatory drugs	14
Bronchodilators	11
Cardiac drugs (digitalis, A, digoxin, and furosemide)	10
Antidepressants	9
Antihypertensives (salts)	8
Oral contraceptives	7
Anticancer drugs	6
Laxatives	5
Anticongestants	4
Oral hypoglycaemics	4
Antibiotics (penicillin, tetracycline, and erythromycin)	4
Topical medicines (steroids, 1, aqueous cream, and antineoplastic)	4
Tranquilisers	3
Eye drops	3
Anticancer supplements	3
Anticancer drugs	3
Quinine	2
Anticancer drugs	2
Sodium nitroprusside	2
Miscellaneous (all new type pills)	2
Total	147

received single prescriptions, but the small total numbers (64) made it difficult to interpret the distribution. There was, however, no apparent tendency for those on higher numbers of repeat prescriptions to be receiving a disproportionately greater number of single prescriptions.

Types of prescribed medicines.—Almost half the medicines prescribed as a repeat were for diuretics, inhalers (bronchodilator and steroid) for asthma, analgesics, non-steroidal anti-inflammatory drugs, and cardiac drugs (table 11). Only 17 (11%) prescriptions were for psychotropic drugs. The 64 single prescriptions were for diuretics (13), antibiotics (12), non-steroidal anti-

inflammatory drugs (eight), analgesics (seven), topical steroids (seven), oral steroids (three), laxatives (two), antineoplastic (two), and miscellaneous (one).

Drug prescribing and consulting patterns.—Almost half of the patients on repeat prescriptions were seen five or more times in the first six months of 1984, and there was an inverse relation between receiving no prescribed medicines and the number of consultations with the general practitioner (table 11). The distribution of single prescriptions did not seem to be related to consultation rates.

TABLE 11—Distribution of prescribing by number of consultations with general practitioner (surgery plus home) during January to November 1984 inclusive (n=146)

No of consultations with general practitioner	Repeat prescriptions	Single prescriptions	No prescriptions
0	4	6	27
1	8	10	11
2	12	14	10
3	15	16	10
4	18	18	10
5	20	20	10
6	22	22	10
7	24	24	10
8	26	26	10
9	28	28	10
10	30	30	10
11	32	32	10
12	34	34	10
13	36	36	10
14	38	38	10
15	40	40	10
16	42	42	10
17	44	44	10
18	46	46	10
19	48	48	10
20	50	50	10
21	52	52	10
22	54	54	10
23	56	56	10
24	58	58	10
25	60	60	10
26	62	62	10
27	64	64	10
28	66	66	10
29	68	68	10
30	70	70	10
31	72	72	10
32	74	74	10
33	76	76	10
34	78	78	10
35	80	80	10
36	82	82	10
37	84	84	10
38	86	86	10
39	88	88	10
40	90	90	10
41	92	92	10
42	94	94	10
43	96	96	10
44	98	98	10
45	100	100	10
46	102	102	10
47	104	104	10
48	106	106	10
49	108	108	10
50	110	110	10
51	112	112	10
52	114	114	10
53	116	116	10
54	118	118	10
55	120	120	10
56	122	122	10
57	124	124	10
58	126	126	10
59	128	128	10
60	130	130	10
61	132	132	10
62	134	134	10
63	136	136	10
64	138	138	10
65	140	140	10
66	142	142	10
67	144	144	10
68	146	146	10
69	148	148	10
70	150	150	10
71	152	152	10
72	154	154	10
73	156	156	10
74	158	158	10
75	160	160	10
76	162	162	10
77	164	164	10
78	166	166	10
79	168	168	10
80	170	170	10
81	172	172	10
82	174	174	10
83	176	176	10
84	178	178	10
85	180	180	10
86	182	182	10
87	184	184	10
88	186	186	10
89	188	188	10
90	190	190	10
91	192	192	10
92	194	194	10
93	196	196	10
94	198	198	10
95	200	200	10
96	202	202	10
97	204	204	10
98	206	206	10
99	208	208	10
100	210	210	10
101	212	212	10
102	214	214	10
103	216	216	10
104	218	218	10
105	220	220	10
106	222	222	10
107	224	224	10
108	226	226	10
109	228	228	10
110	230	230	10
111	232	232	10
112	234	234	10
113	236	236	10
114	238	238	10
115	240	240	10
116	242	242	10
117	244	244	10
118	246	246	10
119	248	248	10
120	250	250	10
121	252	252	10
122	254	254	10
123	256	256	10
124	258	258	10
125	260	260	10
126	262	262	10
127	264	264	10
128	266	266	10
129	268	268	10
130	270	270	10
131	272	272	10
132	274	274	10
133	276	276	10
134	278	278	10
135	280	280	10
136	282	282	10
137	284	284	10
138	286	286	10
139	288	288	10
140	290	290	10
141	292	292	10
142	294	294	10
143	296	296	10
144	298	298	10
145	300	300	10
146	302	302	10
147	304	304	10
148	306	306	10
149	308	308	10
150	310	310	10
151	312	312	10
152	314	314	10
153	316	316	10
154	318	318	10
155	320	320	10
156	322	322	10
157	324	324	10
158	326	326	10
159	328	328	10
160	330	330	10
161	332	332	10
162	334	334	10
163	336	336	10
164	338	338	10
165	340	340	10
166	342	342	10
167	344	344	10
168	346	346	10
169	348	348	10
170	350	350	10
171	352	352	10
172	354	354	10
173	356	356	10
174	358	358	10
175	360	360	10
176	362	362	10
177	364	364	10
178	366	366	10
179	368	368	10
180	370	370	10
181	372	372	10
182	374	374	10
183	376	376	10
184	378	378	10
185	380	380	10
186	382	382	