PRACTICE OBSERVED

Practice Research

Follow up study of angina in patients aged 30 to 59 in general practice

D W HOBKIRK

Abstract

In 1979, 51 general practitioners in the Northern region identified 336 patients aged 30 to 59 who had angina pectoris. Information was obtained on their treatment and how angina affected their daily lives. This study reviews the same patients after an interval of four years. Of the 307 who were not lost to follow up, 18 (12%) had died and 44 (14%) were free from angina in the last year.

Drugs seemed to be used to their best advantage, with a fourfold increase in the use of the newer calcium antagonists and long acting trinitrate preparations. Many patients still suffered from angina, which was severely affecting their daily lives, but only 41 patients had had exercise electrocardiograms recorded or more detailed investigations; this includes 16 (6%) patients who had had coronary artery bypass surgery.

Introduction

In 1982 we reported on 336 young patients who had angina pectoris. This follow up study after an interval of four years was carried out to look at the subsequent morbidity and mortality of this general practice population, with particular reference, firstly, to those who had originally been found to suffer from "severe" angina and, secondly, to changes in drug treatment and the use of coronary artery bypass surgery. Most studies from hospital present a distorted picture as the patients referred to them are presumably those with more severe angina.

Health Centre, Gushborough, Cleveland TS14 7AA
(D W HOBKIRK, senior general practitioner)

Method

During 1979, 51 general practitioners from 20 practices in the Northern region reported on 336 patients aged 30 to 59 who had angina pectoris. All patients also completed a questionnaire about their angina and the effect that it had on their daily lives. In 1983 each general practitioner or successor completed a second questionnaire about the same patients, also distributing a follow up questionnaire to patients in the practice who were still alive. For those patients who had moved the relevant family practitioner committee was usually able to help to track their new general practitioner, who was then contacted.

The doctor's follow up questionnaire inquired whether it was still thought that the patient had angina in 1979 and whether the patient still had it; what had been made of drugs to treat angina; whether any complications such as myocardial infarction had occurred; and to what extent the hospital services had been concerned in the patient's care. The date and cause of death for patients who had died was recorded.

The patients' follow up questionnaire again concentrated on their daily lives and the effect that angina or its absence had on it. For the most part the questions were identical to those in the original study, enabling direct comparisons to be made. Statistical comparisons were made by the χ^2 test.

Results

Of the original 336 patients, 307 (91%) entered the second study. Of the remaining 29 patients, four declined to participate in the follow up study but most had left the region. There was no evidence of an excessive number of deaths in this group.

The analysis relates to the remaining 307 patients. The 38 patients who died are excluded from some analyses indicated to enable more direct comparisons because their medical histories between the first study and their death was not available.

SEVERITY OF ANGINA

As in the first study patients were divided into two groups of "severe" and "not severe" angina. The definition of "severe" angina was the same as before—namely, the patient who had attacks not only when walking at an

ordinary pace on the level but also had one or more attacks a day, and whose angina affected one or more of the following activities: sport, sex, housework, and shopping. The changes in severity over the four years are shown in the figure. In the 30 patients whose angina was no longer "severe" four (13%) had had coronary artery bypass surgery. In all other respects, including drug treatment and daily activities, there was no difference between this group and the remaining 186 with "not severe" angina.

Retrospectively, the doctors thought that 256 (85%) of the surviving patients at the time of the first study did have angina. By the second study the doctor thought that 213 (79%) of the surviving patients (213 of these 51 had no angina in the last year but the histories of 28 were highly suggestive of continuing angina, though "not severe" angina. Seven of these 51 patients had had cardiac surgery.

Over the four years, 38 (12.4%) patients died, 28 men and 10 women (table 1). The angina in those who had died affected their daily lives for all the factors to a much greater extent compared with the 269 known survivors. Eighteen (47%) had severe angina compared with 55 (19%) of the survivors. This was highly significant ($p < 0.001$). No significant differences were found in 1979 between those who died and those who survived, with regard to smoking habits or medication, nor in the percentage who had had a myocardial infarction. One patient died during coronary artery bypass surgery.

Table II gives the medication taken at the time of the first study and the drugs that were introduced or discontinued during the study (excluding patients who died and those lost to follow up). No patient was recorded as starting or discontinuing a drug if the drug was used being changed to another in the same therapeutic group. Thirty seven (14%) patients were taking no

"Severe" angina group at first study ($n=63$)

Outcome of second study, excluding the 29 lost to follow up and 16 with severe angina not known

Changes in severity of angina. Figures are numbers of patients.

TABLE 1—Causes of death of 38 patients

Myocardial infarction	Coronary heart failure	Coronary artery bypass surgery	Cardiomyopathy	Stroke	Other	Total
18	15	1	1	1	1	38

TABLE 2—Drug treatment of 307 patients, excluding those who died and those who were lost to follow up

Drug	No. % of patients taking drug at first study	No. % of patients taking drug at second study	No. % of patients taking drug at second study	Total No. % of patients
Controlled treatment	179 58	11	3	190 61
Long acting trinitrate	14 5	21	7	35 11
Calcium antagonist	18 6	61	21	79 26

medication compared with 28 (8%) in 1979. Fifty seven (41%) of those patients who were taking glyceryl trinitrate in the follow up study were taking it prophylactically, compared with 75 (33%) in the first study. Table III gives the reasons for discontinuing a particular medication. There were recorded only for the 51 of the 73 patients who discontinued blockers and only for 19 of those who stopped glyceryl trinitrate, but were recorded for all but one of those who stopped taking long acting trinitrate preparations and for all who stopped taking calcium antagonists (table III). The large number of reasons for discontinuing drugs arises because some patients were changed from one drug to another in the same therapeutic group. For example, of the 37 patients recorded as discontinuing calcium antagonists, one patient had stopped the calcium antagonist but was now on another.

TABLE III—Reasons for discontinuing drugs, excluding those who died and those who were lost to follow up

Drug	No. % of patients discontinuing drug	No. % of patients discontinuing drug	No. % of patients discontinuing drug	Total No. % of patients
Controlled treatment	6 2	4	1	11 4
Long acting trinitrate	14 5	21	7	35 11
Calcium antagonist	18 6	61	21	79 26

From 1979 onwards 73 (23%) of the surviving patients were under hospital care and 12 (4%) of these remain. Table IV gives the investigations carried out and the number of patients who had coronary artery bypass surgery, while Table V compares the severity of patients' angina in 1979 and 1983. The 16 patients who had had coronary artery bypass surgery, and survived, 11 (69%) had had a previous myocardial infarction. A further three patients had had a myocardial infarction between 1979 and 1983. When compared with the remaining patients but excluding those who died and those who were lost to follow up only 12 (49%) had had a myocardial infarction; this was significant ($p < 0.05$). The severity of angina at the first study was not significantly different from that found for all patients. By the second study eight (10%) patients had been free from angina for at least a year from the date of their surgery, in comparison with the remaining patients; this was highly significant ($p < 0.001$). No improvement in state of employment was found.

TABLE IV—Investigations and surgery, excluding those who died and those who were lost to follow up

Investigation	Number of patients	Number of patients	Number of patients	Total No. % of patients
Controlled treatment	6	4	1	11
Long acting trinitrate	14	21	7	35
Calcium antagonist	18	61	21	79

TABLE V—Severity of angina related to investigations and surgery, excluding those who died and those who were lost to follow up

Investigation	Number of patients	Number of patients	Number of patients	Total No. % of patients
Controlled treatment	6	4	1	11
Long acting trinitrate	14	21	7	35
Calcium antagonist	18	61	21	79

Over the four years 22 patients had myocardial infarctions, two patients suffered two each. Ten of these patients had not had a myocardial infarction before. There was no appreciable change in smoking habits, though the trend was downward—107 (40%) were smoking in the first study and 98 (36%) in the follow up study.

In the first study 12 patients were found to have diabetes mellitus, and at that time there was no difference between these and the rest of the patients. By the second study a further five patients were found to have diabetes. Four

of the original 12 had died and three had had a myocardial infarction, one had had coronary artery bypass surgery. By the second study, 44 patients stated that they had not had angina in the previous year. When they were seen at the time of the first study these patients were younger and their angina was much less severe compared with the remaining patients in all variables; only one patient had severe angina. By 1983 eight (18%) of these patients had had coronary artery bypass surgery. The doctor thought that 20 (46%) still had angina and 14 (32%) still took antianginal medication. Most patients were more active than in 1979, though this was not statistically significant.

Discussion

Angina is still a severe and disabling condition for many people despite recent advances in both drug treatment and surgery. As in the previous study the daily lives of many patients are greatly affected, with a continuing high morbidity and mortality.

The deaths in 34 of the patients were cardiovascular in origin, giving an annual mortality rate of 2.8% compared with an expected rate of 0.3% in the same age group for England and Wales. Fry reported an annual mortality rate of 4.6% but the average age was greater. There was no difference in the mortality rates between men and women. This is at variance with the Framingham study, suggesting a better prognosis for women, but it is in agreement with Oliver, who found a 30% mortality over eight years in women aged under 45.

Of the survivors, 44 (16%) stated that they had not had angina in the last year. This remission rate is comparable with 18% reported by Fry and with the 14% and 19% of women with previously persistent angina in the Framingham study. Suggested mechanisms of remission have been: reduction of daily activity, an interim myocardial infarction destroying ischaemic muscle, and development of collateral circulation. As the patients seemed to be at least as active and had fewer myocardial infarctions, the first two seem unlikely. In 1979 their angina was much less severe, perhaps suggesting that they had less pronounced coronary artery disease, which enabled even a limited amount of revascularisation to free them from angina for a time at least. Eight of these patients had had coronary artery bypass surgery.

The disparity of the 28 patients who still had anginal symptoms despite the doctor's belief to the contrary seems most likely to be due to the fact that as their angina was not severe, they no longer consulted about it. Not unexpectedly, those who were first seen with "severe" angina had the worst prognosis, with 18 (29%) dying. None the less, 30 (48%) had improved, though they still suffered from angina. As in those who are now free from angina, there was no evidence that this was due to increased medication or reduction in daily activities or in smoking. The improvement was therefore presumably due to some revascularisation.

Those with diabetes seem to have fared particularly badly, with a third of those who were originally diagnosed dying. Though the numbers are small the trend is as expected and suggests that all younger patients with diabetes and angina should be referred for a cardiological opinion.

The extent to which angina affected the daily lives of patients was broadly similar in the two studies. Sadly, smoking habits had changed only a little. The most noticeable change was in employment, dropping from 175 (69%) to 99 (37%) in 1983. Accepting that this must partly be due to patients being four years older, those with less than perfect health are undoubtedly the first to be let go in a recession.

In 1979, 252 (75%) patients were taking β -blockers; by 1983 a further 14 (5%) had been prescribed such a drug. Though it is not known whether it was being used in full dosage, it seems that most of those who might benefit have received the drug. The figures are similar for glyceryl trinitrate, though still only a minority of patients is using glyceryl trinitrate prophylactically. The use of calcium

antagonists and long acting trinitrates has more than quadrupled in the four years, reflecting current interest. Twenty six (13%) patients stopped taking calcium antagonists because of side effects or lack of benefit. The numbers are small, however, and another study would be needed to clarify this.

Since 1979 only 73 (23%) of the surviving patients had been under the care of the hospital specialist specifically for their angina. Furthermore, only a few patients (table IV) had undergone detailed hospital investigation, such as an exercise electrocardiogram, despite this being a young population, in most of whom there was appreciable morbidity and mortality. The numbers are too small to be significant, but, of those with severe angina in 1979 who were investigated, over half had sufficiently severe coronary artery stenosis to warrant surgery. The outcome suggests that at least those with severe angina, together with diabetic patients who have angina, should have an exercise electrocardiogram. Raftery *et al.* showed that a properly conducted exercise electrocardiogram can be highly predictive of important coronary arterial lesions.

The results of clinical trials of surgical versus medical treatment in angina have shown that surgery improves morbidity and in those with severe left main branch stenosis or three vessel disease it may reduce mortality, though this is not clear.

In this study the morbidity was less in patients who had surgery compared with those who did not. Only a further 13 (5%) patients underwent coronary artery bypass surgery during the study interval. Using the high reporting practices as defined in the previous study, this gives an annual rate of 35 million. There may be some patients in our study population who have since developed angina for which they have had surgery. This rate may therefore be slightly low. In the Northern region few patients outside the age group studied are operated on. The reported rate in 1982 for the metropolitan regions was 169 million and for the rest of England and Wales 47 million. The questionnaires did not indicate why so few had been referred to the specialist for further investigation, but in discussions with some of the participating doctors it was apparent that they thought that the unavailability of such facilities in the region rather than the attitudes of the general practitioners. More can obviously be done, particularly in relation to smoking, and perhaps in the further use of drugs, but for those with severe angina greater use of surgery must play a greater part at present.

I thank Professor Julian, Freeman Hospital, Newcastle upon Tyne, the Research Committee, Northern Region, Royal College of General Practitioners, and Miss Vivienne Mait of the statistical department, Northern Regional Health Authority, for their help and advice. Mr A. Clare for secretarial work, and all the general practitioners who participated in the study. The study was supported by a grant from the Northern Region Health Authority.

References

1. Committee on Coronary Disease. *Coronary Disease in Great Britain: A Report of the Committee on Coronary Disease*. London: HMSO, 1978.
2. Fry J. Natural history of angina pectoris. *Br Med J* 1978; 2: 102-104.
3. Raftery J, Bland J. Natural history of angina pectoris. *Br Med J* 1978; 2: 102-104.
4. Oliver J. Ischaemic heart disease in young women. *Br Med J* 1978; 2: 102-104.
5. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.
6. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.
7. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.
8. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.
9. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.
10. Raftery J, Bland J. Remission of angina pectoris. *Br Med J* 1978; 2: 102-104.

Accepted 21 February 1985

Factors associated with home visiting in an inner London general practice

D BUCQUET, B JARMAN, PAT WHITE

Abstract

We decided to examine the services provided by doctors in an inner London practice for domiciliary care. It was expected that the study would highlight the most relevant questions and variables related to access and up-to-date use of these services; it would also contribute to the design of an accurate procedure for auditing the pattern of delivery of home care to be conducted in the practice in the future.

During the study period, 1976-81, there were 90 500 doctor-patient contacts. For patients up to the age of 10 years the proportion of home visits was 9.2%, falling to 2.2% in the age group 20 to 29, then there is a quasi plateau till the age of 60. After 60 the proportion of home visits doubles in each of the following 10 year age groups, reaching 54% in the over 80s. The proportion of home visits (standardised by age) rises from social class II (10%) to social class V (10.9%), but is higher in social class I (11.7%).

The proportion of home visits according to distance from the practice rises from 8.2% near the health centre to 9.6% at a distance of 0.25 to 0.50 mile, and drops to 8.8% beyond 0.75 mile. The distance effect is not consistent when the social class dimension is added: social classes I and II have higher proportions of home visits in certain age and distance groups. Single people have the lowest proportion of home visits (6.8%); there are large differences between men and women among widowed (14.1% and 8.6% respectively) and divorced or separated (7.0% and 10.7% respectively) patients. There are important variations in the proportions of home visits made by the doctors in the practice, the trainees carrying out proportionally many more home visits.

Data collected in the practice can be used to define specific issues for future audit exercises. Furthermore, socio-demographic characteristics of patients have been shown to be associated with use and access to medical services.

Introduction

The declining home visiting rate of general practitioners in the past 10 years is well documented in England as well as in the United States, where the alleged poor cost effectiveness of this service was a strong incentive for reducing it. Time spent travelling in the car is largely wasted, and the availability of equipment and other resources argue in favour of the patient coming to the doctor rather than the reverse. General practitioners have expressed the fear of home care of patients being taken over by other professionals if this declining trend persists, there is a claim of "total" responsibility for family care. "If the profession is failing to meet a public need society will find some way of meeting the need, if necessary by turning to a group outside the profession."

Nowadays, rehabilitation in the home, preadmission home visiting for geriatric patients who have been referred for care as inpatients, and the whole person approach to a patient's management are in vogue. "The questions of 'access' and 'utilisation' of

medical services (overutilization and underutilization) have also been discussed; perhaps these terms should no longer be exclusively to the planners' vocabulary but be applied to primary care providers; it might enable them to rationalise their patterns of delivery of care on need rather than on demand to achieve greater equity of access, although the assessment of need and of appropriate use relative to need is not simple. This would lead to a selection of criteria for home visits. "Learn to respond to the need not to the demand," wrote Caution.

We thus decided to examine the services provided by doctors in an inner London practice for domiciliary care. We expected that in highlighting current practice the study would yield clues for selecting questions and criteria to audit in the future the pattern of delivery of domiciliary care, with emphasis on accessibility and usage.

Method

The practice is one of a group of three located in a health centre in inner London that has access through the department of general practice, St Mary's Hospital Medical School, to computer facilities for storing and analysing the data on the registration and consultations of all its patients. The practice had three full time partners and a trainee during the study; the list size was 5800 patients in 1976 and 7100 in 1981. The data that we analysed related to all doctor-patient contacts between 1976 and 1981. The registration data for each patient included age, sex, marital status, social class according to the Office of Population Censuses and Surveys, classification of date of onset of practice, and distance of residence from the health centre. The consultation data included up to two diagnoses for each consultation, whether or not a prescription, a certificate, or a follow up appointment was given, and details of up to three investigations and one referral for each diagnosis.

We studied the proportions of home visits according to age, sex, social class, distance from the surgery, marital status, and doctor; the proportions were standardised by age to the population of England and Wales (1981), using the direct method. The denominators were the total numbers of contacts in the relevant cells. The statistical analysis was performed with the Statistical Package for Social Sciences. The relations between the variables were tested with the χ^2 test.

Results

During 1976-81, 90 500 patient contacts, consultations and home visits, were coded and stored on computer. The results refer to home visits only.

Age, sex, social class, distance, marital status, and doctor

The effect by age on home visits is clear (fig. 1). The proportion of home visits in the age group up to 10 years is similar to the overall practice average (9.2%); the lowest proportion of home visits, 2.2%, is a characteristic feature of the age group 20 to 29, increasing to 10.7% in the age group 50 to 59, doubling in each of the following 10 year age groups to reach 54% in the over 80s; $p < 0.001$. Men had a higher proportion of home visits than women as a whole, 9.8% v. 8.5%, $p < 0.001$, which, after standardising for age, is 8.4% for men and 8.8% for women.

The relation between home visits and social class shows an increase in the proportion of home visits from social class II (10%) to III (8.5%), to IV (9.2%), to V (10.9%), but the proportion is highest in social class I (11.4%). With regard to distance, the proportion of home visits rises from 8.2% near the health centre (0-0.25 mile) to 9.6% at a distance of 0.25 to 0.5 mile, dropping to 8.8% at 0.50 to 0.75 mile and 8.8% when the distance is over 0.75 mile.

Figure 2 shows the variations in the proportion of home visits according to marital status. Single people had the lowest proportion of home visits. There

Department of General Practice, St Mary's Hospital Medical School, London.

DENIS BUCQUET, MD, epidemiologist, visiting research fellow.
BRIAN JARMAN, MB, MRCP, professor of primary health care.
PAT WHITE, BSc, research assistant.

Correspondence to: D. Bucquet, St Mary's Hospital Health Centre, London NW6 6EG.

are large differences between men and women who are widowed and separated.

The variations in the proportion of home visits made by doctors in the practice (partners and trainees) are important (fig 3), the range being 6.6% to 15.5%. In the age band 45 to 64, for example, the proportions of home visits for each doctor are: 6.7%, 5.2%, 13.8% (trainee), 5.5%, and 10.6%.

Combination of variables

Age, sex, and social class—A detailed analysis (table 1, fig 4) shows higher proportions of home visits in social classes I and II for girls aged 0-4 ($p<0.001$), boys aged 0-4, 5-9, and 10-19 ($p<0.001$), women aged 20-29 ($p<0.001$), and men aged 30-39 ($p<0.05$) and 70-79 ($p<0.001$). Through out the ages in the different social classes the higher proportion of

FIG. 1—Proportion of home visits by age and sex.

FIG. 2—Proportion of home visits by marital state, standardized by age.

TABLE 1—Proportion of home visits by age, sex, and social class

Social class	Age group years									
	0-4	5-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+
I	10.1	11.1	17.1	10.6	10.1	1.1	2.1	7.9	6.0	1.0
II	12.4	12.8	15.4	10.1	4.4	2.9	1.5	1.2	1.6	4.1
III	6.0	7.2	10.0	10.0	4.1	2.6	1.1	1.1	2.1	1.1
IV	10.8	6.1	7.1	9.2	4.6	4.1	2.2	4.1	2.7	1.8
V	6.2	6.0	6.0	5.1	4.1	1.4	1.9	2.7	2.6	4.6

* $p<0.05$, ** $p<0.01$, *** $p<0.001$, NS, not significant.

TABLE 2—Proportion of home visits by marital state, age, and sex

Marital state	Age group years									
	0-4	5-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+
Single	2.6	1.4	7.6	5.2	8.1	9.9	24.2	41.7	41.7	41.7
Married	1.4	1.9	9.0	8.1	17.1	12.1	48.6	49.7	49.7	49.7
Widowed	22.1	8.1	28.1	28.1	14.8	17.4	41.8	41.8	41.8	41.8
Divorced/separated	1.0	1.9	1.9	1.1	6.6	20.1	36.1	52.1	52.1	52.1

* $p<0.05$, ** $p<0.01$, *** $p<0.001$, NS, not significant.

FIG. 3—Proportion of home visits by age and sex, standardized by social class.

Doctor, social class, and age—Table IV shows further variations in the proportions of home visits. The highest proportion of home visits is a characteristic of the trainee practice throughout the age groups and social classes (except for social classes I-II, age 75 and over, seen by Dr 4, and social class III, age 45-64, seen by Dr 4).

Discussion

It is relevant to analyse aggregated data over a five year period when a review of published reports suggests that there has been an important decrease in home visits in the past 10 years.¹ The reductions reported are 40% in 1964-77, and in 1969-80, 15% in 1965-78, and 21% in the General Household Survey (1972-83). We checked for this by calculating standardised rates of home visits for

FIG. 4—Proportion of home visits by doctor, standardized by age.

TABLE 3—Proportion of home visits standardized by age, by distance from the health centre, and by social class

Distance from health centre	Social class									
	I	II	III	IV	V	VI	VII	VIII	IX	X
0-25 m	12.1	10.1	11.0	10.1	10.1	10.1	10.1	10.1	10.1	10.1
25-50 m	12.1	10.1	11.0	10.1	10.1	10.1	10.1	10.1	10.1	10.1
50-75 m	12.1	10.1	11.0	10.1	10.1	10.1	10.1	10.1	10.1	10.1

home visits in men is not consistent. For example, women had a higher proportion of home visits than men in social class I, age 20-29 ($p<0.05$), in social class II, age 70-79 ($p<0.001$), and in social class V, age 60-69 ($p<0.001$).

Distance, age, and social class—Here again the pattern of home visits among patients in social class I is different, especially in the immediate surroundings of the health centre. Table II shows clearly the higher proportion of home visits in social class I, particularly in the two extreme distance categories. The gradient between social classes is clear from II to V in the third distance category (0-50 m) and from III to V in the second distance category (0-25 m). The last distance category (over 75 m) is characterized by high proportions of home visits in social classes I and IV and a low proportion of home visits in social class V. Merging social classes I and II, in the age group 75+, living less than 0.25 mile from the health centre, the proportion of home visits in social classes I-II is 65.8% (47.5% in social class V ($p<0.001$)). 15 year olds and under in social classes I and II have a higher proportion of home visits: 11.4% v 5.9% in social class V for distances between 0.25 and 0.50 mile and 12.5% v 5.7% for distances between 0.50 and 0.75 mile.

Marital state, age, and sex—A striking finding is the differences between men and women, men having a higher proportion of home visits in all groups except for those aged 65-74 who are divorced or separated and after the age of 75, whatever the marital state (table III). The addition of the social class dimension brings out no consistent pattern of association between proportion of home visits and marital state.

TABLE 4—Proportion of home visits by doctor, social class, and age

Doctor	Age group years									
	0-4	5-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+
Dr 1	10.1	11.1	17.1	10.6	10.1	1.1	2.1	7.9	6.0	1.0
Dr 2	12.4	12.8	15.4	10.1	4.4	2.9	1.5	1.2	1.6	4.1
Trainee	6.0	7.2	10.0	10.0	4.1	2.6	1.1	1.1	2.1	1.1
Dr 4	10.8	6.1	7.1	9.2	4.6	4.1	2.2	4.1	2.7	1.8
Dr 5	6.2	6.0	6.0	5.1	4.1	1.4	1.9	2.7	2.6	4.6

* $p<0.05$, ** $p<0.01$, *** $p<0.001$, NS, not significant.

FIG. 5—Proportion of home visits by doctor, standardized by age, social class, and sex.

each year in the period covered by the study and standardised rates of patient contacts. Table V shows the results. Although this shows a slightly decreasing trend, and thus it is interesting information for the practice, it is not far from the 13% found by Cartwright and in 1977.¹ We thought that it was reasonable therefore to attempt to identify factors related to home visits that are not obvious through the daily clinical observation to consider setting up criteria for the delivery of this particular service and the need for implementing a more targeted home visiting policy.

The accuracy of the data has been checked on a random sample of 500 records and was found reasonable; less than 1% of the data on age and sex, 2.5% of that on marital state, and less than 5% of that on social class were wrongly classified. The proportion of unrecorded home visits was subjectively estimated by the doctors in the practice to be around 5%.

The differences in the proportion of home visits by social class and distance are not different from those found by Cartwright and Anderson in 1977, in which there was a clear cut social class gradient explained mainly by differences in the availability of private transport—... patients being less likely to go to their doctor by private transport and therefore ask for more home visits.¹ This does not seem to apply in this practice, as 50% of patients in social class V on the list live within a quarter of a mile of the premises (80% within half a mile). Unfortunately, car ownership is not a variable available in our registration data, but it is available from the census or the General Household Survey. Figure 5 shows the variations of car ownership by socioeconomic groups (General Household Survey, 1971 and 1982). Some evidence has been discussed regarding the relation between car ownership and state of health and take up of medical services.¹ Although crude distance is a less useful measure of access than time spent travelling,¹ the influence of distance, in the initiation of care until symptoms are clear and demand immediate attention, is highlighted by Knott, Weiss and Greenfield, and Parkin.¹ Distance affects the medical

FIG. 6—Proportion of home visits by doctor, standardized by age, social class, and sex, for Great Britain.

care process differently by social class and interacts with social class as an explanatory variable.¹

Parkin also raises the issue of the cost of journey time and travel, which does not represent the same constraints for each social class. If one assumes that the cost to the patient of a home visit is less than the cost of going to the surgery it might be argued that arranging more home visits for patients in the lower social classes would counteract inequalities in income. Level of education influences the way in which patients use doctors in the sense that more educated patients tend to conform to the pattern expected by the medical profession, as reflected by a lower propensity to consult for trivial illnesses.¹ The inappropriate anxiety of the patient or of the person who initiates the call for a home visit is also linked with level of education.¹

In the opposite direction is the fact—minor in terms of workload and activity—that the highest proportion of home visits is made to patients in social class I who live in the immediate surroundings of the health centre. It might illustrate the easier class to class relations experienced by some doctors, corroborated by the longer duration of consultations with patients in social class I (practice data, unpublished). Home visits to patients in social class I might be considered more attractive than visits to patients in other social classes. Other variables such as how long a patient has been on the list, quality of the doctor-patient relationship, and emotional input might explain what appears to be a "special service" to patients in social class I. The argument of the unattractiveness of home visits (because of travelling time or lack of facilities available at the surgery)¹ is not always the rule. The higher social classes also handle communication systems more easily; this can work in two ways: either to explain their complaint more clearly and then receive advice that seems professionally satisfactory to the practitioner or to order a home visit as they would order any other service.

Marital state seems to be a relevant variable to include in the construction of a policy for domiciliary care. Further analysis on larger or multicentre data sets is required to assess its additional contribution in the delivery or uptake of home visits. The differences between men and women in the widowed category and between single, divorced, and separated people as opposed to married or widowed people are interesting as no published information seems to be available on this.

The variations among partners in the practice are a further indication of the idiosyncratic behaviour of practitioners, as shown recently by Hartley et al in test ordering.² The variations between trainees and partners confirm what Harris et al found when

analysing the prescriptions of doctors according to their status—trainees prescribing mainly for acute onset of conditions with very low involvement with patients with chronic conditions.

Gray underlined the privilege that home visits provide in allowing doctors to observe their patients on their own territory.³ Young et al pointed out that home visits place the doctor in the patient's domain, therefore shifting the relationship to one of more equal status and allowing the patient to "bring in" the doctor rather than the reverse.⁴ Guy et al and Guyan Schuman showed how home visiting allows the doctor and patient to "get acquainted and establish communication, to facilitate observational skills and awareness of the community (cross cultural aspects), to improve research parameters in the family record."⁵ Clark and Berolzheimer emphasized the "great informational value" of home visits, particularly in assessing the emotional dynamics of the family.⁶ Young et al think that home visiting helps to develop a holistic view of the patient, the ability to evaluate the patient from a psychological perspective, and a treatment plan that integrates psychosocial and medical issues, and helps to teach the values of a multidisciplinary approach to medical care.

The variability in the delivery of domiciliary care provided by the practice has been shown in terms of the behaviour of both providers and consumers. This raises the question of how the practice might use this information: it helps the doctors to review their behaviour to bear in mind this differential pattern of delivery of home care. But they strongly feel that they work in a system based on demand, and the problem of providing access to home visiting based on need (assessed by the practitioner or by consensus on need reached by both practitioner and patient) is delicate and solving it would be a long term educational process.

This is an example of the use of practice data collected systematically; it provides information and questions (appointing a visit initiator, for instance), which might be used to design accurately a purposive audit procedure on home visiting.

We thank Drs Constantinos, Elder, Levi, Read, and Webb, who were also concerned in this study, and Mrs Lewis and Mrs Stephan, who helped with the recording.

References

1. Harris J. Home visiting as a rural practice. *Br Med J* 1981;282:1154.
2. Gray TJ. Practice as a home visit. *J R Coll Gen Pract* 1978;28:617.
3. Young TJ. The home visit—what is it? *J Fam Pract* 1981;13:181-2.
4. Clark WJ, Berolzheimer N. The home visit. *Fam Med* 1981;13:131-9.
5. Guy A, Guyan S. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
6. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
7. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
8. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
9. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
10. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
11. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
12. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
13. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
14. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
15. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
16. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
17. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
18. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
19. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
20. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
21. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
22. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
23. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
24. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
25. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
26. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
27. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
28. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
29. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.
30. Young TJ. The home visit: a review of the literature. *Br Med J* 1981;282:1154.

Vocational Training

Vocational training schemes for general practice: what happens to those not appointed?

A J BALFOUR

General practice is a popular career for many doctors, and most want to prepare for it by entering a three year linked vocational training scheme. Since the Vocational Training Act was passed there has been a great increase in the numbers of doctors applying for schemes, without any increase in the number of places available. This leads doctors to make multiple applications until they are appointed to a place on a scheme. But what happens to those who are not appointed to three year schemes? Can they obtain suitable training posts to prepare them for general practice?

To find out I made a survey of the unsuccessful applicants to the Bedford Vocational Training Scheme in 1984. Advertisements were placed in the *BMJ* on 7 April, the closing date was 18 April, and 132 applications were received. Interviews were held on 3 May 1984, when two doctors were appointed to the scheme. The following week letters of rejection were sent to the 130 unsuccessful applicants, enclosing a questionnaire asking what posts they had eventually obtained. In July 1984 the same questionnaire was sent to all those who had not replied, with a stamped, addressed envelope: 73 replies were received, which is a response rate of 56%.

I had originally sorted the written applications into 46 "possible" and 84 "less suitable" candidates. Subsequent questionnaires were received from 58% of the "possible" candidates and 52% of the "less suitable" candidates, but I cannot exclude any hidden differences between responders and non-responders. Sixty five doctors still intended to enter general practice, but eight (11%) had changed their minds. Three of those who had obtained posts in anaesthetics, two in psychiatry, and one in medicine and paediatrics. All the doctors had made multiple applications for posts, although 15 had made fewer than 10 applications; 16 doctors had made from 11 to 20 applications and 33 had made from 21 to 50; seven had made over 50 applications.

After all that effort 20 doctors (27%) had been appointed to places on three year linked vocational training schemes, of which only one was in the same region as Bedford. Applicants can no longer be too particular where they think the difficulty lies in getting their training where they want. Forty four doctors (66%) had obtained posts as senior house officers, which could count as equivalent experience: 15 were in accident and emergency, nine in medicine, nine in psychiatry, two in ear, nose, and throat, two in paediatrics, one in obstetrics and gynaecology, and three in anaesthetics. Obviously, paediatrics and

obstetrics are difficult specialties in which to obtain experience, even though they are so valuable as preparation for general practice. Two doctors had been appointed as trainees in general practice, although they had apparently not yet completed sufficient post registration hospital experience to gain certificates of prescribed experience. Seven doctors (10%) stated that they would be unemployed in August 1984. Measuring unemployment is notoriously difficult, and a 56% response rate may include a disproportionate number of disappointed doctors.

It has been suggested that multiple applications might be avoided if there was a common clearing house for applications similar to the Universities Central Council on Admissions (UCAA). This would greatly reduce the number of forms that applicants would need to complete and require fewer days off work to attend interviews. Applicants, however, would have to be careful in choosing their schemes to avoid too many over-subscribed schemes and apply for some schemes because of alleged unpopularity rather than a real desire to join. The advantage to the schemes might be fewer applications, but good candidates who had put a scheme low on their list might be missed. In the questionnaire they were asked their opinion of this—32 (44%) were in favour of it, though this was most noticeable among those who obtained posts as senior house officers. Sixteen were definitely against the idea, and a further 14 uncertain. Their remarks indicated that it would restrict the number of applications that a candidate could make and not improve their chances of employment unless there was a safety net.

Most comments that were made on the questionnaire complained of the large number of applicants for a small number of posts and expressed a desire for more places to be made available. This presupposes, however, that there are not sufficient vacancies in general practice for those who complete their vocational training. Some doctors feared that posts had already been promised to local doctors and pledged for a far chance for outsiders. They also commented that it was difficult to put forward their qualities when all candidates had completed only two house officer posts, and thus shortlisting relied heavily on the presentation of their curriculum vitae. Some candidates wanted to visit the scheme in advance but found it difficult to identify who the local consultants and trainers were or to ascertain whether a visit would be appreciated.

I believe that this small survey of the future employment of unsuccessful applicants to one vocational training scheme confirms the competition for such places but that it is encouraging in showing that 27% will get on to a linked scheme and 60% will start equivalent training.

Accepted 20 March 1985.

1 Golding Road, Bedford MK40 3NF
A J BALFOUR, MRCP, course organizer