PRACTICE OBSERVED

Practice Research

Computer assisted screening: effect on the patient and his consultation

MIKE PRINGLE, SALLY ROBINS, GEORGE BROWN

Abstract

The initial impact of computer assisted preventive screening in general practice consultations has been monitored. The technology has not been found stressful by patients, and the power of the consultation to alleviate low arousal has been increased by computer use. No appreciable increase in the duration of consultation was detected, despite an average computer initiated input of two minutes eight seconds. The computer has successfully prompted preventive screening and health education with a sixfold increase in the number of potentially relevant procedures being mentioned.

The actual information presented by the computer has been shown to be crucial, with the terminal's mere presence an ineffective reminder. The computer terminal was used in 65% of the consultations for which it was available, which, if sustained, represents an effective screening programme for attending patients.

Introduction

The doctors at the Colliingham Health Centre have designed a computer system, which allows interactive use during the consultation. The patients' files contain extensive demographic and preventive medicine information and the terminal is used primarily as a reminder of individually appropriate opportunistic screening.

Computers are being used more widely by general practitioners, although only a small minority use terminals during the consultation. The evaluation of the effects of such use on the dynamics of the consultation has been suggested before computer aided consultations become commonplace.¹

Attitude measurement has already identified concerns among some patients over computerisation in general practice. The first such survey received wide methodological criticism after claiming that many patients react strongly against computers—67% said that they would change their general practitioner. Another survey found that 23% of general practice patients would be uncomfortable in a doctor's surgery if they saw a computer. We found that 83% of patients were generally in favour of medical computerisation, but 31% had reservations concerning loss of confidentiality with the advent of general practice computing.²

The present study meant evaluating the actual effects of computerising medical consultations rather than the "blind" attitudes of patients. The low vocabulary "mood adjective checklist" (MALC) stress and arousal scale has previously been outlined and has proved sensitive by showing the effect of a second doctor sitting in during training consultations.³

It is a fast and simple psychometric test, which has shown that about two thirds of patients consulting a general practitioner enter the consultation with a higher level of stress than when they leave. Similarly, most patients leave the consultation with a higher level of arousal than that measured before the consultation, although the prevalence of arousal increase has shown more variation—between 52% and 72% after "control" consultations in different studies. The effect of most consultations in reducing stress and increasing

an average of 48 seconds, two minutes and eight seconds were occupied by the CPMS. No significant difference between consulting times in any group for any doctors were detected by a test.

PREVENTIVE CONTENT OF THE CONSULTATION

Table IV shows that computerised consultations resulted in a large increase in the mention of three preventive procedures: cervical smears, tetanus immunisations, and blood pressure recordings. Of the possible mentions (table V) of these measures in each group, 12.7% were mentioned in group 1, 18% were mentioned in group 2, 78.6% were mentioned in group 3 ($p < 0.001$). In group 3, 23% of the smears mentioned were actually done or arranged, 64.7% of tetanus immunisations mentioned were done or arranged, and 90.9% of blood pressures mentioned were recorded or arranged for a future visit. Preventive measures not done or arranged were either established not to be due or applicable, or they were referred by the patient or deferred for the patient to "think about".

TABLE IV—Number of preventive and educational procedures mentioned during consultations in each of the three groups

Items mentioned by the computer	Group 1 (n=10)			Group 2 (n=21)			Group 3 (n=19)		
	DA	DA	DA	DA	DA	DA	DA	DA	DA
Cervical smears	0	0	0	0	0	0	4	4	4
Tetanus	0	0	0	0	0	0	11	11	11
Blood pressure	0	0	0	0	0	0	11	11	11
Items not mentioned by the computer	0	0	0	0	0	0	0	0	0
Weight	0	0	0	0	0	0	0	0	0

TABLE V—Observed and possible mentions of computer initiated items: cervical smear blood pressure tetanus

Items mentioned	Group 1 (n=10)			Group 2 (n=21)			Group 3 (n=19)		
	DA	DA	DA	DA	DA	DA	DA	DA	DA
Cervical smears	0	0	0	0	0	0	4	4	4
Tetanus	0	0	0	0	0	0	11	11	11
Blood pressure	0	0	0	0	0	0	11	11	11
Weight	0	0	0	0	0	0	0	0	0

When the computer was used there were fewer mentions of health education measures about which it has no information or prompt facility, such as weight and smoking. A Fisher's exact probability test showed that this was not significant, but it indicates nevertheless that the actual information presented is important, and that the machine's presence does not act as an intrinsic prompt to doctor to "think prevention". This is confirmed by the similarity between groups 1 and 2.

Discussion

In view of the estimated 75% of patients who consult their general practitioners each year (the promotion of prophylactic and other screening measures during consultations has wide reaching potential in the maintenance of healthy patients). If computer use continues in 65% of consultations a large proportion of the practice population could be screened each year. Patients who rarely or never attend surgery, however, will inevitably continue to be an ill screened population. Clearly the benefits of this method of prevention must be weighed against other changes in the consultation, such as duration and patient anxiety.

Surveys assessing attitudes towards computerisation in general practice have provided useful subjective guidance on potential areas of concern to patients. The actual use of computers in general practice, however, has been unfamiliar to all respondents participating in such surveys. Patients participating in the present study have been introduced to the use of general practice computing and a more objective measurement of effects has been made.

This study was carried out in the first months after computer

terminals were installed. Their presence was still unfamiliar to patients and, to an extent, doctors. Computer use was limited to the presentation of reminders of deficits in the patients' preventive medicine cover, resulting in discussion or treatment of the items shown.

Survey results may have led us to anticipate a stressing effect and this has been supported by a previous study, which showed that computers used in the presence of patients in a more diagnostic capacity in a hospital outpatient setting produced undesirable effects on the level of patient's postconsultation stress.⁴ A diagnostic mode of use, however, may reinforce some of the patient's greatest fears—that computers may replace skilled practitioners or that computer errors will prove detrimental to their health. In this study computer use had no effect on patient stress.

No marked effect on the arousing potential of the consultation was seen when computers were used in hospital outpatient depart-

ments. The computerisation of general practice consultations, however, produced arousal increases in patients more consistently. As this was, for all participating patients, the first encounter with doctors using computers some of this may have been attributable to novelty.

Consultation durations showed wide variance for all doctors, but even when the computer was used there was no marked increase in the number of consultations exceeding their appointed seven minutes and 30 seconds.

Doctors differed in their style and extent of their use. Doctors A and B initiated computer use at a definite point, often having established with the patient that all they came for had been dealt with. A question such as "Is there anything else?" preceded computer entry and the discussion or action necessary, or both. Doctor C used his spare moments throughout the consultation to enter the patient's identification and plan on the screen before initiating discussion or action at the end. He was the only doctor to use the computer's prompt for routine urine testing, and the only doctor for whom mean consultation times with computer use were increased by the combined duration of computer use and initiated discussion.

Doctor B showed the longest mean consultation duration for those consultations with the computer available for use, but not in fact used. He also used the computer less frequently than the two other doctors, and may consciously or otherwise have used time as a criterion for deciding whether or not to enter the computer phase. Doctor A used the computer most often and for the longest mean duration. Even so, an average of two minutes and 35 seconds of computer initiated activity added only 14 seconds to his average consulting times. Doctors A and B included preventive care in consultations in this way without a comparable increase in consultation times.

While these results seem favourable some modes of use may increase consultation times or where consultations have apparent extra content for no extra time there may be an increase, as indicated by the slight decline in discussion of the two health education topics, smoking and weight.

The effect of the computer on patient stress and arousal, consultation duration, and the uptake of preventive medicine was highly satisfactory. Most procedures thus included are carried out or arranged (about 65% of those observed here). The only potential

Health Centre, Colliingham, N. Newark, Nottinghamshire

MIKE PRINGLE, MRCP, general practitioner

SALLY ROBINS, MRCP, research assistant

Nottingham University

GEORGE BROWN, MRCP, reader in educational methods

Correspondence to Dr Mike Pringle

1710

BRITISH MEDICAL JOURNAL VOLUME 290 8 JUNE 1985

arousal is thought to be desirable, since stress shows in inverse proportion to memory of consultation content, and performance efficiency improves as arousal increases, until excessive arousal causes declining efficiency. Illness itself combined with seeing the doctor may create a state of relatively high stress and low arousal in patients, which the consultation seems to alleviate.

A prime consideration in the feasibility of computerising consultations is whether or not doctors and patients have the time for such an adjunct to the consultation. Consultation duration is important to doctors, but they at least have some control over the time spent with each patient. Longer consultations may be a greater problem to patients who tend to wait verbal or non-verbal cues from the doctor before leaving the consultation. Some have other appointments or wish to return to work. Others already find waiting times frustrating, and longer consultations imply longer waiting times.

The computer in Colliingham has been used initially to prompt extra medical care by issuing preventive reminders. Doctors have not abandoned the conventional medical records and still use these for the routine consultation content. The computer is not needed as such, therefore, but requires the identification of the patient should doctors wish to add preventive care to the consultation agenda. This extra but unforced use of a computer terminal has provided a good opportunity to observe the extent and duration of computer use.

The intrinsic benefit of preventive care to patients cannot be disputed. Other methods of implementing such care, with and without a computer, have been documented.⁵⁻⁷ For this method of computer use it was thought that the efficiency should be assessed in terms of education and prevention by comparing the preventive measures mentioned and done during consultations with and without the computer present.

Since the presence of a video camera has already been shown to effect neither stress nor arousal it was considered an acceptable method of data collection.³

Method

Over a period of eight months a selection of patients attending morning, afternoon, and evening surgeries were chosen from the appointment book. Where practicable they were chosen according to a random selection of appointment numbers (from 1 to 12), but unsuitable patients were replaced by the next suitable attending patient. A maximum of four patients per surgery were included, but only three proved more manageable. Sixty patients were selected for each of two groups. Equal numbers of men and women patients were chosen for each group; children (under 16 years) were excluded, and no patient was selected more than once.

TABLE I—Number of patients and computer status of groups compared

Group	Computer status		Non-computer	
	Men	Women	Men	Women
1	30	30	30	30
2	30	30	30	30

The 60 patients in groups 1, 10 men and 10 women for each of three doctors, were asked to help by accompanying the experimenter (SR) to an interview room, where they completed the low vocabulary MALC stress and arousal scale. This requires the independent presentation of 26 single word items to subjects, who then "pen" each under the most appropriate of four headings (I definitely feel, I think I feel, I don't think I feel, I definitely don't feel). The patients were told that the doctor they had arranged to see would like to video record their consultation in connection with the research project and they were offered the opportunity to decline. If they did not object written consent was obtained. They were asked to return to the experimenter after the consultation. The doctor was told to record selected patients by a marker in the patient's notes. After the consultation patients completed the stress and arousal scale again and confirmed that video recording had taken place.

A further 60 patients in groups 2 and 3 were treated in the same manner, but during these consultations the doctor had access to a computer terminal on his desk, which he used at his discretion.

BRITISH MEDICAL JOURNAL VOLUME 290 8 JUNE 1985

Doctors A, B, and C are the three male principals in a rural general practice, run from a modern purpose built health centre. On average they use the computer in 65% of those consultations for which it was available (55%, 65%, and 55% for doctors A, B, and C, respectively). The video recording of consultations in which the computer was available for use were viewed and subdivided according to whether or not the computer had actually been used. This gave three groups for analysis.

The three groups were then compared with respect to stress and arousal changes, and the video recordings were assessed for consultation duration; preventive care was included in the consultations.

RESULTS

MEASUREMENT OF STRESS

The changes in stress and arousal before and after the consultation are shown in table II. The three groups were compared using 2 x 2 x 2 tests, presuming the most desirable outcome of any consultation to be decreased stress and increased arousal with any other result constituting a less desirable outcome. There were insufficient data to compare individual doctors, so all statistical tests were applied to the totals for all three doctors.

TABLE II—Changes in stress and arousal before and after consultation

Group	Stress				Arousal			
	Low desirable outcome	High desirable outcome	Low undesirable outcome	High undesirable outcome	Low desirable outcome	High desirable outcome	Low undesirable outcome	High undesirable outcome
1	10	10	10	10	10	10	10	10
2	10	10	10	10	10	10	10	10
3	10	10	10	10	10	10	10	10

Notes: 1. No significant difference between groups 1 and 2; between groups 2 and 3; and between groups 1 and 3.

2. Significant difference between groups 1 and 2; significant difference between groups 1 and 3; 10% ($p < 0.01$), and significant difference between groups 2 and 3; 10% ($p < 0.01$).

There was no significant difference between the three groups for stress changes, but more patients from group 1 returned from their consultation showing higher levels of arousal than in group 1 ($p < 0.01$) and 2 ($p < 0.05$).

TIMING AND STYLE OF COMPUTER USE

Two of the doctors developed a clear cut stage of computer use, which was timed as the "computer initiated preventive medicine stage" (CPMS). The third began with a well defined stage, but developed a more subtle and spread out mode of use. His overall CPMS therefore comprised short sections of input to the keyboard, viewing the screen, discussing the information with the patient, and carrying out any necessary procedures.

As table III shows, while the computer increased consultation duration by

TABLE III—Mean duration in minutes and seconds of total consultation for the three groups and for CPMS computer initiated preventive medicine stage for group 1

Group	DA		DA		DA		DA		Standard deviation	
	Min	Sec	Min	Sec	Min	Sec	Min	Sec	Min	Sec
1	2	42	2	42	2	42	2	42	2	42
2	2	42	2	42	2	42	2	42	2	42
3	2	42	2	42	2	42	2	42	2	42

Notes: 1. No significant difference between consultations in which the computer was used and those in which it was not used; 2. No significant difference between consultations in which the computer was used and those in which it was not used; 3. No significant difference between consultations in which the computer was used and those in which it was not used.

1712

BRITISH MEDICAL JOURNAL VOLUME 290 8 JUNE 1985

area for concern is the possible constricting of the "normal consultation" to accommodate computer use.

We thank the doctors, the staff, and the patients of Colliingham Health Centre for their help with the study, and the Trent Regional Health Authority for funding the research.

References

1. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
2. Brackenbury SM, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
3. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
4. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
5. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
6. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.
7. Pringle M, Jones D, Brown A. Computerisation in the clinic. *Br Med J* 1984; 289: 1053.

Use of medications on the restricted list by the elderly

NORMAN J VETTER, DEE A JONES, CHRISTINA R VICTOR

Abstract

On 1 April 1985 the range of drugs that may be prescribed on the National Health Service was limited by the government. We studied the possible effects of this new policy on 106 elderly patients in two general practices in South Wales. Many patients who were prescribed drugs from the restricted list were also disabled, had psychological problems, or were on supplementary pension. We intend later this year to study the consequences of the changes in prescribing on elderly patients.

Introduction

On 8 November 1984 the Secretary of State for Social Services announced in parliament the government's intention to limit from 1 April 1985 the range of drugs available for prescription on the National Health Service. The statement began:

"The number and cost of prescriptions issued under the NHS has increased very substantially in recent years. In 1983 some 334 million prescriptions were dispensed through the family practitioner service in England at a cost of almost £1400m. This is about 100 million more prescriptions a year than were issued 25 years ago, and the range of drugs prescribed has doubled in the same period from less than 8500 different items to more than 17 000."

"Many of these increases are fully justified and reflect the enormous therapeutic advances made during the last quarter of a century, but expenditure on the present scale can be maintained only at the expense of other parts of the health service. It is now necessary to consider carefully whether all the drugs being prescribed need to be provided by the NHS and whether there are areas in which sensible economies can be made in the drugs bill without detriment to patients."

Research Team for the Care of the Elderly, University of Wales College of Medicine, Cardiff

NORMAN J VETTER, MRCP, senior research fellow

DEE A JONES, MRCP, senior research fellow

CHRISTINA R VICTOR, MRCP, senior research fellow

Correspondence to Dr Norman J Vetter, St David's Hospital, Cardiff CF1 7EL

It then suggested that medicines for the relief of minor symptoms, tonics, cough and cold remedies, antacid laxatives, analgesics for mild to moderate pain, low dose vitamin preparations, benzodiazepine sedatives, and tranquilisers should be restricted to certain brands, some generic, some proprietary.

At the time of the announcement it was thought that such changes would have an effect on the prescribing patterns for elderly people by causing them to alter their medication or to reduce the amount taken, or they might choose to pay for some medicines to keep those that they preferred. As the elderly have previously been exempt from prescription charges this was expected to affect either their use of medicines or their financial state considerably.

This paper describes prescribing for elderly patients in two general practices in South Wales before the limited list was instituted to estimate the likely consequences of such limitation. We intend to examine the actual consequences of the change in detail later in the year when changes in the prescribing pattern have become established.

Method

The study was based in two general practices. The population consisted of 529 persons in one practice and 552 in the other who had been born in or before 1909 and lived at home. They were randomly selected from the age-sex register of the two general practices. All of the patients were selected from a typical rural practice of four general practitioners, covering a large area in southern Powys, 552 patients were randomly selected from a typical urban practice of six doctors in Gwent.

Each member of the study group was visited at home by interviewers and asked a series of structured questions to investigate their physical, mental, and social characteristics. Physical disability was measured using a questionnaire based on a nine item score that was first described by Townsend.⁸ Mental disability was measured using standard questionnaires for anxiety and depression, based on those developed by Bedford and Foulds.⁹ This allocates a score to a series of symptoms and signs that are characteristic of patients with these neurones. The social variables measured included composition of household, frequency of social contacts, use of medical and social services, and the availability of and characteristics of the old person's main carer. The type and quality of housing and subjective feelings on their quality of life were also measured.

Completed responses were obtained from 524 people in Powys and from 542 in Gwent, a response rate of both areas of 90%. Information was obtained about the medication these people had taken in the past 24 hours by checking the medicine containers and repeat prescription cards and by the interviewer interrogating other family members.