Discussion

The usual career for a general practitioner covers about 30 years, so that the total of 31 new doctors becoming new principals in FPC A during the years 1981, 1982, and 1983 is to be expected. FPC B, which was short of doctors during those years, had a higher recruitment rate. This is unlikely to continue, however, as almost all of its area is now classified as either "intermediate" or "restricted". Among the reasons given as to why working in primary care teams is unusual in these family practitioner committee areas are: (a) difficulties in improving surgery premises to cope with the larger number of people working in them; (b) many senior partners are reluctant to change their working methods, particularly when they can see little financial reward in such alterations; (c) nursing managers, who are responsible to the district health authorities, express a preference for community nurses, health visitors, and midwives to be based geographically rather than attached to practices; and (d) some new doctors do not wish for changes and some were fearful of mentioning such matters to their senior partners—among those who wished to make changes, many were unaware of how to bring them about. Yet it is the responsibility of the principals to provide the facilities for primary care and to bring about the changes deemed necessary.

Though nearly equal numbers of men and women doctors were recruited, the younger doctors, particularly those who had completed vocational training, tended to go into the group practices and health centres, where improvements were often not so necessary. Older men seemed to be preferred by the senior partners in the two doctor practices.

Most doctors had joined their preferred type of practice. Stockton *et al.* reported in 1980 that among doctors undergoing vocational training courses in the North East Thames Region there was a wide discrepancy between their wishes while undergoing training and the probability that their expensive training would be met. They suggested that most doctors, particularly overseas graduates, wished to work in group practices in health centres. Others have suggested that overseas graduates really wished to work in singlehanded practices and that this perhaps accounted for some of the difficulties of singlehanded practice in London.¹¹ This survey supports neither of these contentions, but it does seem clear that vocationally trained women doctors wish to go into large practices, which are willing to take them into partnership, often on a part time basis.

Small practices have the greatest difficulties in establishing primary care teams, and here the turnover of new doctors is greatest. There must be many reasons for this, including inappropriate selection by the senior partner and frustrations felt by the new junior partner. An almost inexplicable feature, however, is that in three practices where the new partner became the senior partner after a short interval and then took in a new partner the relationship rapidly deteriorated, and the partnership was dissolved.

New partners were willing to discuss the interpersonal problems leading to serious tensions that they were finding in their practices. A third of the new doctors, indeed, seemed to welcome the opportunity for discussion, and several more indicated that they had such problems but were unwilling to discuss them in detail. Some doubts are clearly raised by the training given to doctors for general practice. Some doctors thought that the training that they had received was inappropriate for the practices that they had joined, but of more concern were the complaints of over two thirds of the recently trained doctors that they had received too little training in how to manage a practice.

The Department of Health and Social Security has expressed a wish to see primary care teams developed, with increasing emphasis on preventive medical care and improved facilities for patients. This survey shows that the highly trained and motivated doctors were choosing and being chosen by the large group practices, often based in health centres, where improvements were not so necessary. The less well trained and the older doctors, who were often not so concerned about change, were entering the smaller practices, where the difficulties in bringing about the often necessary improvements are greatest. Financial inducements do not seem to be effective in bringing about such changes in the small practices. There must be many reasons for this, but the least expected ones include unsatisfactory selection procedures and poor interpersonal relationships between the partners. Where the need and the difficulties are the greatest, these human factors must produce an insuperable barrier to bringing about improvements.

Suggested remedies are: (1) Doctors who have vacancies in their practices in London might inform the secretary of the London local medical committees of their need for a partner. The committee will keep a list of doctors looking for vacancies, thus allowing for a wider choice on both sides.

(2) The training programmes for general practitioners should include an attachment to a small practice, so as to provide some experience of how they work. In addition, the importance of teaching management in general practice should be realised.

(3) New principals in general practice need some means of discussing their worries. They do not like doing so with their nearby colleagues, with whom they often work competitively, or with the local medical committees, partly for the same reason and partly because of the perceived authoritarian nature of these committees.

Nearly all the doctors interviewed stated that they were trying to give a good service to their patients, and all expressed a real concern for the quality of care that they were giving.

I thank all the doctors for spending time to be interviewed and for their friendliness and cooperation. I also thank my secretary for helping to prepare this paper. The survey was carried out between November 1983 and June 1984.

References

1. *Final Report of the problems of primary care in London*. (Lancet 1984; 1:109-12).
2. *Editorial: General practice in London—a personal view*. *Health Trends* 1984; 7:4-9.
3. *London Health Planning Commission: Primary health care in London*. LHPN, 1981.
4. *General Practice in London*. London: LHPN, 1981.
5. *General Practice in London*. London: LHPN, 1981.
6. *General Practice in London*. London: LHPN, 1981.
7. *General Practice in London*. London: LHPN, 1981.
8. *General Practice in London*. London: LHPN, 1981.
9. *General Practice in London*. London: LHPN, 1981.
10. *General Practice in London*. London: LHPN, 1981.
11. *General Practice in London*. London: LHPN, 1981.

100 YEARS AGO

Some day, when the Local Government Board can spare time from the investigation of obscure epidemics in the mountain villages of Cumberland or Wales, it may possibly be able to give a little attention to the sanitary condition of our metropolitan suburbs, especially of those comprised within the "outer ring." There is no need to use refinements of language in speaking of the public health administration of many of these suburbs. It is disgracefully and scandalously inefficient. The late Mr. Netten Radcliffe made a start upon a survey of the suburbs in the year 1871, under the direction of Mr. Simon, but, for some reason or other, the work was never more than begun. The fault of the present inefficiency does not rest altogether with the local boards. It needs an especially and unusually foresighted body to supervise and safeguard the sanitary interests of places with the mushroom-like rapidity of growth of the suburbs. There is a Bill before Parliament, giving certain powers to the Metropolitan Board of Works over the area comprised in a six miles radius from Charing Cross, instead of four miles as at present. To give the Board of Works any more authority over the local boards, it needs an especially and unusually forward-looking and sympathetic with suburban local boards; and, we think, to be at once established by the central authorities. An excellent example is West Ham, which is technically outside the metropolitan area, but which has proved an agency of suggestions to it. It goes its own way until a serious prevalence of infectious disease in its midst recalls the attention of the Local Government Board to the fact of its existence. Widespread in another suburb, whose colour white has been neglected, with consequences that are already revealing themselves. If there is to be a sanitary survey of the kingdom, let us begin in the centre and move outwards, and set the feet of wrong, overgrown, and inexperienced communities in the right path, before we begin to teach new ways to the care-hardened and obstinate sanitary defectors of past generations. (*British Medical Journal* 1883; 1:955.)

Evaluation of a computer assisted repeat prescribing programme in a general practice

M O ROLAND, L J ZANDER, MARY EVANS, RICHARD MORRIS, R A SAVAGE

Abstract

After introducing a computer assisted repeat prescribing programme into a south London practice improvements were made in several aspects of practice organisation. Time was saved by doctors and receptionists, prescriptions were produced more rapidly, information in the records about drugs that were available for repeat prescription was improved, and queries from chemists about prescriptions were reduced. The costs of the system in terms of computer operator time were estimated. Computerising prescribing data provides a source of data that may be used to enable doctors to audit their own prescribing.

Introduction

Recent developments in microtechnology have resulted in a rapid increase in the introduction of computers into general practice.

They are commonly used to process repeat prescriptions. Patients who have stable chronic conditions and require continuous medication often obtain their prescriptions without seeing the doctor. Such "repeat prescriptions" may account for nearly half of the total prescribing for adults in a practice. Information that is held on a computer about repeat prescribing may be used to audit both prescribing costs and drug management.¹ Processing repeat prescriptions should be well suited to computerisation, but initial experience suggests that the benefits have to be weighed against substantial costs—particularly the increased demands on the clerical staff.

St Thomas's Hospital Medical School, London
M O ROLAND, MRSC, MRCGP, lecturer in general practice
L J ZANDER, MRSC, senior lecturer in general practice
MARY EVANS, MRSC, lecturer in medical statistics, department of community medicine

Lambeth Road Group Practice, London SE11
MARY EVANS, research administrator

Stockwell Group Practice, London SW9
R A SAVAGE, MRSCGP, general practitioner

Correspondence to: Dr L J Zander, Department of General Practice, 10 Kennington Road, London SE11 8SP

This study was undertaken to quantify the costs and benefits of installing a computer assisted repeat prescribing programme in a general practice.

Method

In late 1982 a North East London practice was installed in the Lambeth Road Group Practice, a south London practice that provides primary medical care for roughly 9000 patients. Computer software was developed by the Sharpey Schaffer Computer Centre at St Thomas's Hospital. To identify patients who received repeat prescriptions, all requests for repeat prescriptions over a three month period were noted. The records of patients who requested these prescriptions were examined by the patients' doctors to decide which had requested prescriptions that were suitable for issuing without face to face contact between patient and doctor. These 658 patients formed the study population. Details of items of medication authorised by the doctor for repeat prescription were entered on the computer.

Then for one month several aspects of practice organisation were

measured before any prescriptions were produced by the computer. The study sample of patients was then randomly split into two groups, one of which continued to receive repeat prescriptions by the conventional system, the other receiving prescriptions by the computerised system. After a settling in period several aspects of practice organisation were measured over a three month period. The conventional group was then computerised, and measurements were made for a further month when the entire sample of patients was receiving prescriptions from the computerised system. To test all the hypotheses in the study it was necessary to combine a "before and after" study design with that of a randomised controlled trial. The study design is summarised in the figure.

In the conventional system requests for repeat prescriptions were noted by the receptionist and passed to the doctor with the patient's medical record together with a prescription on which the name and address of the patient had been entered. The doctor filled in the prescription and signed it. In the computerised system the request was passed to the computer operator without the medical records. If the requested item was authorised by the computer, a prescription was generated, which was then signed by the doctor. A dummy prescription containing items that were authorised by the computer was kept in the patient's medical record.

MEASUREMENT OF PRACTICE ORGANISATION

Use of doctor's time—During the randomised controlled trial doctors were given four batches of 10 requests for repeat prescriptions to process: two batches containing requests from the conventional group and two containing requests from the computerised group. The doctors were timed while completing these requests.

Use of receptionist's time—A research worker sat in the reception area for two hours on each of 10 consecutive weekdays mornings during both the "before" and "after" period to measure the time spent by receptionists on processing repeat prescriptions. This was broken down into the time spent receiving requests, extracting medical records, and writing prescriptions on headings.

Use of computer operator's time—A log was kept of time spent processing repeat prescriptions throughout the study period.

Prescription turnover time—The date and time of the arrival of all requests for prescriptions at the reception desk were recorded and the time when the completed prescription arrived back at the reception desk.

Queries from chemists—These were recorded throughout the study period.

Method of requesting repeat prescriptions—Throughout the study a record was kept of the way prescriptions were requested, for example by phone, in person, using repeat prescription card, or by computer slip.

Availability of prescribing details in records—During the randomised controlled trial the training general practitioners in the practice examined the medical records of all the patients in the conventional group and tried to identify those drugs that the patient was taking and which were suitable for repeat prescription. Earlier in the study the patient's own doctor had drawn up a similar list. These two lists were then compared. (It is a policy of the practice that regularly prescribed drugs should be recorded at the top of the current A4 consultation sheet.)

Results

Sixty eight patients left the practice or were taken off repeat prescriptions during the study. The results in this paper relate to the remaining sample of 590 patients.

TABLE I—Analysis of receptionist's time

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

There was also a substantial change in the way in which patients made their requests for repeat prescriptions. Before computerisation 46% of requests were made by telephone. After computerisation only 19% of requests were made by telephone, with 83% of requests being made by the patient either sending or bringing their computerised repeat prescription slip (as requested). This represented a considerable benefit to the practice, which had been trying for some time to persuade patients from asking for repeat prescriptions by telephone.

TABLE II—Analysis of doctor's time

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

The mean duration of time to complete one batch of 10 conventional prescriptions requests was 12 minutes 44 seconds compared with only one minute six seconds for 10 computerised prescriptions requests. For the conventional group the time taken to complete one batch of 10 requests was 12 minutes 44 seconds, with 83% of requests being made by the patient either sending or bringing their computerised repeat prescription slip (as requested). This represented a considerable benefit to the practice, which had been trying for some time to persuade patients from asking for repeat prescriptions by telephone.

TABLE III—Queries from chemists

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

therefore have resulted from a general improvement in processing repeat prescriptions, in addition to that resulting from the improved accuracy and legibility of computerised prescriptions.

TABLE IV—Improvement in information in medical records

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

When a patient was authorised by the doctor to receive computerised repeat prescriptions, a copy of the authorised prescription was kept in the medical records, so that it was easy to see which items were available for repeat prescription. When the notes of the "conventional" group were examined by trainers they were able to identify all drugs available on repeat prescription in 71% of the sample, but for 38% of patients only were all the drugs identified and the information found in the part of the records previously agreed by the partners (the top of the current A4 consultation sheet). Using the computer therefore resulted in an improvement in the standard of the medical records.

TABLE V—Costs of computer operator's time

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

The installation of the computer caused a substantial amount of thought, work, and frustration. Part of this resulted from the computer not being

TABLE VI—Costs of computer operator's time

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

commercially available, and so there were considerable difficulties with the software that might not be encountered by a practice buying a computer "off the shelf".

During the study period the computer operator spent a mean of five minutes 27 seconds per prescription produced with a mean of 2.2 items per prescription, and mean time per item of two minutes 15 seconds. This included printing the prescriptions, reassembling items, and keeping the repeat prescription file up to date.

Not all of the patients in the practice who were receiving repeat prescriptions were included in the study. At present the practice receives roughly 160 requests for repeat prescriptions a week. Assuming that a maximum of 90% of patients who request repeat prescriptions are likely to be on the computer at any one time, the figure in the practice is 80% at present; the repeat prescribing system would take 13 hours of computer operator time per week for this figure of 9000 patients. The system currently operates more efficiently, than during the study period—the operator takes roughly one and a half minutes per item requested, including time for housekeeping of the computer programme, equivalent to 8.6 hours of operator time a week.

WHAT PROVISION FOR REPEAT PRESCRIBING CAN THE COMPUTER DO?

Of requests made by patients who have a file on the repeat prescription programme, roughly 13% contain requests for at least one item not on the computer. Eighty per cent of patients who request repeats have a computer file, which means that the computer can process about 70% of requests in the practice.

Discussion

The results of this study show clear benefits from the introduction of a computerised repeat prescribing programme. Time was saved by receptionists and doctors, queries from chemists were substantially reduced, prescriptions were produced more rapidly, and informa-

tion in the records about drugs available for repeat prescription was improved. These improvements were probably related directly to the introduction of the computer, except turnover time for prescriptions and queries from chemists, where the improvement may in part have resulted from other changes in practice organisation that were indirectly associated with the introduction of the computer. A major additional benefit of the repeat prescribing programme is that the doctors now have a source of data on their repeat prescriptions that will enable them to audit their own repeat prescribing. The main cost, apart from that of the hardware, is the computer operator's time. These figures cannot be extrapolated with accuracy to other practices because of the wide variations that exist in practice organisation. Nevertheless, the results provide firm evidence of the tangible benefits that can be derived from a computer assisted repeat prescribing programme.

We thank the following people who helped make this study possible: Mrs Gladys Evans, who operated the computer at the Lambeth Road Group Practice; Jenni Bartholomew, research assistant; Professor David Morrell, for advice throughout the study; Chris Deves and the staff of the Sharpey Schaffer computer centre, who developed the software; Dinah Shah who helped with the statistical analysis; the doctors and staff of the Lambeth Road Group Practice and the Stockwell Group Practice, and the Inner Cities Partnership, which provided substantial funding towards the purchase of the computer.

References

1. Durrant R, Laverick A. *Medicine, practice and medicine*. London: Routledge and Kegan Paul, 1972.
2. Durrant R. *Repeating prescriptions: how computer controlled repeat prescribing*. *Br Med J* 1984; 1:1084-1086.
3. Durrant R. *Repeating prescriptions: how computer controlled repeat prescribing*. *Br Med J* 1984; 1:1084-1086.
4. Gierber R, Fitter M, Evans A, Ashdown P. *Repeating prescriptions: how computer controlled repeat prescribing*. *Br Med J* 1984; 1:1084-1086.

Accepted 16 May 1985

TABLE VII—Median prescription turnover time in hours: number of prescriptions

	Before computerisation (mean SD of 10)	During computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)	9 min (3.5)

TABLE VIII—Queries from chemists

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

The percentage of queries resulting in a query from a chemist fell over the study period from 6.0% before computerisation to 1.1% after computerisation ($\chi^2 = 7.8$, $p < 0.01$). Further analysis (table III) shows that there was a steady reduction in queries from chemists in both the computerised and the conventional group throughout the study. Some of the improvement may

therefore have resulted from a general improvement in processing repeat prescriptions, in addition to that resulting from the improved accuracy and legibility of computerised prescriptions.

TABLE IX—Improvement in information in medical records

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

therefore have resulted from a general improvement in processing repeat prescriptions, in addition to that resulting from the improved accuracy and legibility of computerised prescriptions.

TABLE X—Improvement in information in medical records

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

therefore have resulted from a general improvement in processing repeat prescriptions, in addition to that resulting from the improved accuracy and legibility of computerised prescriptions.

TABLE XI—Costs of computer operator's time

	Before computerisation (mean SD of 10)	After computerisation (mean SD of 10)
No. of prescriptions requested	9.4 (4.1)	9.2 (3.0)
No. of prescriptions issued	11.6 (5.2)	11.1 (4.1)
Time taken to issue prescriptions	22 min (5.5)	17 min (5.5)
Receiving requests	9 min (2.5)	8 min (2.5)
Writing prescriptions	9 min (3.5)	9 min (3.5)
Looking for records	9 min (3.5)	9 min (3.5)

The installation of the computer caused a substantial amount of thought, work, and frustration. Part of this resulted from the computer not being

commercially available, and so there were considerable difficulties with the software that might not be encountered by a practice buying a computer "off the shelf".

During the study period the computer operator spent a mean of five minutes 27 seconds per prescription produced with a mean of 2.2 items per prescription, and mean time per item of two minutes 15 seconds. This included printing the prescriptions, reassembling items, and keeping the repeat prescription file up to date.

Not all of the patients in the practice who were receiving repeat prescriptions were included in the study. At present the practice receives roughly 160 requests for repeat prescriptions a week. Assuming that a maximum of 90% of patients who request repeat prescriptions are likely to be on the computer at any one time, the figure in the practice is 80% at present; the repeat prescribing system would take 13 hours of computer operator time per week for this figure of 9000 patients. The system currently operates more efficiently, than during the study period—the operator takes roughly one and a half minutes per item requested, including time for housekeeping of the computer programme, equivalent to 8.6 hours of operator time a week.

WHAT PROVISION FOR REPEAT PRESCRIBING CAN THE COMPUTER DO?

Of requests made by patients who have a file on the repeat prescription programme, roughly 13% contain requests for at least one item not on the computer. Eighty per cent of patients who request repeats have a computer file, which means that the computer can process about 70% of requests in the practice.

Discussion

The results of this study show clear benefits from the introduction of a computerised repeat prescribing programme. Time was saved by receptionists and doctors, queries from chemists were substantially reduced, prescriptions were produced more rapidly, and informa-

tion in the records about drugs available for repeat prescription was improved. These improvements were probably related directly to the introduction of the computer, except turnover time for prescriptions and queries from chemists, where the improvement may in part have resulted from other changes in practice organisation that were indirectly associated with the introduction of the computer. A major additional benefit of the repeat prescribing programme is that the doctors now have a source of data on their repeat prescriptions that will enable them to audit their own repeat prescribing. The main cost, apart from that of the hardware, is the computer operator's time. These figures cannot be extrapolated with accuracy to other practices because of the wide variations that exist in practice organisation. Nevertheless, the results provide firm evidence of the tangible benefits that can be derived from a computer assisted repeat prescribing programme.

We thank the following people who helped make this study possible: Mrs Gladys Evans, who operated the computer at the Lambeth Road Group Practice; Jenni Bartholomew, research assistant; Professor David Morrell, for advice throughout the study; Chris Deves and the staff of the Sharpey Schaffer computer centre, who developed the software; Dinah Shah who helped with the statistical analysis; the doctors and staff of the Lambeth Road Group Practice and the Stockwell Group Practice, and the Inner Cities