References

- 1 Ciba Guest Symposium. Terminology, definitions and classifications of chronic pulmonary emphysema and related conditions. *Thorax* 1959;14:286-99.
- 2 Reid L. Measurement of the bronchial mucous gland layer: a diagnostic yardstick in chronic bronchitis. *Thorax* 1960;15:132-41.
- 3 Thurlbeck WM, Angus GE. A distribution curve for chronic bronchitis. *Thorax* 1964;19:436-42.
- 4 Hayes JA. Distribution of bronchial gland measurements in a Jamaican population. *Thorax* 1969;24:619-22.
- 5 McKenzie HI, Glick M, Outhred KG. Chronic bronchitis in coal miners: ante-mortem/post-mortem comparisons. *Thorax* 1969;24:527-35.
- 6 Scott KWM. An autopsy study of bronchial mucous gland hypertrophy in Glasgow. *Am Rev Respir Dis* 1973;107:239-45.
- 7 Fletcher CM, Pride NB. Definitions of emphysema, chronic bronchitis, asthma, and airflow obstruction: 25 years on from the Ciba symposium. *Thorax* 1984;39:81-5.
- 8 Reid L. Pathology of chronic bronchitis. *Lancet* 1954;i:275-8.
- 9 MacLean KH. The histology of generalized pulmonary emphysema. I. The genesis of the early centrolobular lesion: focal emphysema. *Australasian Annals of Medicine* 1957;6:124-40.
- 10 Sharp JT, Paul O, McKean H, Best WR. A longitudinal study of bronchitic symptoms and spirometry in a middle-aged, male, industrial population. *Am Rev Respir Dis* 1973;108:1066-77.
- 11 Bates DV. The fate of the chronic bronchitic. *Am Rev Respir Dis* 1973;108:1043-65.
- 12 Fletcher C, Peto R. The natural history of chronic airflow obstruction. *Br Med J* 1977;ii:1645-8.
- 13 Peto R, Speizer FE, Cochrane AL, et al. The relevance in adults of airflow obstruction, but not of mucus hypersecretion, to mortality from chronic lung disease. *Am Rev Respir Dis* 1983;128:491-500.
- 14 Hogg JC, Macklem PT, Thurlbeck W. Site and nature of airway obstruction in chronic obstructive lung disease. *N Engl J Med* 1968;278:1355-60.
- 15 Cosio M, Ghezzi H, Hogg JC, et al. The relations between structural changes in small airways and pulmonary-function tests. *N Engl J Med* 1978;298:1277-81.
- 16 Berend N, Woolcock AJ, Marlin GE. Correlation between the function and structure of the lung in smokers. *Am Rev Respir Dis* 1979;119:695-705.
- 17 Cosio MG, Hale KA, Niewoehner DE. Morphologic and morphometric effects of prolonged cigarette smoking on the small airways. *Am Rev Respir Dis* 1980;122:265-71.
- 18 Petty TL, Silvers W, Stanford RE, Baird D, Mitchel RS. Small airways pathology is related to increased closing capacity and abnormal slope of phase III in excised human lungs. *Am Rev Respir Dis* 1980;121:449-56.
- 19 Berend N, Wright JL, Thurlbeck WM, Marlin GE, Woolcock AJ. Small airways disease: reproducibility of measurements and correlation with lung function. *Chest* 1981;79:263-8.
- 20 Wright JL, Lawson LM, Pare PD, Kennedy S, Wiggs B, Hogg JC. The detection of small airways disease. *Am Rev Respir Dis* 1984;129:989-94.
- 21 Pare PD, Harvey K, Mildenberger M, Brooks LA. The effects of balloon volume and position on the pressure volume curve of the lung. *Clin Invest Med* 1983;6:143-6.
- 22 Dubois AB, Botelho SY, Bedel GW, Marshall R, Comroe JH. A rapid plethysmographic method for measuring thoracic gas volume: a comparison with a nitrogen washout method for measuring functional residual capacity in normal subjects. *J Clin Invest* 1956;35:322-6.
- 23 Buist AS, Ross BB. Quantitative analysis of the alveolar plateau in the diagnosis of early airway obstruction. *Am Rev Respir Dis* 1973;108:1078-87.
- 24 Bates DV, Woolf CR, Paul GI. Chronic bronchitis: a report on the first two stages of the coordinated study of chronic bronchitis in the Department of Veterans Affairs, Canada. *Medical Services Journal, Canada* 1962;18:211-303.
- 25 Gaensler EA, Wright GW. Evaluation of respiratory impairment. *Arch Environ Health* 1966;12:146-89.
- 26 Morris JF, Koski A, Johnson LC. Spirometric standards for healthy nonsmoking adults. *Am Rev Respir Dis* 1971;103:57-67.
- 27 Dosman J, Bode F, Urbanetti J, Martin R, Macklem PT. The use of a helium-oxygen mixture during expiratory flow to demonstrate obstruction in small airways in smokers. *J Clin Invest* 1975;55:1090-9.
- 28 Thurlbeck WM, Dunnill MS, Hartung W, Heard BE, Heppleston AG, Ryder RC. A comparison of three methods of measuring emphysema. *Hum Pathol* 1972;1:215-26.
- 29 Hollander M, Wolfe DA. *Nonparametric statistical methods*. New York: John Wiley and Sons, 1973:71-82.
- 29a Hollander M, Wolfe DA. *Nonparametric statistical methods*. New York: John Wiley and Sons, 1973:185-99.
- 30 Hale FC, Olsen CR, Mickey MR. The measurement of bronchial wall components. *Am Rev Respir Dis* 1968;98:978-87.
- 31 Dunnill MS, Massarella GR, Anderson JA. A comparison of the quantitative anatomy of the bronchi in normal subjects, in status asthmaticus, in chronic bronchitis, and in emphysema. *Thorax* 1969;24:176-9.
- 32 Takizawa T, Thurlbeck WM. Muscle and mucous gland size in the major bronchi of patients with chronic bronchitis, asthma and asthmatic bronchitis. *Am Rev Respir Dis* 1971;104:331-6.
- 33 Niewoehner DE, Kleinerman J, Knoke JD. Regional chronic bronchitis. *Am Rev Respir Dis* 1972;105:586-93.
- 34 Oberholzer M, Dalquen P, Rohr HP. Stereology, a complement to respiration research. Bronchus morphometry: methodology and baseline data. *Microsc Acta* 1977;79:205-23.
- 35 Florey HW. The secretion of mucus and inflammation of mucous membranes. In: Florey HW, ed. *General pathology*. 3rd ed. London: Lloyd-Luke Ltd, 1962:167-96.
- 36 Takizawa T, Thurlbeck WM. A comparative study of four methods of assessing the morphologic changes in chronic bronchitis. *Am Rev Respir Dis* 1971;103:774-83.
- 37 Carlile A, Edwards C. Structural variation in the named bronchi of the left lung. A morphometric study. *Br J Dis Chest* 1983;77:344-8.
- 38 Wright JL, Wiggs BJ, Hogg JC. Airway disease in upper and lower lobes in lungs of patients with and without emphysema. *Thorax* 1984;39:282-5.
- 39 Thurlbeck WM, Angus GE, Pare JAP. Mucous gland hypertrophy in chronic bronchitis, and its occurrence in smokers. *Br J Dis Chest* 1963;57:73-8.
- 40 Restrepo G, Heard BE. The size of the bronchial glands in chronic bronchitis. *Journal of Pathology and Bacteriology* 1963;85:305-10.
- 41 Ryder RC, Dunnill MS, Anderson JA. A quantitative study of the bronchial mucous gland volume, emphysema and smoking in a necropsy population. *J Pathol* 1971;104:59-71.
- 42 Field WEH, Davey EN, Reid L, Roe FJC. Bronchial mucus gland hypertrophy: its relation to symptoms and environment. *Br J Dis Chest* 1966;60:66-80.
- 43 Salvato G. Some histological changes in chronic bronchitis and asthma. *Thorax* 1968;23:168-72.
- 44 Martin C, Katsura S, Cochran TH. The relationship of chronic bronchitis to the diffuse obstructive pulmonary syndrome. *Am Rev Respir Dis* 1970;102:362-9.
- 45 Hossain S, Heard BE. Hyperplasia of bronchial muscle in chronic bronchitis. *J Pathol* 1970;101:171-84.
- 46 Restrepo GL, Heard BE. Air trapping in chronic bronchitis and emphysema: measurement of the bronchial cartilage. *Am Rev Respir Dis* 1964;90:395-400.
- 47 Greenberg SD, Boushy SF, Jenkins DE. Chronic bronchitis and emphysema: correlation of pathologic findings. *Am Rev Respir Dis* 1967;96:918-28.
- 48 Tandon MK, Campbell AH. Bronchial cartilage in chronic bronchitis. *Thorax* 1969;24:607-12.
- 49 Thurlbeck WM, Pun R, Toth J, Frazer RG. Bronchial cartilage in chronic obstructive lung disease. *Am Rev Respir Dis* 1974;109:73-80.
- 50 Tobin MJ, Jenouri G, Sackner MA. Subjective and objective measurement of cigarette smoke inhalation. *Chest* 1982;82:696-700.
- 51 Valberg PA, Brain JD, Sneddon SL, LeMott SR. Breathing patterns influence aerosol deposition sites in excised dog lungs. *J Appl Physiol* 1982;53:824-37.
- 52 Higenbottom T, Feyerabend C, Clark TJH. Cigarette smoke inhalation and the acute airway response. *Thorax* 1980;35:246-54.
- 53 Falk HL, Tremmer HM, Kotin P. Effect of cigarette smoke and its constituents on ciliated mucus-secreting epithelium. *Journal of the National Cancer Institute* 1959;23:999-1012.
- 54 Ailsby RL, Ghadially FN. A typical cilia in human bronchial mucosa. *J Pathol* 1973;109:75-8.
- 55 McDowell EM, Barrett LA, Harris CC, Trump BF. Abnormal cilia in human bronchial epithelium. *Arch Pathol Lab Med* 1976;100:429-326.
- 56 Rossman CM, Lee RMKW, Forrest JB, Newhouse MT. Nasal ciliary ultrastructure and function in patients with primary ciliary dyskinesia compared with that in normal subjects and in subjects with various respiratory diseases. *Am Rev Respir Dis* 1984;129:161-7.
- 57 Camner P, Philipson K. Tracheobronchial clearance in smoking—discordant twins. *Arch Environ Health* 1972;25:60-3.
- 58 Sanchis J, Dolovich M, Rossman C, Newhouse M. Lung clearance in patients with airways obstruction. *Bulletin de Physio-Pathologie Respiratoire (Nancy)* 1973;9:325-35.
- 59 Katz DH, Mead R. Control of antibody synthesis. In: Middleton E, Reed C, Ellis EF, eds. *Allergy: principles and practice*. 2nd ed. St Louis: C V Mosby, 1983:31-42.

(Accepted 10 July 1985)

SHORT REPORTS

Toxocariasis and eosinophilic meningitis

Toxocariasis is usually manifested as visceral larva migrans in young children with pica.¹ Lesions in the eye are more common in older children and adults,^{1,2} and rarely other parts of the central nervous system may be affected.² A benign, self limiting meningitis has not been described and we therefore report such a case.

Case history

A previously healthy 11 year old girl was referred by her general practitioner with a 10 day history of worsening neck stiffness and back pain. She had initially presented with severe bilateral frontal headache and dizziness but these symptoms had resolved within 48 hours after beginning treatment with paracetamol.

On examination the child looked alert and well and was not feverish. There was pronounced meningism and she complained of severe pain at the level of T12-L1, accentuated by movement. Kernig's sign was positive and both optic discs were blurred. There was generalised hyperreflexia. The rest of the physical examination, routine blood tests, and radiography showed nothing unusual. An urgent CT scan showed no abnormality. Lumbar punc-

ture the day after admission yielded 150 white cells, reported initially as being predominantly lymphocytes with normal protein and glucose. In view of the atypical history Romanovsky staining was performed on a cytospin of cerebrospinal fluid. This showed that 30% of the cells were eosinophils and that the remainder were lymphocytes. Culture and serology for common viral and bacterial infection were unhelpful. Two days after admission an eosinophilia of $1.3 \times 10^9/l$ was noted, which persisted to follow up at six weeks.

The patient recovered spontaneously over the next three days and was sent home. Results of serological screening for toxocaral infection suggested active infection (see table), both acute and convalescent serum samples having appreciable titres. Anti-A agglutinin titres were also raised but fell during convalescence, were IgG only, and the patient had never had a blood transfusion, all points suggestive of acute toxocaral infection. She denied pica but both neighbours had recently acquired puppies, ensuring a plentiful supply of toxocara ova.

She was treated with diethylcarbamazine 8 mg/kg for 10 days and tolerated the treatment well. She continued to be asymptomatic nine months after presentation.

Comment

This case was unusual in its presentation and is instructive in the method of diagnosis. Pronounced eosinophilia in the cerebrospinal fluid is rare in Britain but there are many recognised causes.³ Other

reports have emphasised the severity of neurological and systemic disorder in patients with toxocaral meningoencephalitis.¹⁻⁴ Apart from neck stiffness our patient was surprisingly well with no evidence of encephalitis. Her diagnosis might have been that of an atypical viral meningitis had Romanovsky staining not been performed. We think that this investigation may be useful in culture negative meningitis when there are no signs of viral infection.

Toxocara enzyme linked immunosorbent assay titres using T canis excretory-secretory products as antigen

	Acute serum	Convalescent serum
IgG	1/400	1/200
IgE	1/320	1/320
Anti-A agglutinins:		
IgM { A ₁	1/16	1/16
A ₂	1/8	1/8
IgG { A ₁	1/8192	1/4096
A ₂	1/8192	1/2048

Note: erythrocytes 0+.

We decided to give specific antitoxocaral treatment because of the risk of ocular lesions or recurrence in the central nervous system,¹ although there is little evidence for the efficacy of diethylcarbamazine or thiabendazole¹⁻² and there is the risk of release of larval antigen and reaction to this.

This is the first reported case of a mild, self limiting eosinophilic meningitis due to *Toxocara canis*, but overt manifestations of toxocaral infection are uncommon in Britain. It is interesting to speculate how common this presentation of toxocariasis may be in view of the fact that over 2% of the population in Britain have seroconverted.⁵

Thanks are due to Drs R W A Girdwood and H V Smith, of the Scottish Parasite Reference Laboratory, for helpful advice and for performing the assays and to Mrs D Hammond for typing the manuscript.

- Schantz PM, Glickman LT. Toxocaral visceral larva migrans. *N Engl J Med* 1978;**298**:436-9.
- Bell WE, McCormick WF. *Neurologic infections in childhood*. 2nd ed. Philadelphia: W B Saunders Company, 1981:572-6.
- Bosch I, Oehmichen M. Eosinophilic granulocytes in cerebrospinal fluid: analysis of 94 cerebrospinal fluid specimens and review of the literature. *J Neurol* 1978;**219**:93-105.
- Schochet SS. Human *Toxocara canis* encephalopathy in a case of visceral larva migrans. *Neurology* 1967;**17**:227-9.
- Woodruff AW. Toxocariasis. *Br Med J* 1970;**iii**:663-9.

(Accepted 19 July 1985)

Departments of Microbiology and Paediatric Neurology, The Children's Hospital, Birmingham

I M GOULD, MRCP, senior registrar in microbiology
S NEWELL, MRCP, registrar
S H GREEN, MRCP, consultant paediatrician
R H GEORGE, MRCPATH, consultant medical microbiologist

Correspondence to: Dr I M Gould, Department of Medical Microbiology, Dudley Road Hospital, Birmingham B18 7QH.

Size of pancreas in diabetes mellitus: a study based on ultrasound

We have shown that exocrine pancreatic deficit is common in diabetes and that the degree of this deficit parallels the endocrine deficit.¹⁻⁴ From necropsy data we also know that the size of the pancreas in patients with insulin dependent diabetes is smaller than in normal people.⁵ In order to confirm this observation in insulin dependent diabetes⁵ and to determine whether pancreatic size is also altered in non-insulin dependent diabetes mellitus we studied the pancreatic size in patients with diabetes mellitus using ultrasound techniques.

Patients, methods, and results

Sixty adult diabetic patients were investigated: 22 had insulin dependent diabetes (group 1) and 19 non-insulin dependent diabetes (group 2) and 19 were non-ketotic patients who had to be given insulin because of inadequacy of diabetic control with oral hypoglycaemic agents (group 3). Nineteen healthy controls were also studied.

All scans were performed by one of us (LAB), using a real time linear array system (Picker LS 3000).

The patients and controls were scanned in the morning after an overnight fast. Scans were performed with the patients supine and erect. A more complete and clearer visualisation of the pancreas was achieved by scanning with the patient erect. The head (defined as the area medial to the superior mesenteric vein) and the body of the pancreas were measured separately, since these were often visualised to best advantage in different views. As the head is often oriented in the longitudinal plane parallel to the inferior vena cava, measurements were made in this plane below the portal vein as well as in the transverse or oblique plane (taking the midpoint of the confluence of the superior mesenteric and splenic veins as the marker point). In normal subjects the longitudinal section of the head was frequently larger than the transverse sections. This did not occur in any of the diabetic patients. The tail of the pancreas was well seen as it passed anterior to the left kidney, but the more distal portion extending into the splenic hilum (which represents a very small part of the pancreatic mass) was rarely seen.

The scans were recorded on photographic paper. The outline of the pancreas was mapped out and the areas computed using a Numonics Graphic Analyzer (Numonics, Japan).

Both the head and the body of the pancreas in patients of group 1 and 2 were noticeably smaller than in the controls (table). The head and body of the pancreas in patients of group 2 were significantly larger than in group 1.

Total area of pancreas and area of head and body of pancreas (in cm²) in diabetic patients and controls. Values are medians (ranges in parentheses)

Group	No of subjects	Total area	Area of head	Area of body
1	22	8.9*† (4.6-14.7)	3.2*† (1.1-5.7)	5.7*‡ (2.0-11.2)
2	19	12.9‡ (6.6-17.8)	5.7‡ (2.2-11.8)	7.8‡ (4.4-11.8)
3	19	11.9‡ (6.7-17.3)	4.8‡ (2.5-6.4)	6.7‡ (3.4-13.6)
Controls	19	18.7 (13.0-26.1)	8.8 (6.2-11.1)	10.4 (7.1-19.8)

*Compared with controls $p < 0.001$.

†Compared with group 2 $p < 0.002$.

‡Compared with controls $p < 0.002$.

§Compared with group 2 $p < 0.05$.

In group 3 the sizes of the head and body of the pancreas were intermediate between those in groups 1 and 2.

There was no correlation between size of the pancreas and body weight or duration of diabetes.

Comment

These results show that the pancreas is significantly smaller in diabetic patients than in healthy controls. Furthermore, patients with insulin dependent diabetes have significantly smaller pancreases than patients with non-insulin dependent disease. Non-ketotic patients whose diabetes was not controlled with maximum doses of oral hypoglycaemic agents and who required insulin had pancreases intermediate in size between those in the other two groups. This is the first study to document these changes systematically. The pattern of diminution in the size of the pancreas in diabetes parallels the impairment of exocrine function previously described by us. Thus patients with insulin dependent diabetes have the lowest serum concentrations of pancreatic enzymes. Patients with non-insulin dependent disease have marginally reduced serum pancreatic enzyme values, while patients with non-ketotic disease who require insulin because of inadequate control with oral hypoglycaemic agents have serum pancreatic enzyme values between those of the other two groups.¹⁻⁴

Pancreatic enzyme values at the time of clinical onset of insulin dependent diabetes mellitus indicate that in these patients exocrine pancreatic reserve may be diminished at the time of presentation, suggesting that exocrine damage occurs not as a long term complication of and sequel to diabetes but probably as a consequence of the process which caused the pancreatic islet and β cell damage.⁴ The factors responsible for the shrinkage of pancreas through a reduction of exocrine tissue may be immunological (lymphocytic infiltration has been observed); genetic; subclinical viral pancreatitis; or lack of trophic effect of insulin on exocrine pancreas. The last is particularly relevant to non-insulin dependent diabetes.

We thank Pamela Dale for secretarial work and Georgina Gray and Rita Stavrou for their help.

- Dandona P, Elias E, Beckett AG. Serum trypsin concentrations in diabetes mellitus. *Br Med J* 1978;**ii**:1125-7.
- Foo Y, Rosalki SB, Ramdial L, Mikhailidis D, Dandona P. Serum isoamylase concentrations in diabetes mellitus. *J Clin Pathol* 1980;**33**:1102-5.
- Junglee D, de Albarran R, Katrak A, Freedman D, Beckett AG, Dandona P. Pancreatic lipase concentration in diabetes mellitus. *J Clin Pathol* 1983;**36**:200-2.
- Dandona P, Freedman DB, Foo Y, et al. Exocrine pancreatic function in diabetes mellitus. *J Clin Pathol* 1984;**37**:302-6.