PRACTICE OBSERVED

Practice Research

Acute otitis media: clinical course among children who received a short course of high dose antibiotic

JOHN BAIN, ELIZABETH MURPHY, FRANCES ROSS

Abstract

A prospective study was carried out in 274 children aged 3 to 10 years with acute otitis media. They were randomly allocated to one of two treatment regimens: (a) a seven day course of amoxycillin 125 mg three times a day, and (b) a two day course of amoxycillin 750 mg twice a day. They were followed up by symptom diaries and clinical examination.

The findings in the 243 children who completed the trial showed that the short course of treatment was as effective as the seven day course in the speed of resolution of symptoms and signs, irrespective of previous history of otitis media or of episodes in which bulging of the eardrums was observed at presentation.

A subgroup of 185 children was followed up for one year after entry to the trial. During this period no appreciable differences emerged between the two antibiotic regimens, either in recurrence rate of otitis media or in the frequency of hearing loss at one month and six months after entry to the study. Side effects of treatment were few, and those that could be attributed to antibiotic use occurred with equal frequency in the two treatment groups.

Introduction

Acute otitis media is usually defined mainly in relation to the physical appearance of the eardrum,¹ but clinical experience indicates that parents and doctors respond to symptoms as much as to signs, epilepsy, and diabetes.

Department of Primary Medical Care, University of Southampton

JOHN BAIN, *MD, FRCS, professor*

ELIZABETH MURPHY, *MA, research assistant*

FRANCES ROSS, *research assistant*

Correspondence to Professor John Bain, Aldermoor Health Centre, Southampton SO1 6ST.

signs in managing children with presumed middle ear infection. It is common practice to prescribe an antibiotic for acute otitis media on the assumption that the pathogenesis includes bacterial infection in most cases. This assumption is based on the findings that bacteria can be isolated in aspirations from the middle ear of children with acute otitis media.²⁻⁴ Because *Haemophilus influenzae* has been isolated in up to a quarter of cases, especially in children under the age of 5, a broad spectrum antibiotic such as ampicillin or amoxycillin is usually recommended.⁵ It is widely believed that such antibiotic treatment will not only shorten an episode and prevent complications, such as mastoiditis, but may reduce the likelihood of recurrent episodes.

A leading article in the *BMJ* in 1979 suggested that traditional regimens for common conditions should be scrutinized.⁶ In the case of otitis media antibiotics are usually prescribed for between five and seven days,⁷ and most suspensions of antibiotics are produced in packages sufficient for either a five or a 10 day course. There is little certainty concerning the optimal duration of treatment, but the findings of two recent small studies suggest that two to three days of either penicillin or amoxycillin at conventional doses may be as effective as the same dose given over 10 days.^{8,9}

The aims of our study were (a) to describe the syndrome of acute otitis media in terms of symptoms and signs and (b) to compare the clinical outcome in children who presented with acute otitis media aged 3 to 10, who received either a two day course of amoxycillin 750 mg twice daily or a seven day course of amoxycillin 125 mg three times a day. A subgroup of the children who were studied was followed up for a year to compare the rates of recurrence and hearing loss in the two treatment regimens.

Methods

The study was conducted in the winter months of 1982-3 and 1983-4 in four practices. The 16 participating general practitioners were asked to recruit 10 consecutive children with acute otitis media each winter, with a target of 20 cases per doctor. Entry to the trial was confined to (a) children

aged 3 to 10 years, (b) those with symptoms and signs of acute otitis media, which in the general practitioner's opinion warranted treatment with an antibiotic, and (c) children whose parents gave their informed consent. Children were excluded who (a) were under age 3 because they present difficulties in communication, and different drug dosages are often recommended for younger children; (b) had had an episode of acute otitis media that required medical attention in the previous month; (c) were currently on antibiotics or had received an antibiotic for any reason in the previous 14 days; (d) were known or suspected of being hypersensitive to antibiotics; and (e) were on other drugs (short or long term) for other conditions such as asthma, epilepsy, and diabetes.

Children were entered into the trial on one occasion only. At entry to the trial doctors completed a medical and social history and recorded information about presenting symptoms and eardrum signs as described in a previous study of the treatment of acute otitis media.¹⁰ The record card was retained in the patient's notes and arrangements made for the child to attend the same general practitioner at one week later when a follow up record of eardrum signs was completed. The follow up of symptoms was measured by using a symptom diary which was issued at the initial consultation and completed by mothers during a 20 day follow up period. Children were then randomly allocated to long (seven days, group A) or short (two days, group B) treatments with amoxycillin using a computerized allocation schedule, which was placed with bottles containing the two treatments labelled A or B in each general practitioner's consulting room. Amoxycillin powder was reconstituted by the general practitioner by adding the appropriate amount of water to bottle A and bottle B. Parents were advised to use analgesics as they considered necessary and to record the use of drugs for pain relief in the symptom diaries.

Within 24 hours of entry to the trial a research assistant visited the child's parents at home to answer queries and ensure that the method of completing the symptom diaries was understood. The research assistant visited each family again at roughly seven to 10 days and at three weeks after entry to the trial. Children who attended our health centre were followed up for one year to identify recurrence of acute otitis media and hearing loss. To ascertain hearing loss audiometry was conducted by a research assistant in those children aged 4 and over at one month and six months after entry to the trial. The assessment of hearing was performed in a soundproof room, and clinically important hearing loss was defined as 30 dB loss at two or more frequencies.

Data were analysed using standard χ^2 statistical tests.

Results

Two hundred and seventy four children were admitted to the trial and 243 completed follow up for three weeks, with 118 in group A (seven days) and 125 in group B (two days). Thirty one (11.3%) children did not complete follow up for three weeks, with 15 in group A and 16 in group B. The reasons for failure to complete follow up were (a) 12 children who moved home, (b) 12 children who were lost to follow up, and (c) 7 children who were unable to attend for follow up. The children who completed follow up for three weeks were followed up for one year to identify recurrence of acute otitis media and hearing loss. To ascertain hearing loss audiometry was conducted by a research assistant in those children aged 4 and over at one month and six months after entry to the trial. The assessment of hearing was performed in a soundproof room, and clinically important hearing loss was defined as 30 dB loss at two or more frequencies.

Table 1—Resolution of symptoms at day 1, day 5, and day 10 after entry into trial. Group A, seven days' treatment (n=118 children); group B, two days' treatment (n=125 children).

Symptoms	Day 1		Day 5		Day 10	
	A (n=118)	B (n=125)	A (n=118)	B (n=125)	A (n=118)	B (n=125)
Earsache	89.5	94.4	9.1	10.4	5.1	5.6
Earsore at night	64.4	67.2	3.4	2.4	2.2	2.4
Earwax	46.6	38.9	13.9	31.2	29.7	29.2
Head discharge	37.6	70.4	28.8	24.6	16.9	19.2
Cough	46.6	56.0	7.6	9.4	4.4	4.0

Table 1—Symptoms and diary follow up: five main symptoms and outcomes on which symptoms were recorded during 20 days after entry to trial.

Symptoms	Group A (seven days' treatment)		Group B (two days' treatment)	
	Day 1	Day 5	Day 1	Day 5
Earsache	2.6	2.6	2.6	2.6
Earsore at night	1.7	0.9	1.7	0.9
Head discharge	4.6	6.0	4.6	6.0
Cough	4.0	1.2	4.0	1.2

Bulging of the drum was selected as a prominent sign, possibly indicating more objectively the presence of infection in the middle ear cleft, and 140 children had this sign with or without other abnormalities. These children were equally distributed between the two treatment groups, and the speed of resolution of their symptoms and the improvement of eardrum appearances at one week were similar for this subgroup in both treatment groups.

PREVIOUS MEDICAL HISTORY OF OTITIS MEDIA AND OUTCOME

A documented history of otitis media was found in 188 children, and these were equally distributed between the two treatment groups. The time taken for symptoms to resolve and the persistence of abnormal eardrum signs were marginally greater in children who had a previous medical history of middle ear infection than in other children in the trial, but this difference was not significant. There were no notable differences in clinical outcome between the two day and seven day courses of treatment in the children with a previous history of otitis media.

RECURRENT OTITIS MEDIA

In the health centre where children were followed up for one year 185 children completed the trial and 76 (41%) had at least one further episode of acute otitis media that required medical attention during the 12 months after entry to the trial. Thirty seven (49%) of these 76 children with a record of

recurrence had the first episode of recurrence within three months of the index episode. There was a significantly higher ($\chi^2=4.04$, p<0.05) number of children with recurrence under age 4, 34 (45%) of the 76 with recurrence being under 4. This compares with 33 (30%) of the 109 children with no recurrence being under 4. There was a trend for children with recurrent episodes to have had a previous history of otitis media, but the trend was not significant.

Table IV shows that irrespective of age there was no appreciable difference in the number of children who had a recurrence in the two treatment groups. One hundred and ten of the 185 children had bulging of at least one eardrum. There was no association between the duration of

TABLE IV—Recurrence of acute otitis media in 12 month period after entry to trial. In group A, seven days' treatment; and group B, two days' treatment.

Group	Total No. (n)	Mean age (years)	No. with recurrence	
			Group A	Group B
A	86	4.8	34 (39.5)	4.2
B	99	5.0	42 (42.4)	4.1
Total	185	4.9	76 (41.1)	4.1

treatment for the index episode and subsequent recurrence in this subgroup. Among the 110 children with initial bulging of the eardrum 52 (47%) had a record of recurrences compared with 24 (32%) of the 75 children with less prominent eardrum signs, and this relation between bulging of the drum and recurrence was significant ($\chi^2=4.296$, p<0.05).

AUDIOMETRY

In the health centre where follow up audiometry was performed 106 children aged 4 and over completed the trial. Of these, 73 had audiometry performed at both one month and six months after entry to the trial. Among these children there were no appreciable differences in hearing loss between the two treatment groups, and only five children (two in group A and three in group B) had a hearing loss of 30 dB or more at both one month and six months after entry to the trial.

Discussion

A course of high dose antibiotic treatment of short duration was as effective as the conventional one week course in resolving symptoms and signs in acute otitis media. The type of treatment received had no appreciable effect on recurrence of otitis media during the one year of follow up and made no difference to hearing loss recorded at one month and six months after entry to the trial.

Compliance with antibiotic treatment for seven to 10 days in children is poor, with up to half failing to complete the full course.¹¹ In a review of compliance with treatment Stewart and Cliff showed that the longer the treatment and the greater the frequency of dosage schedules the less likely a patient is to complete the recommended course of treatment.¹² The results of previous trials with smaller numbers of children (84 and 103 respectively)^{8,9} showed no convincing differences between two and three day treatments and conventional dosages of antibiotics for seven and 10 days. There is even less evidence on the optimum dose beyond the suggestion that more treatment failures occur in patients who are treated with low doses for seven days than with higher doses for 10 days.¹³

When children receive 10 to 12 doses over 48 to 72 hours there is still a possibility of doses being missed and inadequate concentrations of antibiotics being achieved. If mothers are advised to give amoxycillin 750 mg 12 hourly for four days the regimen is simplified even further with the aim of adequate therapeutic concentrations being sustained over the course of treatment.

Ampicillin and amoxycillin are effective against most bacteria that are implicated in acute otitis media.¹⁴ Amoxycillin was chosen as the antibiotic for the study because it has the advantages of better absorption and a lower incidence of gastrointestinal side effects.¹⁵ Of the 12 children who failed to complete the trial, 12 had presumed side effects, but seven only were related to gastrointestinal upset. There were similar numbers in the two treatment groups. Leigh *et al*

found that 17.5% of adult patients who received single high dose treatment with amoxycillin had diarrhoea,¹⁶ but that in studies of high dose treatment in adults gastrointestinal upset were not a problem.¹⁷ The evidence from this trial suggests that a high dose, short course of treatment with amoxycillin does not lead to noticeable gastrointestinal disturbances in children aged 3 and over.

The clinical diagnosis of middle ear infection is often limited to a definition of abnormalities observed on the tympanic membrane. The general practitioner often sees a child during the early stages of the disease, and a precise diagnosis is difficult to make. We have to accept that children were included in this trial who may not have had pus formation in the middle ear cleft. The results, however, emphasise that in reality clinical presentation in general practice encompasses clusters of symptoms and signs, and given that myringotomy is not conducted in general practice decisions about treatment have to be made from a complex of symptoms and signs.

Studies of the treatment of otitis media often omit any reference to symptoms, and the duration of follow up is often short. Though the main aim of this study was to compare different treatment regimens, the results provide valuable information about the clinical course of otitis media in general practice. Earache tended to resolve rapidly, but the signs of associated upper respiratory tract infection were more persistent, with roughly one in five children continuing to have symptoms such as cough and nasal discharge 10 days after entry to the trial. Children under 4 were more likely to have a recurrence, which is consistent with the experience of clinical practice where the preschool child presents more frequently with upper respiratory and middle ear infections.

There is no overall agreement about which eardrum appearances are diagnostic of bacterial infection, but the importance of bulging of the tympanic membrane is cited as one of the most useful indicators of suppurative otitis media.¹⁸⁻²⁰ The presence of a bulging drum was a predictor of a recurrence of otitis media. Thus younger children and those with bulging of the drum should probably be regarded as an important subgroup who require more careful follow up.

Four out of 10 children had at least one further episode of acute otitis media requiring an antibiotic during the one year of follow up, and half of these children had their initial recurrence within three months of the index episode—a result similar to the findings of Onon and Taylor.²¹ With any given episode of middle ear infection the chance of recurrence in that child within one year is between 30% and 50%. After an acute attack the resolution of middle ear effusion associated with infection may take six weeks or longer, and the child is particularly vulnerable to reinfection during this time.

The results of this study show that shortening the course of antibiotic treatment has no adverse effect on outcome among children with acute otitis media. Several studies have attempted to investigate whether antibiotics are necessary in the treatment of at least some children with middle ear infection, but these studies do not provide sufficiently clear guidelines for taking decisions in individual children.²²⁻²⁴ It is likely that a number of children in this study did not require antibiotic treatment. The evidence about the clinical course will be helpful in planning a prospective trial of antibiotic placebo.

In circumstances where a general practitioner decides to use an antibiotic the conventional regimen of three to four times a day for one week has a ring of familiarity, but failure to complete a course of treatment for seven days is well recognized. The main advantage of a short course of treatment is that parents have a simpler regimen to follow, and the time during which the child has to take the antibiotic after the acute pain of earache has disappeared is reduced by several days. Where the general practitioner considers that prescribing an antibiotic is warranted a high dose, short course of amoxycillin is an alternative form of treatment that is acceptable to parents and suitable for children aged 3 and over.

This study was supported by a grant from the Wessex Regional Health Authority, and Bencard provided the amoxycillin used in the trial. We thank Dr David Machin, senior lecturer, department of community medicine, University of Southampton, for statistical advice, and our colleagues at

Aldermoor Health Centre for their advice and support throughout the many stages of the study. We are grateful to the cooperation of all general practitioners who participated in the trial.

References

1. Bain J. Acute otitis media. *Paediatrics* 1975;74:281-94.
2. Parham JF. Otitis media in children. *Paediatrics* 1972;49:917-41.
3. Bessner CD. Recent advances in the pathogenesis, diagnosis and management of otitis media. *Paediatr Child Health* 1980;16:101-10.
4. Horne VM, Pineda H, Lister RL. Otitis media: a clinical and bacteriological correlation. *Arch Dis Child* 1979;54:107-10.
5. Howard J, Nelson J, Chalkin BN, Jackson LH. Otitis media of infancy and early childhood. *Arch Dis Child* 1979;54:107-10.
6. Anonymous. Antibiotics for otitis media. *BMJ* 1979;3:1407.
7. Kline JF. Microbiology and non-invasive treatment of otitis media. *Ann Clin Res* 1979;11:101-10.
8. Anonymous. Non-compliance does not matter. *BMJ* 1979;3:1148.
9. Anonymous. Antibiotics for otitis media. *BMJ* 1979;3:1148.
10. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
11. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
12. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
13. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
14. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
15. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
16. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
17. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
18. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
19. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
20. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
21. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
22. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
23. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.
24. Anonymous. Compliance with antibiotic treatment. *BMJ* 1979;3:1148.

Audit Report

Do patients cash prescriptions?

In 1982 a small survey indicated that nearly 20% of patients failed to cash their prescription within a month of receiving it. I conducted a survey based on carbon copies of 966 prescriptions issued at 1217 surgery consultations by seven doctors in a group practice in Wirbury, which showed that 911 (92.4%) prescriptions had definitely been presented at local chemists. A postal questionnaire of the remaining 75 prescriptions elicited 65 replies; six further replies were obtained by home visiting. Forty eight of the 71 prescriptions were said to have been presented, and the remaining 23 replies included, "I got better before cashing the prescription" (five); "I could not afford the prescription fee" (four); "I did not think it was right for me" (four); "I lost the prescription" (two); and "I was not given a prescription" (two). Nobody replied, "I did not want a prescription," although this was an optional reply.

The prescribing habits of the six principals and trainee (myself) were also analysed. All the doctors were seeing 90-110 patients a week and had similar commitments as clinical assistants in the general practitioner hospital. The number of items prescribed by

each doctor varied from 168 in 182 consultations to 408 in 208 consultations. The use of generic drugs ranged from 7% to 68%. Eighteen per cent of the drugs prescribed would now be on the "blacklist." The heaviest prescribers had the lowest rates of non-enactment. Analysis of drug groups prescribed showed fairly even proportions of cardiovascular, antibiotic, and psychotropic drugs; the use of "symptomatic" treatments was much more variable but accounted for only 10% of the total.

Previous surveys using smaller numbers indicated that 10-20% of prescriptions were not presented.²⁵ In this study the figure was between 2.7% and 7.6%, whether or not all the drugs were consumed may be a different matter—IAM STUART, registrar, Family Medicine Training Programme, Victoria, Australia. (Correspondence to: 4 Martine Crescent, Great Yarmouth, Norfolk.) (Accepted 24 September 1985).

1. Baskin D. Do patients cash prescriptions? *BMJ* 1982;284:24-5.

2. Corbridge A, Ouse R. *Medical audit*. London: Baillière Tindall, 1977.

unusually as it is. Sunk costs are infinitely worse when the water supply is obtained from wells in the immediate neighbourhood than most important point, no information has been afforded. It is gratifying, and rather surprising, to learn that, in one large waterworks, earth-closets for 600 men have been found to work well; and if this system could be extended, it would be more wholesome and less wasteful than any other, as it is not used as the desodorizer. Where this is not possible, good latrines, with an automatic flushing chamber, are the best arrangement, and can generally be erected at a small cost. The replies, incomplete as they are, may be of use in calling the attention of medical officers of health and sanitary inspectors to the matter; they could even be able to prove whether the replies fairly represent the average condition, a point which must at present remain in some doubt.

(British Medical Journal 1985;291:403.)