Philosophical Medical Ethics

Consent

RAANAN GILLON

Vets do not bother with their patients' consent: why should doctors? To address this question it is important to state which meaning of "consent" is being used for the term is ambiguous. Under one definition it simply means agreement, acceptance, or assent. It is fairly obvious (though here unargued) that this meaning of consent is not relevant to medical interventions, whether these are investigations, treatments, or research. For medical interventions it is widely accepted that consent means a voluntary, uncoerced decision, made by a sufficiently competent or autonomous person on the basis of adequate information and deliberation, to accept rather than reject some proposed course of action that will affect him or her.¹⁻¹⁴ Consent in this sense requires action by an autonomous agent based on adequate information and is by definition informed consent.

Autonomy and informed consent

This analysis explains why vets do not bother with consent from their patients: their patients are not autonomous agents (or at least most of them are not—I remain agnostic about certain higher primates and dolphins), and they simply could not give consent. Given that most adult patients could give consent in this sense, why should doctors bother to ensure that they do? The most obvious answer is that generally we have a moral obligation to respect each other's autonomy, at least in so far as to do so is compatible with respect for the autonomy of others—a moral principle that I have previously shown to be supported from various moral perspectives. Doing things to other autonomous agents without their consent generally means overriding their autonomy. It is respect for people's autonomy, or self determination as Mason and McCall Smith¹⁵ and Kirby¹⁶ call it, that morally underpins the requirement of consent. Justice Kirby, then president of the Australian Law Reform Commission, in an excellent article on consent, wrote: "[T]he fundamental principle underlying consent is said to be a right of self determination: the principle or value choice of autonomy of the person The principle is not just a legal rule devised by one profession to harass another. It is an ethical principle which is simply reflected in legal rules because our law has been developed by judges sensitive to the practical application of generally held community ethical principles."¹⁶

According to the above definition, to consent to medical intervention a person requires sufficient information to be able to make an informed and deliberated choice, and it is in this context that doctors often object to the requirement for such consent. Patients, they say, are unnecessarily alarmed and their medical state unnecessarily impaired if they have to be given information about their diagnosis or prognosis or risks associated with their proposed

management and treatment. The alarm is compounded with confusion, the argument continues, if differential diagnoses and risks have to be mentioned and made intolerable if the patient is expected to choose between the options. As Dr Ingelfinger exclaimed, he as a patient would not want "to be in the position of a shopper at the casbah who negotiates and haggles with the physician about what is best." Indeed, the doctor "who merely spreads an array of vendibles in front of the patient and then says 'go ahead and choose, it's your life' is guilty of shirking his duty, if not of malpractice."¹⁷

Three counterarguments

There are at least three problems with such arguments. The first is that it is dangerous for a doctor to extrapolate from his own case to generalisations about his patients (or about anybody else). The fact that the good doctor would feel like that does not ensure or even make probable that any particular patient feels like that; and, as Dr Ingelfinger's argument implicitly acknowledges, the doctor should be trying to meet his patient's wishes rather than his own. As a wise philosopher once wrote, it is not putting yourself into another's shoes that is morally relevant, it is understanding what it is like for that other person to be in his or her own shoes that is morally important.

The second problem is empirical. Is it true that patients are generally made more alarmed by open minded and sympathetic provision of information and an invitation from the doctor to express their own preferences in so far as they have any? I doubt it. In this regard Dr Ingelfinger offers a parody of respect for the patient's autonomy when he suggests that the doctor would "haggle" with the patient or alternatively merely tell the patient the available options and then say "Go ahead and choose, it's your life." The doctor who really respected his patient's autonomy would discover in a sensitive way, which did not demand a particular answer, what and how much each patient really wanted to know and how much he wished to participate in the decision making. Some patients are, like Dr Ingelfinger, undoubtedly keen not to be given unpleasant information and to leave all the decision making to the doctor; if the doctor acts accordingly, thus respecting their autonomy, such patients will probably be happier. Other patients, however, do want to know about their medical condition and its implications and about alternative options; they also wish to be active in decision making when it affects them. Such patients will probably feel more at ease, trusting, and happy if the doctor has respected their wishes and if they have deliberately, comprehendingly, and willingly consented to their doctor's plan of management and have been "incorporated" as key figures in the medical management team. Of course, this second counterargument is no more than an empirical claim based on clinical impressions and common sense, but then so purports to be the claim it opposes—namely, that patients will be more miserable if they are given information they want and genuine opportunities to make their own decisions about their management.

Imperial College of Science and Technology, London SW7 1NA

RAANAN GILLON, MB, MRCP, director, Imperial College Health Service, editor, *Journal of Medical Ethics*, and associate director, Institute of Medical Ethics

The third problem is that even if it were true that in some circumstances a patient would be made more miserable by being given information about his condition and the risks of alternative methods of management, if that is what the patient truly wishes to know is the doctor not morally obliged to tell him? In my articles on autonomy and paternalism I outlined what seem to me to be cogent arguments from utilitarian as well as deontological perspectives for respecting people's autonomy in so far as to do so is compatible with respecting the autonomy of others. These arguments have as much weight against withholding of relevant information required for consent as they do against lying or otherwise deceiving the patient (indeed the deliberate withholding of information relevant to decision making is a form of deceit) and against breaking confidentiality in the patient's own interests.

Doctors' licence under English law

A further set of common medical counterarguments to such emphasis on respect for patients' autonomy in the context of consent is that English law, at least with regard to consent to medical treatment, leaves doctors to decide, in the context of their therapeutic (as distinct from research) relationships, how much information they ought to give patients when obtaining their consent to treatment.

Accounts of what the law stipulates in any particular jurisdiction, however, do not in themselves provide moral justification for a moral claim. At most they can be used as part of a moral claim that there is a general presumption that it is a good thing to obey just laws. As I have argued previously, the possibility that there might be unjust laws (never mind the fact that there are) shows that it is not necessarily a good thing to obey all laws. Similarly, the possibility that there might be morally bad laws shows that what is legal is not necessarily morally justifiable.

The second point is that even if English law did leave it to doctors to decide how much information ought to be given to patients in the course of obtaining their consent to treatment doctors would still be obliged to try to resolve the moral problem: to be given the legal responsibility of making a moral decision is precisely not to be absolved from doing so.

It may, however, be worth noting that English law has recently been changed considerably by the House of Lords' appeal judgment in the Sidaway case.¹⁸ Until that case the law seemed to leave decision making to doctors, although there is no legal unanimity about this.¹⁹ Thus according to Lord Scarman's account of the relevant leading case, in which a Mr Bolam claimed, inter alia, that he had not been adequately informed about the risks of his (then unanaesthetised) electroconvulsive therapy, doctors would not be held negligent in law provided they acted "in accordance with the practice accepted by a responsible body of medical men," even if there existed a different practice advocated by another responsible body of doctors.¹⁸ In the Sidaway judgment, however, only one of the five law lords (Lord Diplock) accepted the so called Bolam doctrine in its unmodified form. Each of the four others judged that the "professional standard" position of the Bolam doctrine required modification towards allowing patients to make their own decisions based on adequate information about treatments that included "material" (Scarman) or substantial (Bridge and Keith) risks, or "which might have disadvantages or dangers" (Templeman). Lords Bridge and Keith added that "When questioned by a patient of apparently sound mind about risks involved in a particular treatment proposed the doctor's duty was to answer both truthfully and as fully as the questioner required."¹⁸

Consent in therapeutic research

Occasionally the argument is heard that although properly informed consent is required in the context of non-therapeutic research on patients, it is not needed in the context of treatment or therapeutic research. The manifold medicomoral issues associated

with medical research are complex but in this context suffice it to assert that doctors are no less morally obliged to respect the autonomy of their patients than that of their research subjects. The moral differences between the two categories relate not to respect for autonomy but to two quite different issues. The first is that there is a substantial danger that research subjects will assume that doctors' normal beneficent Hippocratic concern to do their best for their particular patients will also apply to their research subjects. In the case of non-therapeutic research this cannot be the case (by definition), and doctors therefore have a particularly strong moral obligation to make this clear to such subjects. The second, though related, point is that in the normal therapeutic relationship patients can properly assume that any risks to which the doctor proposes to subject them will be proposed only in the light of an analysis of risk and benefit that is favourable for the particular patient. Such an assumption would again be quite mistaken in the case of non-therapeutic research (in which the benefits, if any, will accrue to future patients, while the risks are taken by the research subjects). But while both these points indicate that doctors should make it crystal clear to patients when they are not acting in a therapeutic role, neither supports the claim that when doctors are acting therapeutically they need not obtain properly informed consent.

In the last three articles I have argued that in normal cases respect for patients' autonomy takes moral priority over medical beneficence and generally precludes lying to or otherwise deceiving patients even in their own interests, breaking their confidences even in their own interests, and failing to obtain their adequately informed consent to medical intervention even in their own interests. In the next article I shall look at circumstances in which the principle of respect for autonomy does not seem to have priority over beneficence.

References

- 1 Wilkinson AW. Consent. In: Duncan AS, Dunstan GR, Welbourn RB, eds. *Dictionary of medical ethics*. London: Darton, Longman and Todd, 1981:113-7.
- 2 Taylor P. Consent, competency and ECT: a psychiatrist's view. *J Med Ethics* 1983;9:146-51.
- 3 Skegg PDG. Informed consent to medical procedures. *Med Sci Law* 1975;15:124-8.
- 4 Herbert V. Informed consent—a legal evaluation. *Cancer* 1980;46:1042-3.
- 5 Dunstan GR, Sellar MJ, eds. *Consent in medicine*. London: King's Fund Publishing Office and Oxford University Press, 1983.
- 6 Beauchamp TL, Childress JF. *Principles of biomedical ethics*. 2nd ed. Oxford: Oxford University Press, 1983:69-93.
- 7 Lidz CW, Meisel A, Zerubavel E, Carter M, Sestak RM, Roth LH. *Informed consent*. London: Guilford Press, 1984.
- 8 Culver CM, Gert B. *Philosophy in medicine*. Oxford: Oxford University Press, 1982:42-63.
- 9 Gorovitz S. *Doctors' dilemmas*. London: Collier Macmillan, 1982:34-54.
- 10 Gorovitz S ed. *Moral problems in medicine*. 2nd ed. Englewood Cliffs: Prentice-Hall, 1983:153-91.
- 11 Reich WT, ed. *Encyclopedia of bioethics*. London: Collier Macmillan, 1978:751-78.
- 12 Levine RJ. *Ethics and regulation of clinical research*. Baltimore: Urban and Schwarzenberg, 1981:69-115.
- 13 Bankowski Z, Howard-Jones N, eds. *Human experimentation and medical ethics*. Geneva: Council for International Organisations of Medical Sciences, 1982:16-121.
- 14 Veatch RM. *Case studies in medical ethics*. Cambridge, Massachusetts: Harvard University Press, 1977:290-316.
- 15 Mason JK, Smith RAM. *Law and medical ethics*. London: Butterworths, 1983:120.
- 16 Kirby MD. Informed consent: what does it mean? *J Med Ethics* 1983;9:69-75.
- 17 Ingelfinger FJ. Arrogance. *N Engl J Med* 1980;303:1507-11.
- 18 Anonymous. Sidaway v Bethlem Royal Hospital and the Maudsley Hospital Health Authority and others (law report). *The Times* 1985 Feb 22:28.
- 19 Norrie K McK. Medical negligence: who sets the standard? *J Med Ethics* 1985;11:135-7.

(Accepted 26 July 1985)

Is there any risk of lead poisoning from using pewter vessels?

Old pewter is a mixture of tin and lead, the lead being added both for cheapness and to make the mix easier to work. In the past pewter plates and vessels were widely used and almost every household would have at least one pewter item. There is a theoretical risk that when using pewter plates some of the metal might be scraped off and swallowed, but I doubt whether this would have posed a serious threat to health. Pewter drinking vessels are another matter, however, and lead poisoning has resulted from drinking home made wine (of low pH) from pewter goblets. Modern pewter does not contain lead so that beer drinkers need have no fear, but it would be safer not to use old—that is pre-1900—pewter vessels for eating or drinking.—H A WALDRON, senior lecturer in occupational medicine, London.