- 39 Yeung CK, Smith RC, Hill GL. Effect of an elemental diet on body composition: a comparison with intravenous nutrition. *Gastroenterology* 1979;77:652-7.
- 40 Askanazi J, Nordenstrom J, Rosenbaum JF, *et al.* Nutrition for the patient with respiratory failure: glucose versus fat. *Anesthesiology* 1981;54:373-7.
- 41 Wolfe RR, Allsop JR, Burke JF. Glucose metabolism in man. Responses to intravenous glucose infusion. *Metabolism* 1979;28:210-6.
- 42 Burke JF, Wolfe RR, Mullaney CJ. Glucose requirements following burn injury. *Ann Surg* 1979;190:274-85.
- 43 Wolfe RR, Burke JF. Isotopic studies of glucose metabolism in humans. *Journal of Clinical Surgery* 1982;1:180-93.
- 44 Wilmore DW. Role of lipid as a source of non-protein calories. In: Johnston IDA, ed. *Advances in parenteral nutrition*. Lancaster: MTP Press, 1977:195-207.
- 45 Anonymous. Adult parenteral nutrition: which preparations? *Drug Ther Bull* 1981;18:85-8.
- 46 Hodges RE. The fat-soluble vitamins. In: *Report of the second Ross conference on medical research*. Columbus, Ohio: Ross Laboratories, 1981:61-3.
- 47 Jeejeebhoy KN. Trace elements in human nutrition; discussion. In: *Report of the second Ross conference on medical research*. Columbus, Ohio: Ross Laboratories, 1981:80.
- 48 Smith JE, Goodman WS. Vitamin A metabolism and transport. In: *Present knowledge in nutrition*. 4th ed. New York: The Nutrition Foundation, 1976.
- 49 Shenkin A. Additives in parenteral nutrition. *Acta Chir Scand* 1980; suppl 507:350-3.
- 50 Jeejeebhoy KN. Micronutrients—state of the art. In: Kleinburger G, Deutsch E, eds. *New aspects of clinical nutrition*. Vienna: Karger, 1983:1-24.
- 51 Ryan JA, Abel RM, Abbott WM, *et al.* Catheter complications in total parenteral nutrition. *N Engl J Med* 1977;290:757-61.
- 52 Keohane PP, Attrill H, Northover J, *et al.* Effect of catheter tunnelling and a nutrition nurse on catheter sepsis during parenteral nutrition. A controlled trial. *Lancet* 1983;iii:1388-90.
- 53 Pollack FP, Kadden M, Byrne WJ, *et al.* 100 Patient years' experience with the Broviac Silastic catheter for central venous nutrition. *JPEN* 1981;5:32-6.
- 54 Welch GW, McKell DW, Silverstein P, *et al.* The role of catheter composition in the development of thrombophlebitis. *Surg Gynecol Obstet* 1974;138:421-4.
- 55 Deitel M, Kaminsky V. Total parenteral nutrition by peripheral vein—the lipid system. *Can Med Assoc J* 1974;111:152-4.
- 56 Freeman JB. Peripheral parenteral nutrition. *Can J Surg* 1978;21:489-92.
- 57 Sanderson I, Kuksis A, Jeejeebhoy KN. Peripheral parenteral nutrition with lipid providing 83% of calories. *Gastroenterology* 1973;64:796.
- 58 Ellis BW, Stenbridge R de L, Fielding LP, Dudley HAF. A rational approach to parenteral nutrition. *Br Med J* 1976;i:1388-91.
- 59 Selzar MH, Assaadi M, Coco ET, *et al.* Use of a simplified standardised hyperalimentation formula. *JPEN* 1978;2:28-30.
- 60 Vanderveen WT, Robinson AL. Total parenteral nutrition solution utilising amino acid sources with and without pre-added electrolytes. A time and cost comparison. *JPEN* 1979;3:84-8.
- 61 Oxford Parenteral Nutrition Team. Total parenteral nutrition: value of a standard feeding regimen. *Br Med J* 1983;286:1323-7.

(Accepted 25 September 1985)

Personal Paper

Stammering

K F HULBERT

I have never known what it is to speak freely without the knowledge that lurking ahead somewhere is an unexpected and insurmountable log jam of words. One of my earliest recollections is of going to a sweet shop to spend my "Saturday penny" and coming away in rage and frustration with something I did not want because the words "liquorice laces" would not come out.

It did not seem to worry me, or anyone else, until I got to the age of about 9. Then some would ask, "What will you do when you grow up?" and there was the inevitable fun making at school, which I hotly resented. The recommended treatment was elocution, which I liked because it entailed reading and learning poetry. There were breathing exercises and relaxation, which all helped the flow of language once the flow was established. But it was the apparent trivialities of conversation that produced such enormous and frightening problems. Saying hello or goodbye, or introducing myself or someone else, was often a nightmare that I would do almost anything to avoid. Silly little things like being asked by an adult the way, or what my name was, I would go to extremes to evade. Friends were never short of advice, such as to talk with a mouthful of pebbles like Demosthenes, the patron saint of stammerers—it can be done but is difficult and quite hazardous. If anyone is to occupy the prime place as the hero and example for all stammerers, it is surely King George VI. They say that stammerers never stammer when they sing, which is largely but not entirely true, but to suggest that you should go through life singing all conversations may be good comic opera but is not a practical proposition. Then, too, there were the psychotherapists, all delightful people with whom I spent many an hour but with no apparent improvement.

This blight on conversation meant that at school I was a loner, had

few friends, was passed over in reading aloud in class, never took part in school plays, and never became a prefect. Talking to myself, aloud, was marvellous, and I would go for long country walks reciting poetry. As I got older the social handicap became more apparent. Standing in a queue shopping or at a ticket office with people waiting behind me was most trying, and I would often resort to a pencil and paper to save embarrassment. Many well meaning people would try to help by guessing what I was trying to say, and this could result in a quite deplorable situation—if they guessed wrong then I had the additional problem of telling them and starting again. If they guessed right I wanted to thank them, and this was one more problem to overcome.

Then there loomed the awful problem of "What are you going to do when you leave school?" Adverts in papers like "Try my method; I can cure you as I cured myself" promised much and disappointed greatly. Some I am sure depend on the natural spontaneous remission in many cases. This all built up to a crisis that produced the state of mind in which I seemed to say, "Never mind what they say, I want to be a doctor." An understanding headmaster, a sympathetic medical school dean, success in gaining an entrance scholarship including success in a viva voce exam, and I was in. From then on it was by no means easy, and viva voce exams in the finals appeared like the Rocky Mountains to the builders of the Canadian Pacific Railway, but it was done.

Once qualified I encountered an entirely new situation. As long as I was with a patient there were few problems. Speaking came naturally because I was the doctor and could speak in my own time, and the patient had to listen. Conversation is not only speaking, it is also holding the attention of the other. If only people would listen and wait it would be so helpful. The concentration sometimes required to navigate a treacherous sentence is such that even if the heavens should fall I would still be concerned with the letter B coming along like a hurdle in a race. Ivan the Terrible, it is said, had a walking stick with a sharp spike with which he transfixed the foot of anyone he wanted to talk to. How I envied him on occasions.

Shoreham, Kent TN14 7TX

K F HULBERT, FRCS, DA, former consultant orthopaedic surgeon

Higher qualifications extended me to the full, and beyond. Then came the army, where the comradeship carried me along with little to worry about. Being a consultant with the prestige that went with it greatly helped in the environment of hospital practice, but there were still the social handicaps in the trivia of conversation. The speaking problems that to the “normal” person are so trivial as to be unnoticed are to me the frightening obstacles, whereas a big problem may well seem a molehill instead. This has been proved so often by the ease with which I have talked to patients and colleagues, only to be floored soon after by having to ask for something in a shop or answer the telephone.

Now that I can look back from the haven of retirement it is possible to make certain comments on this strange and baffling impediment. Yes, it would have been nice not to have to contend with this frustrating and embarrassing condition. It still is most frustrating and embarrassing. My friends tell me that it is much less so to others than it is to me, but I still have to contend with the desire to retire into a shell. There are many ways in which people who stammer can be helped, but the one way above all others is to give us complete attention while we concentrate all our efforts on the verbal steeplechase that lies ahead. To make a snatch at a difficult word and to experience the pleasant surprise of clearing the hurdle, only to find that your addressee was not listening, is made worse by the major disaster as you crash at the fence on the second attempt. We all learn by experience, and there are many ways of circumventing or overcoming obstacles. There are numerous alternative words, or

even quaint phrases, which may sound strange to others but all help in this strange hurdle race of conversation.

I have some regrets, now seen more clearly with hindsight. If only all of us as children could start off under the care of a paediatrician who regards stammering as a symptom rather than a diagnosis. We as children in my time were at the mercy of a diversity of therapists, each confident that his method was sure to succeed if you followed the course with application. If you were not cured then it was your fault; you can speak perfectly well when alone, so it is psychological, it is yourself that is at fault. I am sure that this is a cause of much unhappiness to a lot of children.

The more I experience it the more I am sure that there is a neurological basis for this strange and yet ill understood malady. To suggest that it is due to some Freudian complex, or a subtle way of avoiding something that we do not like, or the result of some far distant event in infancy, is just not good enough. Please, as doctors, take us and our problem seriously and do not farm us out in strange pastures. Looking back now on a lifetime of this handicap, in which I have known no other, it would have been nice to be without it. It warps you in certain ways, cramps you in some of the social aspects of life, and even causes some misery at times. Having said all this, I consider my life to have been interesting and challenging and even to have had its funny side, and it has gained me many many friends. Strange as it may seem, I would not now have had it otherwise.

(Accepted 11 September 1985)

Clinical Topics

Sexual dysfunction in Asian couples

DINESH BHUGRA, CHRISTINE CORDLE

No information is available on the prevalence of sexual dysfunction in Asian couples in the United Kingdom. There are only two papers concerned with Asians out of over 2000 papers published on sexual dysfunction between 1966 and 1985.^{1,2} This study looks at 32 Asian couples who presented to a sexual dysfunction clinic between 1977 and 1983. The drop out rate was high, with lower than generally reported success, and we consider some of the factors that may have been responsible.

Present study and results

The sexual dysfunction clinic at Leicester General Hospital covers the inner city population, which includes some 60 000 Asians; many of these speak Gujarati, Punjabi, Urdu, and Hindi. Referrals are accepted from general practitioners, psychiatrists, physicians, surgeons, and social workers.

The details of all 23 men and nine women with Asian names referred to the

clinic from 1977 to 1983 were retrieved from the referral register. The table shows the presenting symptoms. In at least four couples there were problems in both partners but only the presenting partner's complaint was taken into account. In the men the duration of symptoms ranged from six months to 25 years, and 17 men (73.9%) had had their problem longer than five years. By contrast, the duration of the women's symptoms varied from one to five years. At least nine of the men and two women had initially attended for screening seeking medication and somatic explanations for their problems. Five or more men and two women also had language difficulties.

Of all 32 patients, 14 (43.8%) gave up attending the clinic; some left immediately after screening and others dropped out during treatment. At least four patients dropped out because they or their partners had gone to the Indian subcontinent for a long holiday.

Of the 18 couples taken on for treatment, five (27.7%) reported an improvement.

Presenting symptoms in 32 Asians (23 men, nine women) attending sexual dysfunction clinic

	No (%) of men	No (%) of women
Premature ejaculation	14 (60.9)	6 (66.7)
Secondary erectile dysfunction	14 (60.9)	7 (77.8)
Delayed ejaculation	1 (4.3)	1 (11.1)
Primary erectile dysfunction	6 (26.1)	0
Low drive	5 (21.7)	0
Others:		
Sexual identity	1 (4.3)	
Urethral discharge	1 (4.3)	
Dyspareunia		
Lack of interest		
Primary anorgasmia		
Secondary anorgasmia		

Academic Unit of Psychiatry, Carlton Hayes Hospital, Leicester LE9 5ES
DINESH BHUGRA, MB, MRCPsych, registrar in psychiatry

Hadley House, Leicester General Hospital, Leicester LE5 4PW
CHRISTINE CORDLE, MSc, DCP, principal psychologist

Correspondence to: Dr Bhugra.