PRACTICE OBSERVED

Practice Research

Epidemic of AIDS related virus (HTLV-III/LAV) infection among intravenous drug abusers

J R ROBERTSON, A B V BUCKNALL, P D WELSBY, J J K ROBERTS, J M INGLIS, J F FEUTHERER, R P BRETTE

Abstract

Stored blood samples from 164 intravenous drug abusers who attended a Scottish general practice were tested for HTLV-III/LAV (human T cell lymphotropic virus type III/lymphadenopathy associated virus) infection. Of those tested, 83 (51%) were seropositive, which is well above the prevalence reported elsewhere in Britain and Europe and approaches that observed in New York City. The timing of taking samples of negative sera and continued drug use suggest that as many as 85% of this population might now be infected. The infection became epidemic in late 1983 and early 1984, thereafter becoming endemic. The practice of sharing needles and syringes correlated with seropositivity, which, combined with the almost exclusive intravenous use of heroin and other behavioural patterns, may explain the high prevalence of HTLV-III/LAV infection in the area. Rapid and aggressive intervention is needed to control the spread of infection.

Edinburgh Drug Addiction Study, West Granton Medical Group, Edinburgh EH4 4PL

J R ROBERTSON, MR, MB, general practitioner
J J K ROBERTS, MR, MB, general practitioner
A B V BUCKNALL, BA, research assistant

City Hospital, Edinburgh EH10 5SB
P D WELSBY, MB, PhD, consultant physician, communicable diseases unit
R P BRETTE, MR, PhD, consultant physician, communicable diseases unit
J M INGLIS, MR, PhD, director, regional virus laboratory

Bacteriology Department, University of Edinburgh, Edinburgh EH8 9AG
J F FEUTHERER, MR, MB, PhD, senior lecturer, hepatitis reference and clinical virology laboratories

Introduction

HTLV-III/LAV (human T cell lymphotropic virus type III/lymphadenopathy associated virus) seropositivity has been described among groups of intravenous drug abusers in the United States and lately in Europe. Recently, in a laboratory study of drug users, in which blood samples from patients who had attended various departments of the Royal Infirmary, Edinburgh were used, 38% of heroin users were infected with the virus.¹ It has not yet been explained why there is such a high incidence of infection in heroin users from this city. Previous research has correlated seropositivity with the number of injections but not with sharing needles and syringes, and it is assumed that sexual transmission of the infection may also be important. A combination of these factors, along with local practices and regional variations in the availability of sterile equipment, may explain the wide variation in seropositivity among different groups of heroin users, which is already known to exist in the United States and Europe.²

The West Granton Medical Group serves a population of about 18 000 patients in a deprived area of Edinburgh that is known to have many intravenous drug abusers. The policy has always been to manage drug users who are known to the practice before the onset of drug misuse but to resist taking heroin users for treatment who approach the practice from outside a specific geographical area of the city. In 1982 an outbreak of hepatitis B infection in the heroin users who attended the practice led to routine blood testing for hepatitis B markers, the epidemic being attributed to an increase in sharing needles and syringes owing to an acute shortage after the local, legal retail supplier was closed and a subsequent unofficial prohibition by pharmacists. More recently, increasing anxiety about the risk of the acquired immune deficiency syndrome (AIDS) to intravenous drug abusers and their spouses and sexual contacts and to the children of infected women led to our decision to test retrospectively stored serum samples for the HTLV-III/LAV antibody to obtain information to manage these patients. This study

impurities provided a source of further injectable heroin when these could be soaked in water. These practices appear to have continued through 1984 and have been moderated only recently owing to anxiety about the risk of AIDS.

From National Health Service identification numbers it could be shown that 88% (144) of the study group were born in Edinburgh and only 2% outside Scotland. Specific questioning about travel within the United Kingdom and abroad up to 1984 indicated that 10% of the group had travelled internationally, one to the United States and North Africa but denying drug use in either place, the other to Germany in 1976. The group as a whole rarely travelled outside Edinburgh.

TABLE 1—Reported locations of heroin use of 70 heroin users up to January 1984

Location	No. % of heroin users
Edinburgh practice area	70 100
Edinburgh other area	4 6
London	5 7
Amsterdam	1 1
Geneva	1 1
Geneva	1 1
Geneva	1 1
Newcastle upon Tyne	1 1
Germany Berlin	1 1

Despite recent reports of a brief influenza like illness shortly after initial exposure a study of case notes indicated no such phenomenon among these patients.³ At the time of writing no patient from the study group had been diagnosed as having AIDS, although a small but growing number are showing AIDS related illnesses such as persistent generalised lymphadenopathy and candidiasis.

Discussion

HTLV-III/LAV infection was observed in half of the intravenous drug abusers studied; this is well above the prevalence observed in previous European studies and approaching that reported in New York City.¹ Although it is not clear how the virus was introduced into this population, the rapid spread of infection (see fig 2) appears to have resulted from a combination of the almost exclusive intravenous use of heroin, the frequency of sharing equipment owing to difficulties in obtaining sterile needles and syringes locally, and injection practices that increase the risk of exposure to the virus. Indeed, the frequency of needle and syringe sharing has been positively correlated with seropositivity and seems to be the most likely cause of the high incidence of infection. Preventing the sharing of equipment may therefore be the quickest and most efficient means of containing the spread of HTLV-III/LAV in drug users. The rapidity with which the epidemic spread in this study population, in addition to the presumed risk of infection from drug users to the non-drug using heterosexual population, shows the need for aggressive intervention by all agencies in the community.

Managing HTLV-III/LAV infection in general practice necessarily includes both preventing the spread of the virus and dealing with the medical, social, and psychological problems of those already infected. The general practice in this study has a policy of advising intravenous drug abusers who attend the practice of the risk of infection to themselves, their sexual contacts, and children. Appropriate counselling is offered that deals with reducing the risk through altered drug use and sexual behaviour, the importance and desirability of antibody testing, and the specific consequences of infection. In attempting to further prevent the spread of the AIDS related virus it has been the policy of two clinical officers to supply clean equipment to the drug users on an exchange basis, a clean set being given only on the return of the old one. Local education campaigns, including providing literature with guidelines for safer sex and safer drug use, have been initiated by general practitioners through local self-help groups. An alternative means of reducing equipment sharing, prescribing non-injectable methadone as a heroin substitute, has not yet been introduced because it is

opposed to movements away from prescribing opiate drugs to drug users. This and other interventions, however, must be considered soon and introduced where appropriate to contain the spread of infection.

Many AIDS related medical and social problems are being dealt with, especially with reference to women of child bearing age and those who are pregnant, who on the whole remain reasonably healthy. At the time of writing two women had been recommended terminations, and two had given birth to infants found to be seropositive. An uncertain proportion of these infants will be born with develop AIDS, and a few more will develop AIDS related illnesses which do not fulfil the criteria of AIDS itself. Links with local clinics that specialise in such problems have been established. Recent evidence suggests that heroin users are more at risk of developing AIDS once infected than male homosexuals.⁴ In a city such as Edinburgh, with large numbers of heroin users and a high prevalence of HTLV-III/LAV, but with a prevalence of infection among male homosexuals undoubtedly of less than 5% (A McMillan, personal communication, 1985), heroin users may represent most clinical AIDS cases seen. AIDS is likely to become the leading cause of death among the patients examined in this study.⁵

Our results indicate that geographical variations are likely in the prevalence of HTLV-III/LAV infection among intravenous drug users owing to differing equipment sharing practices and differing times of arrival of the virus into populations of drug users. A city where the seropositivity is known to be low but where needle and syringe sharing is routine among heroin users must expect the virus to spread rapidly in the absence of immediate intervention.

We thank the doctors and staff of the West Granton Medical Group for their help in collecting samples and information, Dr E Edmond and J D Dickson, University of Edinburgh Bacteriology Department, and I Collocott from the Regional Virus Laboratory, City Hospital, Edinburgh. This work was supported by the Scottish Home and Health Department and the Rosebank Hepatitis Research Fund.

References

1. Spivey TJ, De Jorio DC, Marmor M, et al. Prevalence of antibody to lymphadenopathy associated virus among drug detoxification patients in New York. *N Engl J Med* 1984; 311: 44-8.
2. Auer R, Bressi F, Saito M, et al. Light and light microscopy in human T cell lymphotropic virus (HTLV-III) in lymphadenitis syndrome and risk for AIDS in Italy. *Br Med J* 1985; 290: 140-6.
3. Feutheer JF, Edmond J, Vennart D, Dickson JD, Burt L. HTLV-III antibodies in Edinburgh drug addicts. *Lancet* 1985; 1: 1278-9.
4. Melnick M. The natural history of human T cell lymphotropic virus type III infection: the acquired immunodeficiency syndrome (AIDS) and suggestions for its control in drug abusers. *The Journal of Infectious Diseases* 1984; 150: 1-10.
5. Robertson JR. Drug abuse in context with general practice. *Br Med J* 1985; 290: 14-5.
6. Friedman R, Hirsch B, Saito M, et al. HTLV-III antibodies in human T cell lymphotropic virus (HTLV-III) in lymphadenitis syndrome. *AIDS* 1985; 1: 145-147.
7. Cooper D, Gold L, MacLean C, et al. HTLV-III antibodies in human T cell lymphotropic virus (HTLV-III) in lymphadenitis syndrome. *Lancet* 1985; 1: 140-1.
8. Tachibana M, Hirsch B, Hirsch A, et al. HTLV-III antibodies in human T cell lymphotropic virus (HTLV-III) in lymphadenitis syndrome. *Lancet* 1985; 1: 140-1.
9. Burt L, Auer R, Robertson JR. Deaths of heroin users in a general practice population. *Br Med J* 1985; 290: 140-1.

Accepted 16 January 1986

100 YEARS AGO

A correspondent sends us the following:—"An old man, nearly octogenarian, who has been in bed for twenty seven years, being a harmless monomaniac, having the delusion that His Satanic Majesty always stood at his door to prevent him from going out, suddenly one morning early in June, took to his head that the Devil was gone, whereupon he got out of bed, and, finding nothing on his list, walked down to the quay (nearly a quarter of a mile) to inspect the ships. Having been in good working order, he returned to his room, and, although a boat put out from a vessel, he saw aboard a noble-looking fellow, who was taken to a public-house, and a medical man was sent for. He was taken to the house, where he remained two or three days, and, after the better of his ship. He has been in church, and is now about to resume his former calling, that of a preacher." (*British Medical Journal* 1886; 2: 261.)

therefore examines the behaviour, injection practices, and seropositivity of a geographically selected population of known heroin users.

Method

The study group comprised 164 patients of the practice who were known to be or to have been heroin users. There were long term users, casual users, and those who had experimented with heroin but never developed an addiction, making the population representative of heroin users as a whole. Sera that had been taken from each heroin user most recently and stored in one of the region's two virus laboratories were tested for HTLV-III/LAV antibody using commercially available ELISA, and if positive this was confirmed by immunofluorescence or Western blotting technique, or both. If the most recent serum was positive, previous samples, when available, were successively tested until a negative result was obtained, thereby identifying the period during which seroconversion occurred. Hepatitis B marker status was also ascertained. Detailed information about the patient's drug practices, lifestyle, and behaviour was recorded on standard case summaries, having been extracted from the practice case notes and interviews. Most of these patients attended the practice regularly often weekly over several years, which allowed insights to be made into their lifestyle and behaviour that is rarely possible in hospital studies.

Results

Serum from 164 heroin users was tested. Of these, 83 (51%) patients were HTLV-III/LAV positive, 60 men and 23 women (mean SD) age 24 (4.7) years. Eighty one (49%) patients were seropositive, 45 men and 26 women (mean SD) age 29.9 (5.6) years. There were no sex differences between the seropositive and seronegative groups, but the seropositive patients were significantly older ($t = 4.44$, $p < 0.01$). The mean age of onset of intravenous drug abuse in 77 seropositive patients was 19.1 years, and in 60 seronegative patients for whom this information was available it was 19.9 years. The mean duration of heroin use could therefore be estimated as 4.9 years for seropositive patients and 6.1 years for seronegative patients. Figure 1 shows the cumulative number of first recorded seropositive sera; the first positive serum being taken in September 1983. Similarly, the period

may have seroconverted after the final serum sample was taken. If all of these were now infected there would be an overall prevalence of HTLV-III/LAV of 85%. Interestingly, the prevalence of hepatitis B marker was 84%, 27 patients being negative for both hepatitis B markers and HTLV-III/LAV antibody.

FIG 2—Last negative and first positive HTLV-III/LAV antibody tests for 83 intravenous heroin users against time.

Clinical and epidemiological information drawn from the patients' case summaries indicated consistent and repeated patterns and practices of drug misuse. The primary route of heroin administration was almost exclusively intravenous (98%), and 83% of 116 users for whom information was available reported sharing equipment (table 1). The serological differences between those who reported sharing needles "usually," "sometimes," or "never" were statistically significant ($\chi^2 = 12.2$, $df 2$, $p < 0.01$), indicating a correlation between the frequency of sharing equipment and seropositivity.

TABLE 1—Self reported sharing of needles and syringes in 164 heroin users with and without antibodies to the HTLV-III/LAV virus

Sharing needles and syringes	No. % who were HTLV-III/LAV antibody positive	No. % who were HTLV-III/LAV antibody negative	Total
Usual	36 52	12 20	48
Sometimes	27 49	21 35	48
Never	14 17	16 26	30
Total	83 100	49 100	164 99

In an attempt to find the causes of the outbreak and rapidity of onset of the epidemic a group of about 40 early seroconverters were identified who had been interviewed before January 1984. The group formed an intimate equipment sharing community between 1980 and 1983. Initially using personal equipment only, the sharing of needles and syringes became routine as the supply of clean equipment failed. Particularly important were consistent reports of gatherings of 10 to 20 drug users sharing a single needle and syringe during 1983—not unlike the "showing galleries" described in the United States. Because the equipment often belonged to the heroin supplier the people who attended such gatherings each day would vary, increasing the possibility of infection. The equipment was reported as best to be used over from the "showing galleries" each day, and was used until sterilisation despite the routine practice of "washing" (drawing blood back into the syringe after injection to flush out any remaining heroin). Drawing heroin solution through cigarette or cotton wool filters to absorb

FIG 1—Cumulative number of first positive HTLV-III/LAV antibody test results among 83 intravenous drug abusers by quarters.

of time during which seroconversion occurred was known for 33 patients (fig 2). Twenty five (32%) of the seropositive patients and 21 (35%) of the seronegative patients for whom information was available were known to have begun injecting heroin after the infection (thought to have entered this population). There is no indication therefore that seropositivity is related to the duration of drug misuse. The interpretation of seropositive status is complex, however. Twenty antibody negative patients had had no serum taken since the onset of the epidemic, and of those who had, 11 were known to have injected drugs within three months and 26 to have continued to inject drugs after their final serum sample was taken. Thus any of these 57 patients

Audit Report

Monitoring care of patients with hypothyroidism in a Yorkshire practice

Recent papers have pointed out the desirability of regular monitoring of the adequacy of thyroxine replacement treatment in patients with hypothyroidism and the importance of undertaking both clinical and biochemical assessment. Himmelfarb suggested that this should be undertaken yearly, and at the same time haematological screening should be carried out for pernicious anaemia.¹ The names of all patients who were receiving thyroxine replacement treatment were obtained from the practice computer used for repeat prescriptions. From the records of these patients, the care of 32 patients with hypothyroidism was reviewed. This number gives prevalence rates for hypothyroidism in the practice of 1.3 per 1000 men and 5.8 per 1000 women, which is similar to that reported elsewhere.

The aetiology of the hypothyroidism was idiopathic in 22 (69%) patients, followed partial thyroidectomy in three (9%) and treatment of thyrotoxicosis with radioactive iodine in three (9%), was associated with lithium carbonate prophylaxis for manic depressive illness in three (9%), and was congenital in one. All patients had

been seen during the previous year, but systematic clinical examination of their thyroid state had not been carried out, or at least was not recorded in their notes. Ten (31%) patients had had biochemical thyroid assessment carried out during the previous year, 18 (56%) between two and four years previously, and four (13%) more than four years ago. No patient had had haematological tests to screen for the development of pernicious anaemia. This assessment showed that our follow up of patients was inadequate and indicated the need to monitor and review patients with hypothyroidism.—W. HALL, general practitioner, Settle, North Yorkshire BD24 4JA and J. A. WALKER, Evaluation and Planning Centre, London School of Hygiene and Tropical Medicine, London WC1E 7HT. (Accepted 13 February 1986)

1. Hall AJ. Thyroxine replacement treatment: clinical judgement or biochemical control. *Br Med J* 1985; 290: 123-4.
2. Himmelfarb BJ. The importance of diagnostic screens in thyroid disease and the laboratory. *Br Med J* 1985; 290: 124-5.

100 YEARS AGO

Some interesting and graphic notes of the work being done by the National Society for the Sick and Wounded in War, in Belgrade, called from letters sent home by General Laure and Mr. Kenneth Barrington have been furnished for publication by Lord Wintage.

General Laure writes: That it was quite evident that proper work could only be done by undertaking a hospital for ourselves and in our own name. This he has done. He has caused the building given over to him to be altered so as to allow it to be worked to the best advantage, and has retained the Serbian staff of nurses; they are for the most part Serbian ladies, who have offered their services gratuitously. The Government staff is also retained in our hospital and made comfortable. He has added material to the hospital of the patients, as well as provided better clothing for them. This will have a capital effect in removing the prevalent idea in so far as our friends to the Serbian nation. From Sofia Mr. Kenneth Barrington writes that he spent December 7th and 8th in visiting the various hospitals, and making the acquaintance of the authorities. At the present moment there are 24 hospitals in the town of Sofia, of which 12 are Government hospitals, and 12 are mainly supported by voluntary agencies. In some cases, the Government supplies food and furniture, while a Red Cross Society supplies the medical staff and some moving spirit. The last society has done excellent work in the military hospital and lunatic asylum (115 wounded), the Etoile Militaire (about 270 wounded), and the Assemblée Nationale. The surgeons in several of the Government hospitals are selected from foreign ambulances. The principal voluntary agencies are the Venus Red Cross, the Dardanelles Red Cross, the Bulgarian Red Cross of which the Bishop of Sofia is president, and the Societe Internationale, of which Mr. and Mrs. Lancelotti are the moving spirit. The last society has done excellent work with the credit supplied by the English National Aid Society. With their small funds this society has supported four small hospitals, containing in all 90 to 100 patients. Of these four, one established in the Railway Administration Offices (25 beds), one paid nurse and servant, the rest volunteers, the second was formed by a resident Italian officer in his own house, his daughters being the nurses, and at one moment having the care of forty wounded, now about twenty-five. The expenses of keeping up this hospital have been borne by the Comité Internationale. The third is taken over from the "hospital of the wounded" and is the place where the untrained in hospital work. The fourth is established in the convent, the nurses being sisters of charity. These four hospitals have been in full work under the medical charge of the Italian and Bulgarian volunteer doctors

since November 22nd. Their funds consist mainly of the amounts received through Mr. Lancelotti from his credit at Courts and Co. They cannot be continued without our support, so I propose to keep them going for another month at least at a cost of about £1300 to £1500 a month. A Belgian engineer of great experience in this country looks after the economic arrangements and supplies, the whole is worked on an inexpensive but effective system. The wounded in them, as in the other hospitals, look for the most part contented, and even cheerful. Of course, many of the hospitals do not come up to our ideas of cleanliness. However, making all allowances, and remembering the fact that over 4 000 wounded were, in the course of a day or two, thrown upon the hospital, both Dr. Frerksenhaugh, Miss Stewart, and Mr. Barrington consider that the Bulgarians deserve the greatest credit for having met the emergency so promptly and effectively. The wounded Servians are, as if anything, made pets of, and a kind word is said to them by every passer-by. One brave Serbian captain, who defended the standard of his regiment until he fell with five wounds, is a special favourite. Close by are lying, side by side, a wounded Serbian and wounded Bulgarian. They had a hand-to-hand fight in the Svinica trenches, and, after wounding each other severely, helped each other along to the Bulgarian field ambulance. Among the wounded are several Turks. Many of the wounded have to use their own great coats for coverings in bed. It is difficult to keep the wards perfectly clean under these circumstances. The patients are provided with blankets here. The bales which are sent out will come in very useful. The Bulgarian Red Cross Society has been formed by a grant from the Government, and by subscriptions from abroad. The President is the Bishop of Sofia, and the Council is composed of Bulgarian residents in Sofia. In order to systematise the relief offered by the Red Cross and other Societies, the Government has appointed a special Commission, of which Mr. Grekoff, the late Prime Minister, is President, and Dr. Volkovitch, Chief Surgeon of the Army, is an important member. Mr. Barrington had a consultation with these gentlemen, and probably the result will be that Mr. Frerksenhaugh will be given medical charge of a ward in a Government hospital, with special serious cases will be placed. Miss Stewart, with some English-speaking Bulgarian nurses, will be placed in a nursing, and attend the heaviest cases. All will be done with the entire cooperation and sanction of the British Consul General, Captain L. Umley, with the instruments, arrived several days before Mr. Barrington. The instruments were all distributed by Mr. Lancelotti as presents from the National Aid Society, and receipts have been given, which will be duly forwarded. (*British Medical Journal* 1886; 2: 35.)