PRACTICE OBSERVED

Practice Research

Terminal care at home: perspective from general practice

ANDREW HAINES, ANGELA BOOROFF

Abstract

A survey of general practitioners in north west London was undertaken by questionnaire to elicit information about problems that they had had in looking after patients at home who were terminally ill and about their perceived needs for both training and support services. The response rate was 73% (196 of 268 doctors). Thirty two per cent of respondents frequently or always had problems in controlling pain in such patients, and 45% frequently or always had difficulties in coping with the emotional distress of patients or relatives, or both. Between 20 and 39% of respondents often had problems with inadequate support services, poor communication with support services and hospital specialists, and difficulty in admitting patients who were terminally ill.

Roughly half of the respondents thought that more training in managing pain and other symptoms that are associated with terminal illness would be of great help, and a similar response was noted for bereavement counselling. About 40% of respondents indicated that training in communicating with dying patients would be a great help in improving care and three quarters that more home nursing support was necessary.

Introduction

Over the past 30 years the proportion of patients who die at home has fallen from 50% to 25%.¹ In a practice with a list of 2500 patients about six patients a year may require terminal care at home. Having so few patients to look after may result in general practitioners losing their skills in managing terminal illness, together with advances in the care of terminally ill patients in recent years, may lead to a lack of confidence in their ability to manage patients at home. Several reports of terminal care in the community from the perspective of general practice have been published, mainly by practitioners with an interest in this.²⁻⁵ A study from Belfast of general practitioners and carers who looked after 118 patients with terminal illness at home documented problems that they had had. There is, however, little information about the difficulties that general practitioners often experience when looking after terminally ill patients at home and about steps which might be taken to alleviate these difficulties.

We report the results of a questionnaire survey of general practitioners in north west London which was intended to elicit information about such problems and about their perceived needs for both training and support services.

Methods

A questionnaire and a prepaid envelope were mailed to 270 general practitioners who were on the list of Brent and Harrow Family Practitioner Committee and had practice addresses in the boroughs of Brent and Harrow. Two general practitioners had retired. One remainder was sent to those who did not respond initially, and 196 (73%) of 268 general practitioners replied. In addition, two other general practitioners returned the questionnaire uncompleted and indicated that they had insufficient experience of terminal care to answer the questions. One had just entered the practice, and the other was working part time. Some questions were uncompleted, accounting for the slight variation in the denominator in the tables.

The general practitioners were asked the number of terminally ill patients

Department of General Practice, St Mary's Hospital Medical School, London. ANDREW HAINES, MD, MRCP, senior lecturer. ANGELA BOOROFF, research assistant. Correspondence to: Dr A. Haines, Horace Jowles Hall, Central Middlesex Hospital, London NW10 7NS.

1052

who had been cared for at home in the past year, the proportion who, in their opinion, wished to die at home, and the proportion who actually died at home. For group practices where a patient may have been looked after by more than one partner respondents were asked to include only patients for whom they had taken major responsibility. The options in answering the latter two questions were: none, around 25%, around 50%, around 75%, all. Further questions included 10 possible problems in the care of terminally ill patients at home. The question was worded: "In your experience which of the following cause problems in looking after terminally ill patients at home?" Respondents were asked to indicate how helpful extra training and more support might be.

Respondents were also asked: "If somewhat more resources could be provided would this enable you to look after any more terminally ill patients at home?" The possible responses were: no, somewhat more patients, considerably more patients. Finally, they were asked how information should be provided: written material, lectures, small groups of general practitioners, or films and video tapes.

The year of graduation and the use of deputising services were also recorded on the questionnaire. The use of a deputising service was considered relevant because of the implications for the continuity of care for the patients who were terminally ill and their families.

Results

Over 90% of respondents had looked after at least one patient who was terminally ill in the past year; 16 (6%) had not looked after any; 95 (48%) had looked after one to three; 56 (29%) had looked after four to six; 16 (8%) had looked after seven to nine; and 13 (7%) had looked after 10. Sixty six per cent (129 of 194 respondents) believed that half or more terminally ill patients wished to die at home, but only 41% (80/194) stated that half or more actually died at home (table 1).

TABLE 1—Answers from respondents on deaths at home (number, % of respondents)*

	None	Around 25%	Around 50%	Around 75%	All
Patients wishing to die at home	10 (5)	51 (26)	57 (29)	63 (32)	9 (5)
Patients who actually died at home	21 (11)	61 (31)	46 (24)	54 (28)	17 (9)

* For partnerships only the doctor who took responsibility for the patient is included.

Over 90% of respondents had problems at least occasionally with control of pain, incontinence, other symptoms, and the emotional distress of patients or relatives, or both (table 2). More importantly, a third indicated that they experienced problems with pain control frequently or always, more had problems controlling other symptoms either frequently or always, and

TABLE 2—Problems general practitioners experienced in looking after terminally ill patients and problems which users (number, % of respondents)

	Never	Occasionally	Frequently	In all cases
Control of pain	13 (7)	119 (60)	96 (50)	4 (2)
Control of incontinence	12 (6)	129 (67)	50 (26)	2 (1)
Control of other symptoms (dyspnoea, anorexia, confusion, etc.)	6 (3)	108 (55)	79 (40)	3 (1)
Coping with emotional distress of patients and relatives	12 (6)	96 (49)	73 (37)	14 (7)
Coping with your own emotional responses to death and dying	49 (25)	85 (44)	33 (17)	7 (4)
Inadequate support services (nursing, home help, etc.)	45 (23)	91 (47)	53 (27)	6 (3)
Poor communication with support services	56 (29)	88 (45)	46 (24)	4 (2)
Poor communication with hospital specialists	53 (27)	100 (51)	37 (19)	4 (2)
Difficulty in admitting patients to hospital or hospice to give final care or for control of symptoms	49 (25)	99 (51)	40 (21)	6 (3)
Lack of availability of appropriate facilities from local pharmacies (particularly controlled drugs)	119 (61)	71 (36)	5 (3)	1 (0.5)

just under half had difficulties in coping with emotional distress of patients or relatives, or both, frequently or always, although only a fifth had difficulties coping with their own emotional responses to death and dying frequently or always. Overall, 124 (63%) of 196 general practitioners frequently experienced one or more difficulties in patient management. Just under a third reported that support services were inadequate in most or all cases, and a slightly smaller proportion reported poor communication with services, a fifth often had poor communication with hospital specialists (table 3).

Responses to questions about the need for further training were fairly consistent (table 4). Roughly half the respondents felt that further training

TABLE 3—How helpful would the following be in helping you to provide improved care to terminally ill patients at home? (number, % of respondents)

	No help	Some help	Very helpful	Essential
In patient management	21 (11)	79 (41)	69 (36)	24 (12)
In control of other symptoms (dyspnoea, anorexia, etc.)	18 (9)	80 (42)	79 (41)	14 (7)
In communicating with dying patients	24 (12)	67 (35)	62 (32)	19 (10)
In bereavement counselling	20 (10)	77 (40)	60 (31)	20 (10)
More general nursing support	6 (3)	89 (46)	100 (51)	58 (30)
Support from home nurses with special training in terminal care	8 (4)	28 (15)	130 (68)	45 (24)
Specialist district nursing/consultative medical services for terminally ill patients	22 (11)	62 (32)	96 (49)	13 (7)

in symptom relief, communication with dying patients, and bereavement counselling would be helpful or essential to improve the care that they provided for terminally ill patients at home. Respondents were emphatic in indicating that more general nursing support and support from home nurses experienced or training requirements. There was a slight but consistent response for general practitioners who had looked after four or more patients with terminal illness in the past year to have fewer problems with pain control, control of other symptoms (but not incontinence), coping with emotional distress of patients or relatives, and coping with their own emotional responses to death and dying compared with those who had looked after fewer than four patients. The differences, however, were not significant. Those who had looked after fewer than four patients were slightly more likely than the others to think that more training in the four specified areas (pain management, other symptoms, communicating with dying patients, and bereavement counselling) could be helpful or essential for improving care, but the differences were not significant.

Several respondents thought that there was sometimes a discrepancy between the wishes of terminally ill patients, who frequently wanted to die at home, and the wishes of relatives, who were more likely to prefer for hospital admission. More support for relatives was advocated.

Only 15 (7.7%) of the respondents never used a deputising service, a further 31 (15.8%) used one only during holidays or illness, 27 (14.8%) regularly used one in the evenings, 101 (51.5%) regularly used one at night (11 pm-7 am), and 21 (10.7%) sometimes used one at night. Several respondents said that they gave relatives of terminally ill patients their home telephone numbers.

Discussion

The results of this survey suggest that general practitioners commonly experience difficulty in certain aspects of care for

patients with terminal illness. Pain is a prominent symptom for such patients, and the finding that a better response to analgesia often had problems with pain control suggests that further training is required. It is likely that the general practitioners underestimated the amount of pain and other symptoms suffered by patients. Nevertheless, nearly half indicated that they would find training in control of pain and other symptoms helpful or essential in improving the quality of home care. The experience of the Edinburgh domiciliary terminal care team suggested that complete eradication of pain could be obtained in about 90% of men and 70% of women, with good relief in a further 7% of men and 20% of women.⁶

A desire for more support from community nurses, particularly those with training in terminal care, was frequently expressed. In general, however, problems with support services were experienced less often than problems with control of symptoms. In one of the health districts (Brent's) a small team of terminal care nurses had been disbanded before the survey, but it has recently been reconstituted. This team had apparently been relatively isolated and had not been seen as part of a comprehensive community terminal care service. Hospice care is available in the North West Thames region at Michael Sobell House, Mount Vernon Hospital in Northwood, but the hospice receives only a few of its patients from the areas covered by this survey. Most terminally ill patients who require admission are therefore referred to the two district general hospitals (Central Middlesex Hospital in Brent and Northwick Park Hospital in Harrow).

The general practitioners generally seemed accessible to their patients out of surgery hours, with only a few regularly using a deputising service in the evening and at weekends. Community care might be disrupted with those general practitioners who regularly use the deputising service at night (around half), but the deputising service can be instructed to contact the general practitioner personally for patients who are terminally ill. General practitioners were aware of the difficulties in communicating with dying patients and their relatives and in handling emotional

100 YEARS AGO

The condition of the sleeping-room at night calls for attention during winter, the two chief points being the regulation of heat and of ventilation. Much depends upon the size and position of the room, the number, age, and constitutional condition of the occupants, and the actual state of the weather. The indications are, of course, to maintain a reasonably warm, uniform, and pure atmosphere. Probably most people can sleep with comfort at the temperature of the room sunk to 40° Fahr., and possibly this is rather low a limit for the majority. If the morning temperature falls to this point, some means of artificial warming of the sleeping-room is necessary. The convenient or more suitable than a small fire in an open grate, so arranged that the heat shall be as even as possible, and that it shall die out gradually and not suddenly. Here again the state of health and sensibility of the individual must be our guide. To a person of robust health, who has plentiful sources of heat within himself, and who does not know what it is to wake feeling chilly, artificial warmth of any kind is unnecessary, and should, therefore, be avoided. On the other hand, a thin-blooded, anemic, person, especially if prone to bronchial catarrh, should sleep in a moderately warm and equable temperature, however this may be best secured.

As regards ventilation, it is hardly any longer necessary to insist upon the necessity for pure air, both by night and by day. In summer it is a good practice, for those in health, to sleep with open windows (draughts being guarded against), but this wholesome practice should not be persisted in during the winter season, if experience show that it is not readily tolerated. A fall of temperature of 10° during the night is not uncommon in our climate, and would be dangerous to many persons if sleeping with open windows. The bronchitic especially must be watchful in this regard, while on the other hand nervous persons and exhausted brain workers, if free from all tendency to pulmonary catarrh, may justifiably risk a little cold of the pure air.

An impression prevails in many quarters that there is something actually noxious about night-air, *per se*, an idea for which there is no adequate

distress, although only 20% had frequent difficulties with their own emotional responses to death and dying. The results suggested that those who had looked after few terminally ill patients in the past year had more difficulties and saw training as being of more potential benefit than the others, but the differences were small and not significant.

Our results suggest that more patients could be looked after in the community and that standards of care could be considerably improved with further training and support services. The exact contribution can be measured only by carefully assessing the effects of innovations in providing terminal care at home. Some schemes have apparently been used extensively by general practitioners, others have found it harder to gain acceptance in the community.⁷⁻⁹ The difficulties experienced by the doctors seem largely to reflect those experienced by patients. The generally positive response of general practitioners to the questionnaire suggests that they consider terminal care an important part of their work and wish to improve their effectiveness.

We thank the general practitioners who participated in the survey, Mr Judith Diamond for help with data processing, Dr Carlos Doria for help in planning the study, Drs J. Korman and R. J. Jackson for helpful comments, and Ms Sheila Forman for typing the manuscript.

References

1. Fry J. Death and dying. *Lancet* 1981;277:176-7.
2. Dore D. Domiciliary terminal care. *Practitioner* 1982;228:173-82.
3. Butler PR. Care of the dying at home. *J R Coll Gen Pract* 1984;34:44-8.
4. Lambert L. Care of the dying at home. *J R Coll Gen Pract* 1972;22:468-7.
5. Calverly PR. Care of the dying at home. *J R Coll Gen Pract* 1972;22:468-7.
6. Finlay J. Care of the dying at home. *Br Med J* 1982;295:1201-2.
7. Raftery PH. Home care of the terminally ill. *J R Coll Gen Pract* 1981;31:511-7.
8. Bann T, Clarke DC, How ARL, Lamb PR. The St Thomas' Hospital terminal care support team. *Br Med J* 1981;1:1261-3.
9. Cartwright A, Hetherly L, Anderson J. *Life After Death*. London: Routledge and Kegan Paul, 1973.

Accepted 17 February 1986

1054

Audit Report

Reimmunisation of teenagers in a central London practice

The Department of Health and Social Security's guidelines, *Immunisation Against Infectious Diseases*, recommends giving reinforcing doses of tetanus and diphtheria vaccines to school children. Our local health authority these are offered to school leavers, but it was difficult to discover which schools offered reimmunisation, and when, because practitioners registered with our practice attended schools in three districts. At the end of 1983 and 1984 all registered teenagers aged between 14½ and 15½ were identified from the computerised age-sex register and offered reimmunisation. Half were written to directly, and a letter was sent to the parents of the others. In 1984 none responded were written to again and also asked whether they had been reimmunised elsewhere. Of 278 teenagers, 32% responded to the first request, but a total of 59% of the girls and 46% of the boys responded to both letters. There was moderately satisfactory success roughly 10% of our patients move away each year. Girls were more compliant than boys ($\chi^2 = 3.99$; $P < 0.05$). We were informed of only 13 teenagers who had been reimmunised before our study began.

Both general practitioners and district health authorities provide

preventive services, often duplicating each other's work and doing poorly the tasks that each considers to be the other's responsibility. Better preventive services could be delivered in general practice if primary care teams accepted that surveillance of vulnerable groups and preventing ill health are as important as treating the sick. This also seems to be the view of the Royal College of General Practitioners. Vulnerable groups include those with chronic illness, computer. More patients might carry a card recording preventive procedures which may be used as a "shared care card" by general practitioners, clinical medical officers, and school doctors, as in the hospital and community maternity services. I thank our practice nurses Liz Metcalfe and Rose Price for help and Merlin Amable for typing the paper. MICHAEL MODELL, general practitioner, James Wigg Practice, Kenilworth Town Health Centre, London NW5. (Accepted 8 April 1986)

1. Joint Committee on Vaccination and Immunisation. *Immunisation against infectious diseases*. London: Department of Health and Social Security, 1984.

2. Working Party. *Royal College of General Practitioners. Health care—dualing prevention*. London: RCGP, 1982.

100 YEARS AGO

As a proof of the old adage, that "the wish is father to the thought," we are pleasantly startled now and again with the announcement, in the newspapers, that a cure has been discovered for that fell disease, the Saxon name, consumption. This announcement comes round with about the same regularity as that of the famous sea-serpent, whose remains, if the animal could only be caught, would meet with a cordial welcome in our museums of comparative anatomy. These alleged remedies generally reflect, to some limited extent, the then current views as to the pathology of this terrible affection, and they thus secure for themselves a share of the popularity which almost invariably surrounds the last new hypothesis, until by-and-by they are shown to be wrong. During the past week, however, we have been treated to a more novel phase of this tendency, and one possibly more open to criticism than the majority of such effusions which interest and may even amuse, while they can scarcely do any harm. Within a few days of the appearance in a lay contemporary of a sensational article on the reported good results obtained in one of two cases of phthisis by the inhalation of the spores of the "bacillus terminus" by an Italian physician, a letter was published in the same journal from a medical gentleman, named Lambert, residing in Liverpool, in which he claims to have effected—rather partly effected—the cure of a very advanced case of phthisis. So eager was the benefactor to give the world the benefit of his experience, that, unwilling to lose valuable time in checking the results by even one other observation, or even in completing the cure, he had his hand, he confesses, not only in his own but in an enthusiastic and glowing account of the wonders he has achieved. That his conduct is irregular and open to censure, however, is hardly

admits, but he conceives his independence to the writer of the article on the murder held recently at Rodean, in the south of France, where a self-accused was proved to be innocent, and the true criminal detected by medical evidence. Last August, a woman, named Melrose, went to register the death of a child, which she had wrapped up in her apron. The registrar examined the child, as is customary, and observed finger-marks on its wrist. The mother then declared that she had strangled her infant. She was sent to prison, but Dr. Deamont, who had been directed to examine the body of the child, stated that it had certainly been strangled, but not by its mother. Her hand was more delicately formed than that of the murderer, which had left its impression on the child's neck. The fingers of the guilty party must have been short and thick, the index-finger being unusually short, and apparently devoid of a nail, defects which the writers indicated as valuable clues for the discovery of the murderer. Melrose's family admitted that the murderer was a man called Bonnet, and that she was his servant. The accused was arrested, and his index-finger was found to be one centimetre shorter than the average length of that member, and its nail had been destroyed by accident or disease. Bonnet was sentenced to six years' imprisonment. (*British Medical Journal* 1886;2:81.)

Our Paris correspondent forwards particulars of a remarkable trial for murder held recently at Rodean, in the south of France, where a self-accused was proved to be innocent, and the true criminal detected by medical evidence. Last August, a woman, named Melrose, went to register the death of a child, which she had wrapped up in her apron. The registrar examined the child, as is customary, and observed finger-marks on its wrist. The mother then declared that she had strangled her infant. She was sent to prison, but Dr. Deamont, who had been directed to examine the body of the child, stated that it had certainly been strangled, but not by its mother. Her hand was more delicately formed than that of the murderer, which had left its impression on the child's neck. The fingers of the guilty party must have been short and thick, the index-finger being unusually short, and apparently devoid of a nail, defects which the writers indicated as valuable clues for the discovery of the murderer. Melrose's family admitted that the murderer was a man called Bonnet, and that she was his servant. The accused was arrested, and his index-finger was found to be one centimetre shorter than the average length of that member, and its nail had been destroyed by accident or disease. Bonnet was sentenced to six years' imprisonment. (*British Medical Journal* 1886;2:81.)