PRACTICE OBSERVED

Practice Research

General practitioners' retirement plans and what influences them

RICHARD WAKEFORD, MARIAN RODEN, ARTHUR ROTHMAN

Abstract: The results of a study of the attitudes of 197 general practitioners aged 55 and over towards retirement and their plans for retirement are reported. Few wished to retire very early, and only a further 40% definitely planned a clean break from practice. Nearly half planned on taking "14% hour retirement," even though most (78%) will be entitled to full pension rights. Job satisfaction and health were the most important factors influencing the timing of the returement. Severaty seven per cent thought that there should be no formal retirement policy. 79% wanted no compulsory retirement age, and 80% wanted as further controls or safeguards directed at older practising general practitioners as safeguards directed at older practising general practitioners for the processing of the practice of the processing practice of the processing that the practice of the practising general practitioners. Presonal discussions with respondent sage 465 and over Personal discussions with respondents aged 55 and over

and the general practionness in the sample ever not tooking. Personal discussions with respondents aged 65 and over aggressed that there are many older graeral practitioners who has to practice to some time, whether that they will know when the contraction of the contraction of

Office of the Regius Professor of Physic, Cambridge University School of Clinical Medicine, Cambridge RICHARD WASHERDE MA, sensor receipt hassis at RICHARD WASHERDE MA, sensor receipt hassis at ARTHER ROTTHMAN, 15th, visualing fellow

Gerrespondence to: Mr R E Wakeford, Clinical School Offices, Addenbrooke's Hospital, Cambridge CB2 2QQ

Introduction

Whether general practitioners retire early, late, or at an age that is regarded as conventional 65 years has important implications for manpower planning—for example, in connection with the demand for and establishment of sociational training schemes. With the exception of a survey that was limited to inner London, little appears to have been established about general practitioners' retirement plans and the factors that influence these. Indeed, the only relevant study of retirement intentions and related attitudes of which we know was restricted to anaesthetists in Yorkshire. Older medical practitioners in North Amerias were found to be less successful on mesures of clinical performance. and there are to participate in continuing medical education. Coupled with the fact that 29% of practising general practitioners in the UK aged 65 and over a ret in singlehanded practice, compared with 14% overall. Department of Health and Social Security. Statistics and Research Section, personal communication, 1984 and this not receiving daily informal "peer review." this max cause concern about older general practitioners in the UK. It would therefore be useful to know about the retirement plans of these dictors and what influences these plans.

Astudy was undertakenering conventional retirement age or who had passed this personal continuing and the processing of the prevention influence them towards earlier of later retirement. Then the retirement intentions of a national sample of such general practitioners were examined.

Conducting personal interviews with a national sample of respondentials to face to experience, vet postal questionnaire did not seem appropriate for collecting data on such a personal subject, and a poor response rate could be expected. A compromise method where eighbon enterwise. Though the expected A compromise method where eighbon enterwises. Though the discussing them in depth, it has been found reliable in surveying broadly the sees and attributed of general practitioners. We therefore decided in collect primary data by means of automat (elephone interviews word) and automat (elephone interviews word) and admitted a simple of sheal discuss where we have the admitted and applications. The national sample of discuss was drawn from the Madual Discussion of the assumption that the age of discuss at qualifiations of 3. The directions on the assumption that the age of discuss at qualifiations (3.2). The directions excluded from the studie are portained subsects who qualified much later practitioner practitioner practioning in Britain. Sor older, was selected for the study. Exclusive below these found in relighbours are small these were replaced by the control of the control of the production of the study. The control of the production of the study of the production of the production of the study of the production of the study. The production of the production of the discussion of the discussion

Results

Results

Of the 250 doctors in the national sample, 28 were no longer practising counting in criterium to death, and a further 24, who were undervised still to be in practice, could not be reached for interview often because of association with the interview of the December of the Control o

Activity	"+ Of general peacutewisers"
Industrial medicine	45
Private practice	17
Nursing Nome et.	(4
Point surgeon	18
Medical training	Į0
Hospital sessions	7
Research	0
Other:	28
None	ta ta

*More than one response was allowed.

This today, for example, medical examinations for insurance companies and the DHSS and
medical services for souris (earns, boarding schools, and homes for hands apped people.)

BRITISH	MEDICAL	JOURNAL	VOLUME	292	17 MAY	198
·· 18. 4 ··				211		

	"« Of general peachthoner»		
l-urnal-	*		
Meetings and courses at punigraduate centres	x i		
Hospital consultants	in		
Likal medical success	4		
Intraction meetings			
Merrings with other practices			
Other	11		

reportedly obtain their continuing medical education Journals and post-graduate centres were most frequently methodied, but meetings with a final continuing the second of the properties of the Result College of termined Participients (4%). The Laurel 25% of Tournal of the Result College of termined Participients (4%), solds and sectories 5% of and their excellences with a felf and Palie 41%. Retirement plane 10% and palie 41% of the properties of the propert

Most general practitioners in the sample. 78% is with be entitled to full persons on retriement and most 78% has perchased additional vales to ready frid persons. The area reflection and with register and produced to the produced and with right influence and with right influence them to retire taker rather than earlier. Also, if 4% of the respondence would offer no reasons for each other stems on the stems of the stems of the respondence. Lack of she windstorn would leaf no early retirement for the stems. 18% for each term, tables III and IV. Job satisfaction was the dominant influence. Lack of she windstorn would leaf no early retirement of most over half. Flow health was the excend fact the would retirement for out-over half. Flow health was the excend fact the would formshot to early retirement, and then being needed by patients and financial reasons would contribute to delaying retirement. Several disknown expressed citizene rolls, time to leave peached intend on interest but medium when adde alread forward to return medium. We could not say, 42% were fooking forward to returnment more than those who were 55-60 to those over 55.

Vieta on retirement policies.— Most of the general practitioners in the sample.

TABLE 111- "What do you see as being the major factors which would influence you to

	"- Of general peaculastics
Lack of soft satisfaction	4.2
Pour health	19
Not keeping abreast not providing good service	11
Lo pursue other interests	
Family	2
Changes in the law after ting pensions retirement	i
Other	4
No amount	4

	"> Of general peachtisoner
let stole in o	**
Needed by patients	15
More	13
Encouraged to stay by partners	*
Health	4
turning grand service	
Souther interests	1
hher	2
No answer	ii.

BRITISH MEDICAL JOURNAL VOLUME 292 17 MAY 1986

BRITISH MEDICAL JOURNAL. VOLUME 292. 17 MAY 1986

beld vews concerning formal retirement policies consistent with their position as self-employed professionals. 77% and that there should not be a reduced to the policy of the policy of the policy of the policy of the decision were less are about the that there delect obligators. 77% wasted no compulsion retirement age, and 87% wanted no further formal controls, steparate of the policy of the polic

Classification of retirement intension see lext	Present age			
	n NI	61.65	nn ts	respondents n 189
Want out wen	1	6	0	
Clean break	4n	42		tic
Gradual	49	42	57	490
Sever	4	10	4	- 11

Discussion

For almost all general practitioners, general practice is a lifetime career, almost always spent in the same place. For many, their only medical activity consists of providing care for the patients on their MISE list, many abor device their whole life to their practice. With list and the same place is the providing care for the patients on their MISE list, many aborded and sensitive states of the same place place of the same place of the same place of the same place place of the same place pla

(DHSS (SRIA), General medical services some base statistics, 1984). If women's unreations of retirement are different from men's they will not be referred in this study.

Although the need for income and the possibility of poor health were important in influencing general practitioners to think of retiring later or earlier (respectively), job sarisfaction or lack of in had the most influence. Thus many wished to "die in harness" and quoted their perceptions of their patients urging them not to retire they were horrified at the prospect of compulsory retirement. It is of course open to discussion how accurate these perceptions more decision of ware of the prospect of compulsory retirement. It is of course open to discussion how accurate these perceptions more decision of the prospect of compulsory retirement. It is of course they have been also also decision of the prospect of the patient oriented—particularly when nearly half of our respondents reported that they did not look forward to retirement. Whether it is in the interests of patients that their general practitioners were asked about the didition of the prospective of the prospec

This study was funded by a grant from the King Edward's Hospital Fund for London. We thank all the dectors who participated in the study, the DHSS Statistics and Research Division (SRTA) for providing statistical information, and Professor Gord Norman for statistical advice.

Emilia Bladin Blading Levertion, Primer health air is mer finder. Leukar EMR (2011).

2 Mills Will Scrimmen of autochronic attailed in Vestilland. Section 1998; 50–50.

3 Mills Scrimmen of Section 1999; 50–50.

3 Mills Scrimmen of Blading Scrimmen or consenting primer in the slight of Blading in Mills Scrimmen of Section 1999; 50–50.

3 Mills Scrimmen of Blading Scrimmen or Section 1999; 50–50.

5 Mills Scrimmen of Section 1999; 50–50.

5 Mills S

Appendix

Views of De, A and De Romentmens.

De A is Tan Bose to mich same there partner rural practice to 80 years. The practice provides its own is used bours over but uses a locum one; a week. His practice provides its own is used bours over but uses a locum one; a week. His practice provides and assimptions returned age. Patients should be the ones to deside whether ones continued. A storage of the search would be the ones to deside whether ones continued. A storage is the search would be the ones to deside whether ones of the storage. A storage is the search would be the ones to deside whether ones of the storage is the search of the storage of the storage is the storage of the storage of the storage of the storage is the storage of the storage of the storage of the storage is the storage of the

not soking forward to returned to the period of the form of the fo

BRITISH MEDICAL JOURNAL VOLUME 292 17 MAY 1986

Questioning Practice

Chickenpox isolation: is it worth the trouble?

We received the following letter from De F. C. Katalite and whed they control practitioners and a community parliamy to unities his question. It is obtained while the his hopes necessary:

Children who have, bukenpox should be isolated for one week from the appearance of spots or until dry scabs have formed, the text-took rell us. Some schools will not allow shidren back until all their spots have disriperated But is resistant on children with shickenpox necessars. And is it effective. "Chickenpox is highly contagnous. Over 85% of susceptible household contacts will cach the disease. 9% of American women contacts and the shickenpox necessars." It is, however, a clustered mild disease, and lives shidren develop serious complications. The idea behind toolaing, shidren with this kenjor will be disease. The short period of interview feeling the proper of their form activing the disease. The short period of interview before the appearance of the risks, however, it offers out prevent others from activing the disease. The short period of interview before the appearance of the risks, however, it offers out in the shid with his high with his classmates before he goes into isolation. So what is being gained by our attempts to contain the endemandeese Not much, I suspect. Plents of well children with spirs are enforted and the shid of t

Retinum RE - Guidher ST - ols - Nove violente et pudous. 12th (ol. Polidelphi) (S. B. Nordale, 1987-25.

Nordale Brit of the State of the British of the State of

De Goree R Lector—Dr Kavalier presents a reasonable argument, with which I am sure many peatimenes agree. The rodation of shalden with an illness that is southly recognised only when the rach appears, set in most infections on the prodomal vage, is noneme. The fast that shickenpors is contracted by a regernal woman means that all children booked be exposed to the virus. The real problem is that society has an aversion to people with spear and an error practice onto children with right, sides until which the proposed to the virus. The real problem is that society has an aversion to people with spear and on the practice onto children with right, sides until which who have reportability for children. Ethocating a section is the medical profession mass above needed.

It a propulation of children can grow up happs in the knowledge that it is better to have child expose a unit and up age and that it is not across in terminal profession mass above needed.

educate the staff. This should also attempt to make it local police that sports shidden come to should their account plantationer should also privace three discharged makes the staff and the staff in the should also private three discharged or when asked to talk about intertious discusse to groups—timel Post Health Center. Surchimmerization Design Staff and the Post Health Center. Surchimmerization Design Staff and the staff and the

Kibel M. (widomin) | Mile A. (n. instal.). Amen't in Translation paradiams. Translat. Condenses the child in congress. 1975; 1940; 4.

De Staart Carne - For most of its outmoschickenpox is a mild dosess. It is infectious, being spread mainly by the respirators staat. The degree of infectitivity, however, is not is great as for example, in measles, and about his control of the production of the pr

Ib Notice Adaptive—It is difficult to due the origin of evolution the individual-white the keeping from the rest of the community appears in which is had solution when the found it was into our technole. It is recombile to assume that the National physicians considered it necessary to indiate the person with shickingoin topical order of the National Policy of the N

1 McGregor Robertson J. Honochold physician. London: Gredham, 1894. 2 Kragman N. Katz NJ. Julic non-disease or studies. NJ Jugis. Modes, 1981.