metabolic needs or energy sparing mechanisms they were left with only one other possibility: the underestimation of food intake. They concluded that the obese people must have been consuming an average of 530 kcal a day more than the lean controls in order to sustain their obesity. The cause of the obesity must, therefore, be sought in the control of food intake and eating behaviour.

This raises several questions. Is the deception deliberate? In this study did the obese misread the scales when they weighed their food or did they "forget" to weigh and record some foods? The size of the postulated underestimation appears too great to countenance accidental omission—it is equivalent to nearly a third of all the food they did record. Apparently these obese women were unable to admit to the experimenters (and probably to themselves) their actual food intake. Furthermore, the data from the lean group also included one substantial underestimation of intake; it might be difficult to prove that the lean were more consistent than the obese.

The paper also adds to the growing evidence that energy expenditures in normal life are lower than the currently accepted estimates of energy requirements.⁶ Nutritionists will need to bear this in mind if obesity is not to become more

common than it is. The central finding, however, is that this type of obese patient should no longer be able to claim inherently lower metabolic requirements. We should treat with scepticism all claims from obese patients that their dietary intakes are modest. In such people, possibly even more than in the lean ones, habitual intake is impossible to measure accurately, at least with existing techniques—a message that needs to be appreciated much more widely. The measurement of food intake is so crucial that it might be argued that research on these techniques is more important than other more prestigious topics.

D A T SOUTHGATE

Head,
Nutrition and Food Quality Division,
Agricultural and Food Research Council Food Research Institute,
Norwich NR4 7UA

- 1 Garrow JS. *Treat obesity seriously: a clinical manual*. London: Churchill Livingstone, 1981.
- 2 Royal College of Physicians. Obesity. *J R Coll Physicians Lond* 1983;17:3-58.
- 3 James WPT, Trayhurn P. Thermogenesis and obesity. *Br Med Bull* 1981;37:43-8.
- 4 Stockley L. Changes in habitual intake during weighed inventory surveys and duplication diet collections. A short review. *Ecology of Food and Nutrition* 1985;17:263-9.
- 5 Lifson N, McClintock R. Theory of the use of the turnover rates of body water for measuring energy and material balance. *J Theor Biol* 1966;12:46-74.
- 6 Department of Health and Social Security. *Recommended daily amounts of food energy and nutrients for groups of people in the United Kingdom*. London: HMSO, 1979. (Report on Health and Social Subjects No 15.)

Regular Review

Psoriasis

R H CHAMPION

Psoriasis was last reviewed in the *BMJ* five years ago,¹ and, though no major breakthroughs have been made in our understanding or treatment of the disease, many small but useful advances have been made and have influenced clinical practice. The ramifications of the disease make it of interest to both scientists and clinicians and some 1500 papers have been written about psoriasis in these five years, including some good reviews and an excellent 600 page monograph.²⁻⁶ Psoriasis remains one of the most common skin complaints, affecting about 1·5% of the population—or even up to 5% at some stage of their lives.⁷ Once patients have developed psoriasis only 40% have prolonged remissions on long follow-up.⁸ The severity may range from a minimal occasional cosmetic problem to a life threatening emergency.

The aetiology of psoriasis is multifactorial. Most of the clinical and pathological changes may be explained on the basis of the rapid turnover of the epidermis, but this does not immediately explain the other main ingredient of psoriasis— inflammation. We have come a long way from the observations of Van Scott and Ekel in 1963 that there is 27 times the normal rate of mitosis in the epidermis.⁹ Normal skin produces 1246 new cells cm² a day from a total 27 000 cells;

but psoriatic skin produces 35 000 new cells cm² a day from 52 000.^{10 11} To achieve this dramatic increase the cell cycle is reduced from 311 to 36 hours; the proliferative cell population is doubled; and the proportion of basal cells actually concerned in division is also roughly doubled.

What causes this overactivity of the epidermis is still a vexed question. A genetic factor is undisputed but variable, probably with polygenic inheritance. Our usual concept of a powerful genetic influence has to be stretched to accommodate a disease which may appear for the first time at the age of 108.¹² The genetic influence seems less important in patients whose psoriasis starts later in adult life. Other trigger factors include trauma (a useful model for studying the disease), infection, drugs (notably lithium, chloroquine, and rarely β blockers), and rather more nebulously climate and hormonal factors, and perhaps even smoking.¹³ The contribution of psychological stress is disputed. Most clinicians would agree that stress may modify the course of the disease, though seldom to the extent that some patients are led to believe, and there is only meagre scientific evidence for this.^{6 14 15}

The whole skin is potentially abnormal in patients with

psoriasis, and appropriate tests may show several changes found in greater degree in the lesions themselves. Moreover, when epidermis is grown in culture *in vitro* or as explants on the nude mouse it still maintains its increased potential deoxyribonucleic acid production.¹⁶ Rather paradoxically, an influence of the dermis is shown by the ability of cells from the psoriatic dermis to provoke hyperproliferative activity in normal keratinocytes *in vitro*.¹⁷ Why lesions are often spotty and why they have such characteristic sharp edges are mysteries which apply to many skin diseases.¹⁸

The nature of the biochemical fault underlying the rapid turnover in psoriasis is still attracting much interest. As with much research into psoriasis the work is hampered by there being no good experimental model in animals—and there is no naturally occurring psoriasis in animals. The problem is to determine which of the many measurable changes are the cause of the rapid turnover and which are the result. Most come in the latter category. The undoubted immunological changes are quite likely, too, to be secondary though perhaps perpetuating factors⁶; the same may be said for the changes in leucocyte function and other evidence that psoriasis is a systemic disease.^{6,19,20} whether psoriasis conveys any biological advantage so that it has not been eliminated by natural selection remains uncertain.

Approach to treatment

No major new therapeutic weapons have been introduced this decade, but our knowledge of established remedies has been painstakingly consolidated. There are five main therapeutic approaches:

- remove any trigger causes;
- general—explanation, reassurance, rest, holiday;
- topical—tar, dithranol, corticosteroids, keratolytics, emollients;
- ultraviolet radiation—ultraviolet B (UVB) and psoralen plus ultraviolet A (PUVA) (these differ in their action);
- oral drugs—methotrexate, hydroxyurea, and so on, and etretinate.

With each of these remedies a decision has to be made about the need for maintenance treatment or whether a short sharp course will give a prolonged remission. The momentum and inertia of psoriasis have been discussed before¹; for many patients it is indeed fortunate that a course of treatment lasting two to four weeks may achieve a remission lasting six to 12 months or even longer. For other patients it seems that long term, lower dose maintenance treatment to “keep the brakes on” the psoriatic process may cause less hazard and less hassle. There are no clear rules, and each patient has to be judged on his or her merits. A symposium on the long term management of psoriasis will be held in Helsinki later this year.

The treatment listed above has about 10 variables, and in the past few years there have been many efforts to show a genuine synergistic effect of two or more modalities. The possible permutations and combinations are enormous. Some of these date back many years and are now established regimens—for example, the Goeckerman regimen of tar and ultraviolet radiation^{21,23} and the Ingram or Leeds regimen of tar baths, ultraviolet radiation, and dithranol.²⁴

Removal of triggers is seldom very easy but should not be neglected.²⁵ Explanation and reassurance are cheap, apart from the time entailed, and without side effects. Efforts to clean up the various messy local applications have not been completely successful. Tar is still only a moderately effective remedy, but even the newer rather more elegant preparations are seldom able to be both effective and cosmetically pleasing except in mild cases. Without doubt tar may measurably potentiate the action of ultraviolet radiation. All the evidence indicates that the theoretical risk of carcinogenesis with tar is not reflected in clinical practice, even after years of use and prolonged follow up, except for a mere handful of isolated case reports.

For the common chronic plaque type of psoriasis dithranol (anthralin) remains a favourite remedy with dermatologists in Britain and is gaining ground in America. It should be used with the greatest caution in other types of the disease. The staining of the skin and clothing restricts its popularity, and it may cause inflammation, especially of the surrounding skin. The dose should be chosen with care. For many patients low concentrations are adequate (for example, 0.1% to 0.2% in Lassar's paste or in a cream base) without any incremental increases. Other patients seem to benefit from progressing to much stronger concentrations. Attempts have been made to isolate other derivatives of dithranol that may be easier to use. None of these has yet won through.

Careful studies have been reported of the mode of action of dithranol and of how it can best be used.^{26,28} The use of short term treatment is a real advance: the application is left on for a short time and then washed off, still leaving the drug in the skin, where it is needed, but not on the surface, where it makes a mess—though some may leach out again later.^{29,30} This concept was introduced on good pharmacokinetic theory—any active drug is likely to be better absorbed through the abnormally permeable, albeit thick, horny layer of lesions than through normal stratum corneum. The application of 1% or 2% dithranol, often with the addition of 5% salicylic acid, is left on for 10–20 minutes. This method may not clear psoriasis any faster than conventional treatment for 23 hours a day but it is very popular with many patients.³¹ One recent paper has recommended treatment of the whole body with low doses of dithranol cream short term.³² In theory the same principles might be applied to treatment with tar³³ and topical steroids not only in psoriasis but also in eczema and other dermatoses.

Topical steroids are still widely used for psoriasis but with diminishing enthusiasm from those who look after patients long term. The problems of atrophy and of other side effects compound the problem that steroids seem to become decreasingly effective as time goes by; they may even possibly alter the course of the disease for the worst if overused. Nevertheless, they may be valuable at times—for example, in patients with the not uncommon mixtures of psoriasis with eczema or seborrheic dermatitis and also rather bizarrely on those sites where one might expect them to be contraindicated—the face, flexures, genitalia, and hands. In theory the combination of steroids with dithranol or tar should be desirable, but the published evidence is confusing. A few patients with chronic plaque psoriasis, resistant perhaps even to strong dithranol as an inpatient, may be greatly improved by just a few nights using a really strong topical steroid such as clobetasol propionate, perhaps even under polyethylene occlusion. This seems to apply especially to the unusual rough velvety plaques which may occur over the lower legs or sacrum. We have found that such

treatment is most helpful in patients already having some other active treatment. In general dermatologists nowadays rarely prescribe strong steroids under polyethylene occlusion.

Ultraviolet

The dramatic results of treating psoriasis with PUVA have been tempered by its long term potential for producing aging of the skin, neoplasms, and other problems. The actual risks cannot be measured after only 10 years, but they cannot be discounted and seem to be greater in those who have had prolonged exposure to other carcinogens such as arsenic or radiotherapy.^{2 34 35} The many publications on PUVA show general agreement that it is a very appropriate treatment for patients with severe psoriasis, especially those over the age of 45.^{2 6 36} There is still no consensus on whether intensive courses or long term maintenance treatment expose the patient to less total radiation and whether there is a useful synergistic effect when treatment is combined with dithranol or with etretinate, both of which can be useful, or with UVB.³⁷ PUVA treatment using topical psoralens may be useful for localised areas of psoriasis—for example, on the hands. More widespread topical treatment with psoralen baths is popular in some countries, sometimes combined with other measures.³⁸

The careful dosimetry required with PUVA has led to a re-examination of the use of conventional ultraviolet radiation without psoralen. Long wave UVA, the active ingredient of PUVA, is almost inactive by itself except after very prolonged exposures.³⁹ Aggressive treatment with UVB (the sunburn wavelengths) may, however, achieve results almost comparable with those of PUVA, and much smaller doses of radiation may be effective.⁴⁰⁻⁴² Topical tar preparations seem to potentiate the action of moderate doses of UVB but not to add much to maximally aggressive doses. Topical emollients are of value applied just before radiation.²² They improve the penetration of the ultraviolet through the often opaque scales and yet may filter off some of the rays penetrating into normal skin—and it is generally the burning of the normal skin which limits treatment. A vigorous course of UVB may be comparable in its radiation with a good sunny holiday,⁴³ so that if prolonged treatment is contemplated (which it usually is not) there might be problems.

Systemic drugs

Systemic drugs come and go. Even after a quarter of a century methotrexate deserves a prime place for life threatening or life ruining psoriasis not responding to other remedies.^{2 44 45} It is very popular with patients who need consume only a few tablets once a week and can dispense with all local applications. Evidence of liver damage (cirrhosis in 5% of patients having long term treatment,⁴⁶) must still be sought by liver biopsies every year or two; scans⁴⁷ and ultrasound⁴⁸ can provide only limited reassurance, and biochemical screening by itself is inadequate. Hydroxyurea is less effective, more toxic to the bone marrow, but virtually lacks the liver problems.^{2 6} It is a useful second line drug in the not uncommon context of a patient with bad psoriasis with alcohol problems. Other cytotoxic and immunosuppressive drugs have only occasional uses. Thioguanine shows some promise.³ Benoxaprofen has yet to be replaced.

The retinoid etretinate is a useful drug, less effective than methotrexate, and with its own problems on the skin, lipids, and especially on the developing fetus. Enthusiastic reports before it was introduced into Britain led many dermatologists to hope that it would be a panacea for any patient with bad psoriasis. Now available (on hospital prescription) for some six years it has not quite lived up to these lofty aspirations but is nevertheless a welcome addition.⁴⁹⁻⁵² It may be very useful for awkward clinical variants and for plaque psoriasis—perhaps especially if combined with some other treatment such as dithranol or PUVA, where there seems to be a definite synergistic effect. Even better retinoids may be coming.

Many other drugs have been the subject of isolated or limited case reports but cannot yet be acclaimed as more than interesting—colchicine, clofazimine, cyclosporin, somatostatin, bromocriptine, and so on. Systemic corticosteroids are contraindicated in ordinary psoriasis. They may very occasionally be of benefit in life threatening pustular psoriasis or in psoriatic erythroderma not responding to methotrexate. It is my conviction that in such rare cases triamcinolone is a more effective drug than prednisolone despite the greater toxicity. Steroids are also used for an occasional case of psoriatic arthropathy, a disease which is usually somewhat milder than rheumatoid arthritis and whose pathogenesis is still obscure.^{2 6 53} It is not due to rapid turnover of synovial cells.

Other treatments

The beneficial effects of haemodialysis and of peritoneal dialysis are most interesting—but at this stage mostly for any light they throw on pathogenesis.⁵⁴ Even surgical ablation of psoriatic plaques has been advocated.⁵⁵

Should treatment be on an inpatient or outpatient basis? Throughout this review the implication has been that a short sharp two to four week course of treatment with many varied remedies may produce a prolonged remission. Many of the treatments are messy, require careful nursing technique or careful monitoring, and undoubtedly work better on inpatients, but attempts have been made to forgo this expensive approach. Day care centres have proved very effective both in Britain and abroad.^{24 56} Another useful compromise may be to have the patient in the ward for two weeks followed by intensive treatment for another week or two as an outpatient. Patients treated in this way have been reported to get better just as quickly as those who spent their whole treatment time as an inpatient.⁵⁷ Which regimen is used must often depend on the facilities available, but there can be no doubt of the importance of teaching the patients precisely how to use the remedies prescribed for them.

Patients with atypical variants of psoriasis require special types of treatment which I shall not discuss.^{2 6} Patients with atypical psoriasis, whether inflamed, flexural, pustular, admixed with eczema, or acute guttate, do not tolerate many of the remedies worked out so carefully for patients with plaque psoriasis, and if in doubt simple emollients may be best.

In a hypothetical personal series of 1000 patients with straightforward plaque psoriasis attending hospital, and to ignore any double treatments prescribed to one patient, the choice of main active ingredients might be 820 having topical treatment (tar 200, dithranol 300, corticosteroids 150, keratolytics 50, emollients 100, and others 20); 120 having

ultraviolet treatment (UVB 100, PUVA 20); and 60 having drugs (methotrexate 20, hydroxyurea 5, etretinate 25, others 10).

In conclusion, the management of a patient with severe psoriasis may cause less heartache to the physician than the management of one with stubborn, moderate disease. There are now several excellent remedies—all with their own problems, but which are worth while in severe psoriasis. It is much harder to decide whether a patient should be asked to

take a 1%, 0.1%, or even 0.01% risk of a severe side effect in order to be able to enjoy life to the full, and a long term plan has to be made for every sufferer. Let us go on hoping for an effective and safe orally active drug to alleviate a disease which may cause so much real ill health and even more anguish.

R H CHAMPION

Consultant Dermatologist,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

- 1 Champion RH. Psoriasis and its treatment. *Br Med J* 1981;282:343-6.
- 2 Baker H. Psoriasis. In: Rook AJ, Wilkinson DS, Ebling FJ, Champion RH, Burton JL, eds. *Textbook of dermatology*. 4th ed. Oxford: Blackwell (in press). (3rd ed 1979.)
- 3 Roenigk HH, Maibach HI. Psoriasis. *Seminars in Dermatology* 1985;4:271-326.
- 4 Farber EM, Cox AJ, eds. *Psoriasis—proceedings of third international symposium*. New York: Grune and Stratton, 1982. (Fourth symposium July 1986.)
- 5 Weinstein GD, Vorhees JJ. Psoriasis. *Dermatologic Clinics* 1982;2:355-519.
- 6 Roenigk HH, Maibach HI, eds. *Psoriasis*. New York: Marcel Dekker, 1985.
- 7 Kavli G, Førde OH, Arnesen E, Stenvold SE. Psoriasis: familial predisposition and environmental factors. *Br Med J* 1985;291:999-1000.
- 8 Farber EM, Nall ML. The natural history of psoriasis in 5800 patients. *Dermatologica* 1974;148:1-18.
- 9 Van Scott EJ, Ekel TM. Kinetics of hyperplasia in psoriasis. *Arch Dermatol* 1963;88:373-80.
- 10 Weinstein GD, McCullough JL. Cell proliferation kinetics. In: Roenigk HH, Maibach HI, eds. *Psoriasis*. New York: Marcel Dekker, 1985:233-47.
- 11 Wright NA, Camplejohn RS. *Psoriasis: cell proliferation*. Edinburgh: Churchill Livingstone, 1983.
- 12 Buntin DM, Skinner RB, Rosenberg EW. Onset of psoriasis at age 108. *J Am Acad Dermatol* 1983;9:276-7.
- 13 O'Doherty CJ, MacIntyre C. Palmoplantar pustulosis and smoking. *Br Med J* 1985;291:862-4.
- 14 Whitlock FA. *Psychophysiological aspects of skin disease*. London: Saunders, 1976.
- 15 Payne RA, Payne CMER, Marks R. Stress does not worsen psoriasis? A controlled study of 32 patients. *Clin Exp Dermatol* 1985;10:239-45.
- 16 Kragballe K, Desjarlais L, Marcelo CL. Increased DNA synthesis of uninvolved psoriatic epidermis is maintained in vitro. *Br J Dermatol* 1985;112:263-70.
- 17 Saig P, Coulomb B, Lebreton C, Bell E, Dubertret L. Psoriatic fibroblasts induced hyperproliferation of normal keratinocytes in a skin equivalent model in vitro. *Science* 1985;230:669-72.
- 18 Whimster IW. An experimental approach to the problem of spottiness. *Br J Dermatol* 1965;77:397-420.
- 19 Preissner WC, Schröder J-M, Christophers E. Altered polymorphonuclear leukocyte responses in psoriasis: chemotaxis and degranulation. *Br J Dermatol* 1983;109:1-8.
- 20 Geerdink JPM, de Mulder PHM, Franzen MR, et al. Monocyte function is normal in quiescent psoriasis. *J Invest Dermatol* 1984;82:122-5.
- 21 Goeckerman WH. Treatment of psoriasis. *Northwest Medicine* 1925;24:229-31.
- 22 Adrian RM, Parrish JA, Momtaz-TK, Karlin MJ. Outpatient phototherapy for psoriasis. *Arch Dermatol* 1981;117:623-6.
- 23 Le Vine MJ, White HAD, Parrish JA. Components of the Goeckerman regimen. *J Invest Dermatol* 1979;73:170-3.
- 24 Ingram JT. The significance and management of psoriasis. *Br Med J* 1954;ii:823-8.
- 25 Farber EM, Nall L. An appraisal of means to prevent and control psoriasis. *J Am Acad Dermatol* 1984;10:511-7.
- 26 Gorsulowsky DC, Vorhees JJ, Ellis CN. Anthralin therapy for psoriasis—a new look at an old compound. *Arch Dermatol* 1985;121:1509-11.
- 27 Shroot B, Schaeffer H, Juhlin L, Greaves MW, eds. Current concepts in the mode of action in anthralin in the treatment of psoriasis. *Br J Dermatol* 1981;105(suppl 20):1-114.
- 28 Molière P, Dubertret L, Melo TSE, Salet C, Fosse M, Santus R. The effect of anthralin (dithranol) on mitochondria. *Br J Dermatol* 1985;112:509-15.
- 29 Schaeffer H, Farber EM, Goldberg L, Schalla W. Limited application period for dithranol in psoriasis. *Br J Dermatol* 1980;102:571-3.
- 30 Schaeffer H. Short contact therapy. *Arch Dermatol* 1985;121:1505-9.
- 31 Statham BN, Ryatt KS, Rowell NR. Short contact dithranol therapy—a comparison with the Ingram regime. *Br J Dermatol* 1984;110:703-8.
- 32 Schwarz T, Gschnait F. Anthralin minute entire skin treatment: a new outpatient therapy for psoriasis. *Arch Dermatol* 1985;121:1512-5.
- 33 Finlay AY, Young DW. Short contact crude coal tar therapy for psoriasis. *Clin Exp Dermatol* 1985;10:371-4.
- 34 Farber EM, Abel EA, Cox AJ. Long term risks of psoralen and UVA therapy for psoriasis. *Arch Dermatol* 1983;119:426-31.
- 35 Reshad H, Challoner F, Pollock DJ, Baker H. Cutaneous carcinoma in psoriatic patients treated with PUVA. *Br J Dermatol* 1984;110:299-305.
- 36 Hönigsman H, Stingl G, eds. *Therapeutic photomedicine. Current problems in dermatology 15*. Basle: Karger, 1986.
- 37 Väättäinen N, Hollmen A, Fräki JE. Trimethylpsoralen bath plus ultraviolet A combined with oral retinoid (etretinate) in the treatment of severe psoriasis. *J Am Acad Dermatol* 1985;12:52-5.
- 38 Momtaz-TK, Parrish JA. Combination of psoralens and ultraviolet A and ultraviolet B in the treatment of psoriasis vulgaris. A bilateral comparison study. *J Am Acad Dermatol* 1984;10:481-6.
- 39 Diette KM, Momtaz-TK, Stern RS, Arndt KA, Parrish JA. Role of ultraviolet A in phototherapy for psoriasis. *J Am Acad Dermatol* 1984;11:441-7.
- 40 Menkes A, Stern RS, Arndt KA. Psoriasis treatment with suberythemogenic ultraviolet B radiation and a coal tar extract. *J Am Acad Dermatol* 1985;12:21-5.
- 41 Lowe NJ, Wortzman MS, Breeding J, Koudsi H, Taylor L. Coal tar phototherapy for psoriasis re-evaluated: erythemogenic versus suberythemogenic ultraviolet light with a tar extract in oil and crude coal tar. *J Am Acad Dermatol* 1983;8:781-9.
- 42 Eells LD, Wolff JM, Garloff J, Eaglstein WH. Comparison of suberythemogenic and maximally aggressive ultraviolet B therapy for psoriasis. *J Am Acad Dermatol* 1984;11:105-10.
- 43 Diffey BL, Larko O, Swanbeck GS. UV-B doses received during different outdoor activities and UB-B treatment of psoriasis. *Br J Dermatol* 1982;106:33-41.
- 44 Roenigk HH, Auerbach R, Maibach HI, Weinstein GD. Methotrexate guidelines—revised. *J Am Acad Dermatol* 1982;6:145-55.
- 45 Rau R, ed. *Low dose methotrexate therapy in rheumatic diseases*. Basle: Karger, 1986.
- 46 Van de Kerhof PCM, Hoefnagels WHL, van Haelst UJGM, Mali JWH. Methotrexate maintenance therapy and liver damage in psoriasis. *Clin Exper Dermatol* 1985;10:194-200.
- 47 Geronemus RG, Auerbach R, Tobias H. Liver biopsies v liver scans in methotrexate-treated patients with psoriasis. *Arch Dermatol* 1982;118:649-51.
- 48 Miller JA, Dodd H, Rustin MHA, et al. Ultrasound as a screening procedure for methotrexate induced hepatic damage in severe psoriasis. *Br J Dermatol* 1985;113:699-705.
- 49 Cunliffe WJ, Miller AJ. *Retinoid therapy: a review of clinical and laboratory research*. Lancaster: MTP Press, 1983.
- 50 Saurat JH, ed. *Retinoids: new trends in research and therapy*. Basle: Karger, 1985.
- 51 Sporn MB, Roberts AB, Goodman DS. *The retinoids*. Orlando, Florida: Academic Press, 1984.
- 52 Ward A, Brogden RN, Heel RC, Speight TM, Avery GS. Etretinate. A review of the pharmacological properties and therapeutic efficacy in psoriasis and other skin disorders. *Drugs* 1983;26:9-43.
- 53 Wright V, Moll JMH. *Seronegative polyarthritis*. Amsterdam: North Holland, 1976.
- 54 Ward RA, Wathen RL. Principles of dialysis. Utilization in nonuremic psoriatic patients. *Int J Dermatol* 1982;21:154-8.
- 55 Kiil J, Kiil J, Sogaard H. Surgical treatment of psoriasis. *Lancet* 1985;iii:16-8.
- 56 Menter A, Cram DL. The Goeckerman regimen in two psoriasis day care centers. *J Am Acad Dermatol* 1983;9:59-65.
- 57 Armstrong RB, Leach EE, Fleiss JL, Harber LC. Modified Goeckerman therapy for psoriasis. *Arch Dermatol* 1984;120:313-8.