mg/100 ml)). Blood urine, stool, and sputum cultures; throat and vaginal swabs; and a viral screen all yielded negative results. The chest radiograph, electrocardiogram, and renal ultrasound scan were normal. A plain abdominal radiograph showed an intrauterine device. She was resuscitated with intravenous fluids and hydrocortisone infusion (200 mg/eight hours for 36 hours). Cefotaxime and metronidazole were given but withdrawn when all cultures were reported to have yielded negative results. The patient improved clinically within 24 hours, her renal and hepatic function returning to normal over the subsequent weeks.

On the fourth day after the Mantoux test an intense erythematous induration (5×6 cm) was noted at the site of injection and this persisted for eight weeks. The culture of the lymph node biopsy specimen grew a fully sensitive *Mycobacterium tuberculosis*. The patient was given rifampicin and isoniazid, initially at a dose of 150 mg and 100 mg daily, respectively, to prevent rifampicin induced toxic shock,¹ which was titrated to a maintenance dose of 600 mg and 300 mg daily, respectively, over a week. She was discharged receiving this regimen and remained clinically well.

Comment

There have been previous reports of transient fever, shivering, vomiting, and arthralgia after tuberculin testing.² Severe anaphylactoid reactions, however, as seen in our patient, are rarely reported. In the past they were observed when preparations of tuberculin were used to desensitise patients with known tuberculosis. The so called "paraphylactoid shock" or "systemic tuberculin shock"^{3,4} occurred within 24 hours of the injection and included rigors, nausea and vomiting, arthralgia, headaches, jaundice, adrenal insufficiency, wheezing, and renal failure with proteinuria and oliguria. The reactions either subsided within 48 hours or progressed with further clinical deterioration. Such adverse effects were noted despite the use of a highly purified form of tuberculin.

In our case we had used a dilute solution of 1:10 000 tuberculin (1 IU). We were reassured by the manufacturers that no fault had been found with the rest of the supplied batch. Our patient was healthy apart from tuberculous lymphadenitis and had no history of any drug reaction or allergies. Her serum immunoglobulin concentrations were normal. The delay in the type IV hypersensitivity reaction, occurring on the fourth day after the Mantoux test, may be partly attributed to the hydrocortisone infusion.⁵ The abnormal results obtained in the liver and renal function tests reflected the patient's hypovolaemic state secondary to the anaphylactoid shock.

As the tuberculin test will remain the most simple and important part of our antituberculosis programme in the community all doctors should be aware of its potential hazards, which may require immediate medical attention.

1 Reynolds JEF. *Martindale: the extra pharmacopoeia*. 28th ed. London: Pharmaceutical Press, 1982:1577.

2 Morrison JB. Lymphangitis after tuberculin tests. *Br Med J* 1984;289:413.

3 Jacquelin A, Cornet A, Vilernova P. Incidents during tuberculosis therapy and their significance. *Presse Médicale* 1941;49:884-5.

4 Besiedka A. Tuberculin anaphylaxis and Koch phenomena. *Presse Médicale* 1940;48:973-6.

5 Horne NW. A critical evaluation of corticosteroids in tuberculosis. *Adv Tuberc Res* 1966;15:1-54.

(Accepted 28 April 1986)

Department of Thoracic Medicine, Royal Free Hospital, London NW3 2QG

M A SPITERI, MRCP, registrar

A BOWMAN, MRCP, senior house officer

A R ASSEFI, MB, senior house officer

S W CLARKE, MD, FRCP, consultant physician

Correspondence to: Dr Spiteri.

Natural killer cells in insulin dependent diabetes mellitus

Patients with diabetes mellitus have long been known to be predisposed to certain infections.¹ Less well known is the higher incidence of certain tumours—particularly pancreatic and endometrial tumours—in diabetes.² The aetiology of these infective and neoplastic lesions is likely to be multifactorial, although much recent attention has focused on the estimation of peripheral blood lymphocyte subpopulations.³ These reports, however, have produced conflicting results. None of them mention natural killer cells, which are responsible for non-specific cytotoxicity and are said to represent the body's first line of defence as they attack cells with altered surface membranes such as virus infected and tumour cells.⁴ To respond in this way these cells do not need to be primed by any other mechanism.

We, therefore, investigated the proportions and absolute numbers of

natural killer cells in healthy, insulin dependent diabetics with no known infection or neoplasia.

Patients, methods, and results

We randomly selected 18 insulin dependent diabetics (10 men, eight women) for the study. All had been taking one of the highly purified forms of insulin for more than one year (median time taking insulin 4.5 years, range 2-10) and none had evidence of infection or neoplasia. Natural killer cell proportions and numbers were measured in these and in 48 age and sex matched healthy volunteers.

Peripheral blood lymphocytes from 10 ml of heparinised blood were isolated by density centrifugation on Ficoll-Hypaque and cell suspensions were incubated on ice with 2.5 µl of the various monoclonal antibodies used. These were anti-Leu 11 for natural killer cells,⁵ anti-Leu 4 for T cells, anti-Leu 3 and 2 respectively for T helper and suppressor subpopulations, and finally anti-Leu 12 for B cells (all antibodies Becton Dickinson, USA). Analysis of cell populations was performed on a fluorescence activated cell sorter (FACS 420 Becton Dickinson, USA). In each case 5000 cells were analysed for fluorescence within the population criteria corresponding to lymphocytes defined by forward and right angle scatter produced by the cells as they interrupt the laser beam of the machine. By imposing limits of size and granularity that are proportional to this forward and right angle scatter not only can we concentrate on the cell population we are interested in but we can also exclude red cells, monocytes, and granulocytes.

The table shows that there were no significant differences between the diabetic and control groups for total number of lymphocytes (as assessed in a Coulter counter) or between the proportions of T cells, T helper or suppressor cells, or B cells. There was, however, a significantly lower proportion ($p<0.005$) and absolute number ($p<0.001$) of natural killer cells in diabetics.

Lymphocyte subpopulations in diabetics and healthy volunteers (mean SD)

Lymphocytes	Healthy volunteers (n=48)	Diabetics (n=18)
Lymphocyte count ($\times 10^9/l$)	2.0 (0.8)	1.89 (0.75) NS
B cells (%)	6.4 (1.4)	4.51 (2.7) NS
T cells (%)	77.9 (3.5)	84.13 (5.33) NS
T helper (%)	53.6 (5.4)	56.95 (11.05) NS
T suppressor (%)	27.9 (4.1)	30.55 (12.05) NS
Natural killer cells (%)	16.0 (4.9)	11.21 (3.69) $p<0.005$
Natural killer cells ($\times 10^9/l$)	0.3 (0.1)	0.21 (0.09) $p<0.001$

Statistics by Student's *t* test. Natural killer cell number was calculated from the product of percentage natural killer cells and the total lymphocyte count.

Comment

This study has shown that insulin dependent diabetics have a significantly lower proportion and absolute number of cells bearing the Leu 11 surface marker (interpreted as being natural killer cells) circulating in their peripheral blood than a group of healthy volunteers studied under the same conditions. We suggest that this may be a contributory factor when considering the causes of the known increased incidence of certain infections and tumours in these patients. The explanation of this phenomenon, however, remains unclear, and we do not know, for example, whether it is a result of the insulin used or a cause or effect of the underlying disease. Despite the obvious problems with controlling a number of variables in such a study, we believe that the phenomenon that we have described is worthy of more detailed analysis.

1 Rayfield EJ, Ault M, Keusch G, Brothers MJ, Nechemias C, Smith H. Infection and diabetes: the case for glucose control. *Am J Med* 1982;72:439-50.

2 Kessler II. Cancer and diabetes mellitus: a review of the literature. *J Chronic Dis* 1971;23:579-600.

3 Pozzilli P, Zuccarini O, Lavicoli M, et al. Monoclonal antibodies defined abnormalities of T lymphocytes in type 1 (insulin dependent) diabetes. *Diabetes* 1983;32:91-4.

4 Klein E, Masucci G, Masucci MG, Vanky F. Cell mediated cytotoxicity of natural and activated killer lymphocytes. In: Serrou B, Rosenfield C, eds. *Natural killer cells*. Amsterdam: Elsevier Biomedical Press, 1982.

5 Lanier LL, Le AM, Phillips JH, Warner NL, Babcock GF. Subpopulations of human natural killer cells defined by expression of the Leu 7 (HNK-1) and Leu 11 (NK 15) antigens. *J Immunol* 1983;131:1789-96.

(Accepted 9 May 1986)

Departments of Surgery and Medicine, University of Newcastle upon Tyne, Newcastle upon Tyne NE2 4HH

R G WILSON, FRCS, honorary senior registrar

J ANDERSON, FRCP, FRCOG, honorary consultant physician

B K SHENTON, PHD, top grade scientific officer

M D WHITE, technician

R M R TAYLOR, FRCS, consultant surgeon

G PROUD, FRCS, consultant surgeon

Correspondence to: Mr Wilson.