PRACTICE OBSERVED

Effect on prescribing of the limited list in a computerised group practice

W G IRWIN, K A MILLS, K STEELE

Abstract
The prescribing of drugs in the therapeutic classes that are affected by the government's limited list was investigated in a computerised group practice of just over 3000 patients. Prescribable drugs in categories that are affected by the list were identified for two consecutive six month periods before and one six month period after the introduction of the list. A significant decrease in the prescribing of cough and cold remedies, vitamins, and antacids occurred after the list was introduced, whereas no change occurred in the prescribing of laxatives, benzodiazepines, or analgesics. The prescribing of iron and penicillin increased significantly after the list was introduced, whereas the use of H₂ antagonists and non-steroidal anti-inflammatory drugs showed no significant change.

Introduction

The limited list was introduced on 1 April 1985. Its limitations produced apprehension in the medical profession. It was born out of a desire by the Department of Health and Social Security to reduce prescribing costs, which in 1983 were £1400 million. A blacklist of medicines was introduced which restricted prescribing under the National Health Service in certain categories—namely, tonics and bitters, cough and cold remedies, antacids, laxatives, analgesics to mild to moderate pain, and benzodiazepine sedatives and tranquillisers—to select medicines, some generic, some proprietary. The Audit Great Britain Drug Index, a survey set up to monitor the list by a panel of 200 pharmacists throughout Britain, claimed an

overall decrease of 25% in the number of drugs prescribed in the categories affected by the list.¹ The aim of our study was to analyse retrospectively the data stored in the computer in one group practice to identify any change in prescribing patterns after the list was introduced.

Method

The study was carried out in a group practice of just over 3000 patients that is attached to the department of general practice. Three general practitioners, each a principal holding a joint contract with the university and the Eastern Area Health and Social Services Board (Northern Ireland), provide medical care. On 1 August 1984 they were joined by another practitioner on a half time service contract.

The practice records were first put onto computer early in 1983. The original system consisted of an Apple II microcomputer with a Winchester hard disk to provide sufficient storage space. The system has recently been upgraded to a Stride 440 minicomputer to cope with increased storage requirements and meet research needs. The recording of patient consultations began in October 1983 with the general practitioners filling in encounter forms (figure). The drugs prescribed were entered in the appropriate section (prescription). When a general practitioner wished a particular drug to be prescribed through the computerised repeat medication system he indicated this by filling in the number of repeats in the column marked "Rep."

Drugs that had been prescribed in the therapeutic groups that are affected by the limited list (whether in generic or proprietary form) were identified for two consecutive six month periods (1 April 1984 to 30 September 1984 and 1 October 1984 to 31 March 1985) before and one six month period (1 April 1985 to 30 September 1985) after the introduction of the list. These included drugs that had been issued both by the repeat medication system and during consultations. Figures for the six month period, 1 October 1983 to 31 March 1984, after the general practitioners had started to record encounters were ignored to allow for user mishandling.

The prescribing of drugs in four therapeutic classes that are not directly affected by the list—iron, the penicillins, the H₂ antagonists (cimetidine and ranitidine), and non-steroidal anti-inflammatory drugs—was also averaged for the study period. We hypothesised that prescribing in those groups might increase because practitioners could offer fewer alternatives or previously unused generic formulations in some of the affected categories, namely, vitamins (which include iron and vitamin combinations), cough and cold remedies (excluding inhalations and topical nasal decongestants), antacids, and analgesics. The participating doctors never prescribe bitters and tonics, so these were excluded from the study.

Department of General Practice, The Queen's University of Belfast
W G IRWIN, MD, FRCP, professor
K A MILLS, BSc, research assistant
K STEELE, BSc, MRCP, senior lecturer

Correspondence to: Dr K Steele, Dunlucan Health Centre, 1 Dunlucan Avenue, Belfast BT9 7TH, Northern Ireland.

Encounter form

No. 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

1. General 2. Specialist 3. Traction 4. Traction 5. Other

6. Specialist 7. Specialist 8. Specialist 9. Specialist 10. Specialist 11. Specialist 12. Specialist 13. Specialist 14. Specialist 15. Specialist 16. Specialist 17. Specialist 18. Specialist 19. Specialist 20. Specialist 21. Specialist 22. Specialist 23. Specialist 24. Specialist 25. Specialist 26. Specialist 27. Specialist 28. Specialist 29. Specialist 30. Specialist 31. Specialist 32. Specialist 33. Specialist 34. Specialist 35. Specialist 36. Specialist 37. Specialist 38. Specialist 39. Specialist 40. Specialist 41. Specialist 42. Specialist 43. Specialist 44. Specialist 45. Specialist 46. Specialist 47. Specialist 48. Specialist 49. Specialist 50. Specialist 51. Specialist 52. Specialist 53. Specialist 54. Specialist 55. Specialist 56. Specialist 57. Specialist 58. Specialist 59. Specialist 60. Specialist 61. Specialist 62. Specialist 63. Specialist 64. Specialist 65. Specialist 66. Specialist 67. Specialist 68. Specialist 69. Specialist 70. Specialist 71. Specialist 72. Specialist 73. Specialist 74. Specialist 75. Specialist 76. Specialist 77. Specialist 78. Specialist 79. Specialist 80. Specialist 81. Specialist 82. Specialist 83. Specialist 84. Specialist 85. Specialist 86. Specialist 87. Specialist 88. Specialist 89. Specialist 90. Specialist 91. Specialist 92. Specialist 93. Specialist 94. Specialist 95. Specialist 96. Specialist 97. Specialist 98. Specialist 99. Specialist 100. Specialist

1. General 2. Specialist 3. Traction 4. Traction 5. Other

6. Specialist 7. Specialist 8. Specialist 9. Specialist 10. Specialist 11. Specialist 12. Specialist 13. Specialist 14. Specialist 15. Specialist 16. Specialist 17. Specialist 18. Specialist 19. Specialist 20. Specialist 21. Specialist 22. Specialist 23. Specialist 24. Specialist 25. Specialist 26. Specialist 27. Specialist 28. Specialist 29. Specialist 30. Specialist 31. Specialist 32. Specialist 33. Specialist 34. Specialist 35. Specialist 36. Specialist 37. Specialist 38. Specialist 39. Specialist 40. Specialist 41. Specialist 42. Specialist 43. Specialist 44. Specialist 45. Specialist 46. Specialist 47. Specialist 48. Specialist 49. Specialist 50. Specialist 51. Specialist 52. Specialist 53. Specialist 54. Specialist 55. Specialist 56. Specialist 57. Specialist 58. Specialist 59. Specialist 60. Specialist 61. Specialist 62. Specialist 63. Specialist 64. Specialist 65. Specialist 66. Specialist 67. Specialist 68. Specialist 69. Specialist 70. Specialist 71. Specialist 72. Specialist 73. Specialist 74. Specialist 75. Specialist 76. Specialist 77. Specialist 78. Specialist 79. Specialist 80. Specialist 81. Specialist 82. Specialist 83. Specialist 84. Specialist 85. Specialist 86. Specialist 87. Specialist 88. Specialist 89. Specialist 90. Specialist 91. Specialist 92. Specialist 93. Specialist 94. Specialist 95. Specialist 96. Specialist 97. Specialist 98. Specialist 99. Specialist 100. Specialist

The form on which consultations with patients were recorded. RP=Blood pressure; PFK=Peak flow rate; FEV/FVC=Forced expiratory volume/forced vital capacity; NGP=patient (N) or general practitioner (GP) initiated consultation; PS=added to problem summary in A4 card; RGP=Royal College of General Practitioners.

Number of prescription items issued in each therapeutic group

	Period A 1 April 1984 to 30 Sept 1984 (1527 consultations)	Period B 1 Oct 1984 to 31 March 1985 (1790 consultations)	Period C 1 April 1985 to 30 Sept 1985 (1567 consultations)	Significance*			
	No	No/1000 consultations	No	No/1000 consultations	A/B	B/C	A/C
Vitamins							
Patients receiving scripts	16	29.4	97	26.2	48	13.1	NS
Total No. of items	146	44.7	144	37.9	57	15.5	NS
Iron							
Patients receiving scripts	16	29.4	97	26.2	48	13.1	NS
Total No. of items	146	44.7	144	37.9	57	15.5	NS
Cough and cold remedies							
Patients receiving scripts	22	5.5	16	4.5	23	5.7	NS
Total No. of items	18	4.7	34	9.2	32	8.1	NS
Antacids							
Patients receiving scripts	199	60.9	146	40.9	53	14.8	NS
Total No. of items	274	81.9	244	66.5	266	66.5	NS
Penicillins							
Patients receiving scripts	178	54.5	318	85.9	343	85.4	NS
Total No. of items	196	60.0	391	107.6	365	99.5	NS
Aspirin							
Patients receiving scripts	93	28.5	110	29.7	130	30.0	NS
Total No. of items	140	42.9	166	46.2	187	51.0	NS
H₂ antagonists							
Patients receiving scripts	119	35.9	90	24.5	45	17.7	NS
Total No. of items	65	19.9	114	30.8	140	38.2	NS
Laxatives							
Patients receiving scripts	54	16.5	73	19.7	49	18.8	NS
Total No. of items	112	34.3	147	40.8	157	40.2	NS
Analgesics							
Patients receiving scripts	242	74.1	272	74.5	213	68.0	NS
Total No. of items	400	122.4	466	127.9	411	112.4	NS
Non-steroidal anti-inflammatory drugs							
Patients receiving scripts	175	53.5	180	50.3	206	56.7	NS
Total No. of items	340	104.0	357	99.3	382	98.5	NS
Benzodiazepines							
Patients receiving scripts	175	53.5	180	50.3	182	46.6	NS
Total No. of items	418	125.6	444	124.7	525	143.2	NS

*Significance (p < 0.05).
Significance based on the test used for comparison of death rates: $2 \times \sqrt{SE^2 + SE^2}$, where SE = standard error, SE² = m²/d, m = rate, and d = number of years.

similar pattern was noted in this study. The expected seasonal rise in the prescribing of penicillin of 75% between periods A and B was followed by a decrease of only 6% between periods B and C. There was also a significant increase in the prescribing of penicillin between the seasonally equivalent periods A and C. Doctors tended to prescribe the more potent antibiotic when their range of symptomatic remedies was limited. Perhaps this trend will be reversed as practitioners become more familiar with the available symptomatic preparations.

The prescribing of antacids has not changed with regard to the number of patients receiving scripts or the number of items issued on repeat prescriptions. Non-repeat antacid prescribing, which had been increasing significantly between periods A and B, decreased significantly between periods B and C. This decrease was not reflected by a significant rise in H₂ antagonist prescribing between periods B and C (except for repeat items). The significant rise in prescribing of these drugs between periods A and C may be secondary to the addition of H₂ antagonists to the data sheet for non-ulcer dyspepsia in July 1984. The DHSS claim that since the inception of the limited list prescribing of H₂ antagonists has increased by approximately 15% overlooks this trend.¹

Although the prescribing of laxatives decreased this was not significant. The decrease was due to fewer non-repeat items being prescribed, whereas before 1 April 1985 there had been a significant increase in non-repeat laxative prescribing that was mainly attributable to increased prescribing of fibre in period B. There was no significant change in analgesic prescribing throughout the study periods. This probably reflects the genuine needs of patients for pain killing drugs and the reasonable range still prescribable under the NHS. Prescribing of non-steroidal anti-inflammatory drugs did not alter over the study period. Although there was a significant rise in non-repeat benzodiazepine prescribing over the study period, there is no evidence that it was directly affected by the limited list. It is reassuring that any rise has been in non-repeat prescribing. This is reassuring that any rise has been in non-repeat prescribing. This is reassuring that any rise has been in non-repeat prescribing.

reflects the attitude of the practitioners in the study, who felt that repeat prescribing of these drugs can lead to addiction. The figures also reflect the difficulty experienced in changing patients to alternative treatment.

We suggest that the limited list has had a limited effect in the practice studied. The prescribing of vitamins, cough and cold remedies, and antacids decreased significantly, whereas the prescribing of iron and penicillin has increased since the limited list was introduced. Analgesic prescribing appears to be impervious to the influence of the list, whereas benzodiazepine use is still on the increase.

A recent study by Anderson and Reilly showed that generic prescribing can save approximately 8.8% of the drug bill.¹ The Greenfield report recommended not only that general practitioners should participate in making policy on therapeutics but also that local formularies and generic prescribing should be introduced.² If the DHSS would encourage this perhaps it would lead to cost effective changes in prescribing habits instead of the mixed response consequent upon sanctions being imposed.

We thank Dr Desmond Merrett for statistical advice and Miss Jennifer Windrum for typing the manuscript.

References

1. Anonymous. *Audit Great Britain Drug Index*. AGI: Tynemouth, North East, 1985.
2. Greenfield. *Report of the Greenfield Committee on the National Health Service*. London: HCSP 401, 1985.
3. Anonymous. *Generic prescribing*. Negative report. *Gen* 1985; Jan 16.
4. Conley J. Many more drugs are back on the limited list. *Gen* 1985; Nov 29.
5. Anderson RJ, Reilly PM. On the use of generic drugs. *Br Med J* 1984; 289: 1129.
6. Department of Health and Social Security. *Report of the national working group on effective prescribing*. London: DHSS, 1985. (Generalised report).

(Accepted 2 June 1986)

Role of general practitioners in the care of disabled young adults

F S W BRIMBLECOMBE, D L KUH, C J LAWRENCE, R C SMITH

Abstract

Services for handicapped children end when these patients become young adults. In the Exeter Health Authority district 283 disabled young adults were interviewed about their unmet needs. Many wished for advice and counselling. A quarter had not visited their general practitioner in the previous year, and two thirds of these had not had a general assessment or seen a specialist.

Introduction

Generally, handicapped children and their families are well cared for, but what happens when these children grow up? We evaluated

the continuity of care in young chronically disabled adults in the Exeter Health Authority district and studied the nature and extent of their unmet needs.

Methods

A confidential register was made of every person and their families in the district who were aged 16 to 25 and suffering from a life long impairment. Patients in whom a primary psychiatric illness led to the handicap were excluded. Five hundred and two young people who were suffering from mental retardation, physical impairment, and sensory impairment were sampled, of whom 383 (76%) agreed to be interviewed (table 1). The catchment area contains 3075 people aged 16 to 25. From those identified from all the registers we ascertained that 2.6% had an appreciable mental, physical, or sensory impairment.

Our control group comprised 152 non-handicapped young people who were drawn from general practice age-sex registers and matched for age, sex, and place of residence. Both the study group and the control group were interviewed by the same interviewers using similar questionnaires for the two groups. In addition, the parents and the main service provider (where one existed) were interviewed. The main service provider is an outsider who is concerned in the care of the young adult, such as a member of the school staff, a nurse, a social worker, or an occupational therapist. The questionnaires covered the nature and extent of the disability if any; type of education, training, employment, or current occupation; frequency of daily living activities including personal mobility and transport; need for personal

University of Exeter
F S W BRIMBLECOMBE, MR, FRCP, honorary professor of child health
D L KUH, MR, research fellow, department of child health
C J LAWRENCE, MR, lecturer, department of mathematical statistics and operational research
R C SMITH, MR, MRCP, medical research fellow, department of child health
Correspondence to: Professor F S W Brimblecombe, Bowmore House, Royal Devon and Exeter Hospital (Winfold), Barrack Road, Exeter EX2 5DW.

help or aids; general health; contacts with general practitioners, hospitals, specialists, and paramedical services; proficiency at money management; social activities; friends; personal relationships; living arrangements; habits; dietary, behavioural problems; self image and confidence; expectations and ambitions for the future; and areas of need.

TABLE 1—Prevalence of impairment

	No sampled from all registers	No who refused to be interviewed	No (%) who agreed to be interviewed	No in revised category of predominant impairment after interview
Severe mental handicap	141	20	121 (86)	115
Mild mental handicap	91	36	55 (60)	42
Physical handicap	198	45	153 (76)	132
Sensory handicap	54	18	36 (67)	45
Psychiatric handicap	1	0	1 (100)	0
Unappreciated handicap	1	0	1 (100)	27
Total No.	502	119	383 (76)	383

TABLE 2—Type of residence by age of study and control groups

	16-18 years	
	Study group	Control group
	No (%)	No (%)
Living in parental home or with other family members	84 (82)	33 (97)
Living independently	12 (12)	1 (3)
Supervised lodgings	1 (1)	0
Living in residential care: hospital, staffed home, care communities, etc	13 (13)	0

TABLE 3—Visits to family doctor in past year

	No(%) of young adults	
No of visits	Study group (n = 333)	Control group (n = 147)
None	82 (25)	44 (30)
1-5	190 (57)	84 (57)
6-10	33 (10)	12 (8)
10	28 (8)	7 (5)

Note: The handicapped young people living in residential care are not included.

Note: The handicapped young people living in residential care are not included.

The interviews were carried out by three health visitors, one with experience with deaf people, a nurse, two teachers, a social worker, and an occupational therapist. All were middle aged women who were prepared to listen.

Results

More than three quarters (80%) of the 383 handicapped young adults were living at home (table 1) and so might rely on their family doctor for primary medical care. Many of the young adults were not receiving medical services that they, their families, or the main service provider wanted: 66 (17%) thought that a full medical assessment would be helpful and 69 (18%) wanted genetic counselling, 99 (26%) sexual counselling, 68 (18%) family counselling, 37 (10%) advice on behavioural problems, and 100 (26%) advice about basic health problems: many said that they had not had enough information about their disability and did not know who to turn to for medical advice. Several had never seen their family doctor alone but would have liked to do so. Less than half (48%) had seen a specialist in the previous year. The unmet medical need was identified by the young adult in 10.4% of cases, by their families only in 4.2%, and by their main service provider alone in 1.5%.

The handicapped young adults did not visit their family doctor more often

than the control group, except for a small number who made more than 10 visits a year (table 2). Of the 82 who had not seen their family doctor in the previous year, 28 (34%) had not seen a hospital specialist or undergone a general assessment. Almost all of them and their families seemed unaware that their family doctor might be able to help.

Many of the handicapped young people have low self esteem and expectations for their future (table 3). The families emphasised the inadequacy of their own preparation for the transition of their disabled child into adulthood.

Discussion

The findings show that there is often little communication between handicapped young adults and general practitioners. Handicapped children are cared for by specialised services and bypass the family doctor. Such services are withdrawn when the children reach 16 or 19, just when they are likely to experience behavioural, sexual, and depressive problems that can be dealt with by the family doctor. Moreover, a full interdisciplinary reassessment of the handicapped young adult is mandatory in the light of medical advances, possible changes of the patient's condition, and

TABLE 4—Self image and self esteem of young handicapped adults

	Study group (%)	Control group (%)
I often think things are not much for me	30	15
Other people often decide things for me	26	26
The very other people treat me as others make me angry	42	21
Sometimes I think I have to work harder than anyone else to prove how good I am	49	24
I often feel lonely	12	12
People often make fun of me	29	14
In groups of people I often feel the odd one out	40	23
I am very uncomfortable in the company of strangers	40	24

For each of these, p < 0.001. The difference is significant at the 0.1% level.

Note: The maximum sample size used for comparison was 273.

his or her increasing maturity. As specialist medical care outside hospitals is not sufficiently organised to provide this service, the general practitioner should arrange for such a reassessment. Knowledge about all local resources and facilities to help these disabled people is essential. The family doctor is also often uniquely placed to reconcile the conflicts arising in a family in which there is a handicapped young adult.

The second phase of the Exeter project is to work with all statutory and voluntary services to help to create enabling services in each locality that are more patient oriented than those now available. Meanwhile, we ask family doctors to pay more attention to the needs of handicapped young adults in their practice.

References

1. Brimblecombe FSW, Kuh DL, Smith RC. The needs of handicapped young adults. *Br Med J* 1986; 293: 1136-1139.
2. Anonymous. *Report of the Greenfield Committee on the National Health Service*. London: HCSP 401, 1985.

(Accepted 7 August 1986)