- 5 Kitchen WH, Campbell NT, Drew JH, Murton LJ, Roy RND, Yu VYH. Provision of perinatal services and survival of extremely low birthweight infants in Victoria. *Med J Aust* 1983;2:314-8.
- 6 World Health Organisation. Recommended definitions, terminology and format for statistical table related to the perinatal period and use of a new certificate for cause of perinatal deaths. *Acta Obstet Gynecol Scand* 1977;56:247-53.
- 7 Spinnato, Sibai BM, Shaver DC, Anderson GD. Inaccuracy of Dubowitz gestational age in low birth weight infants. *Obstet Gynecol* 1984;63:491-5.
- 8 Kitchen WH, Yu VYH, Orgill AA, *et al*. Collaborative study of very low birthweight infants: outcome of two year old survivors. *Lancet* 1982;ii:1457-61.
- 9 Orgill AA, Astbury J, Bajuk B, Yu VYH. Early development of infants 1000 g or less at birth. *Arch Dis Child* 1982;57:823-7.
- 10 Yu VYH, Wong PY, Bajuk B, Orgill AA, Astbury J. Outcome of extremely low birthweight infants. *Br J Obstet Gynaecol* 1986;93:162-70.
- 11 Stewart AL, Turcan DM, Rawlings G, Reynolds EOR. Prognosis for infants weighing 1000 g or less at birth. *Arch Dis Child* 1977;52:97-104.
- 12 Kitchen WH, Bajuk B, Lissenden JV, Yu VYH. Intrauterine growth charts from 24 to 29 weeks' gestation. *Aust Paediatr J* 1981;17:269-72.
- 13 Britton SB, Fritzhardinge PM, Ashby S. Is intensive care justified for infants weighing less than 801 g at birth? *J Pediatr* 1981;99:937-43.
- 14 Bennett FC, Robinson NM, Sells CJ. Growth and development of infants weighing less than 800 grams at birth. *Pediatrics* 1983;71:319-23.
- 15 Hirata T, Epcar JT, Walsh A, *et al*. Survival and outcome of infants 501 to 750 gm: a six year experience. *J Pediatr* 1983;102:741-8.
- 16 Buckwald S, Zorn WA, Egan EA. Mortality and follow-up data for neonates weighing 500-800 g at birth. *Am J Dis Child* 1984;138:779-82.
- 17 Hack M, Fanaroff AA. Changes in the delivery room care of the extremely small infant (<750 g). Effects on morbidity and outcome. *N Engl J Med* 1986;314:660-4.
- 18 Kitchen W, Ford G, Orgill AA, *et al*. Outcome of infants of birthweight 500-599 g: a regional study of 1979-1980 births. *J Pediatr* 1984;104:921-7.
- 19 Dillon WP, Egan EA. Aggressive obstetric management in late second-trimester deliveries. *Obstet Gynecol* 1981;58:685-90.
- 20 Herschel M, Kennedy JL, Jr, Kayne HL, Henry M, Cetrulo CL. Survival of infants born at 24 to 28 weeks' gestation. *Obstet Gynecol* 1982;60:154-8.
- 21 Yu VYH, Watkins A, Bajuk B. Neonatal and postneonatal mortality in very low birthweight infants. *Arch Dis Child* 1984;138:779-99.
- 22 Bhat R, Raju TNK, Vidyasagar D. Immediate and long-term outcome of infants less than 1000 grams. *Crit Care Med* 1978;6:147-50.
- 23 Pape KE, Buncic RJ, Ashby S, Fitzhardinge PM. The status at two years of low birthweight infants born in 1974 with birthweights of less than 1001 g. *J Pediatr* 1978;92:253-60.
- 24 Shennan AT, Milligan JE, Hoskins EM. Perinatal factors associated with death or handicap in very preterm infants. *Am J Obstet Gynecol* 1985;151:231-8.
- 25 Khoury MJ, Marks JS, McCarthy BJ, Zaro SM. Factors affecting the sex differential in neonatal mortality: the role of respiratory distress syndrome. *Am J Obstet Gynecol* 1985;151:777-82.
- 26 McCarthy BJ, Sachs BP, Layde PM, Burton A, Terry JS, Rochat R. The epidemiology of neonatal deaths in twins. *Am J Obstet Gynecol* 1981;141:252-6.
- 27 Hawrylyshyn PA, Barkin M, Bernstein A, Papsin FR. Twin pregnancies—a continuing perinatal challenge. *Obstet Gynecol* 1982;59:463-6.
- 28 Barrett JM, Staggs SM, Van Hooydonk JE, Growdon JH, Killam AP, Boem FH. The effect of delivery upon neonatal outcome in premature twins. *Am J Obstet Gynecol* 1982;143:360-4.
- 29 Chervenak FA, Johnson RE, Berkowitz RL, Grannum P, Hobbins JC. Is routine cesarean section necessary for vertex-breech and vertex-transverse twin gestations? *Am J Obstet Gynecol* 1984;148:1-5.
- 30 Bell D, Johnansson D, McLean FH, Usher RH. Birth asphyxia, trauma, and mortality in twins: has cesarean section improved outcome? *Am J Obstet Gynecol* 1986;154:235-9.
- 31 Yu VYH, Downe L, Astbury J, Orgill AA, Bajuk B. Perinatal factors associated with adverse outcome in extremely low birthweight infants. *Arch Dis Child* 1986;61:554-8.
- 32 Szymonowicz W, Yu VYH, Bajuk B, Astbury J. Neurodevelopmental outcome of periventricular haemorrhage and leukomalacia in infants 1250 grams or less at birth. *Early Hum Dev* 1986;14:1-7.
- 33 Szymonowicz W, Yu VYH. Periventricular haemorrhage and leukomalacia in extremely low birthweight infants. *Aust Paediatr J* 1986;22:207-10.
- 34 Yu VYH, Orgill AA, Lim SB, Bajuk B, Astbury J. Growth and development of very low birthweight children recovering from bronchopulmonary dysplasia. *Arch Dis Child* 1983;58:791-4.
- 35 Morgan MEI. Late morbidity of very low birthweight infants. *Br Med J* 1985;291:171-3.
- 36 Astbury J, Orgill AA, Bajuk B, Yu VYH. Neonatal and neurodevelopmental significance of behaviour in very low birthweight children. *Early Hum Dev* 1985;11:113-21.
- 37 Astbury J, Orgill AA, Bajuk B, Yu VYH. The physical and neurobehavioural sequelae of growth failure in appropriate for gestational age very low birthweight children. *Dev Med Child Neurol* (in press).
- 38 Enhorning G, Shennan A, Possmayer F, Dunn M, Chen CP, Milligan J. Prevention of neonatal respiratory distress syndrome by tracheal instillation of surfactant: a randomised clinical trial. *Pediatrics* 1986;76:145-53.
- 39 Boynton BR, Mannino FL, Davis RF, Kopotic RJ, Friederichsen G. Combined high-frequency oscillatory ventilation in critically ill neonates. *J Pediatr* 1984;105:297-302.
- 40 Yu VYH, Bajuk B, Orgill AA, Astbury J. Viability of infants born at 24 to 26 weeks' gestation. *Ann Acad Med* 1985;14:563-71.
- 41 Yu VYH, Davis NG, Mercado M, Bajuk B, Astbury J. Subsequent pregnancy following the birth of an extremely low birthweight infant. *Aust NZ J Obstet Gynaecol* 1986;26:115-9.

(Accepted 14 August 1986)

Hip fractures in healthy patients: operative delay versus prognosis

R N VILLAR, S M ALLEN, S J BARNES

Abstract

One hundred and forty five women who had undergone hemiarthroplasty for a subcapital fracture of the femoral neck but who were otherwise fit were studied to determine whether undue delay between injury and operation influenced their social circumstances three months after surgery. The median delay for those patients who showed good rehabilitation at three months was 29 hours, but for those who showed poor rehabilitation it was 57 hours. This difference was significant.

It is suggested that a subcapital fracture in an otherwise fit elderly patient should therefore be regarded as a surgical emergency.

Introduction

It is widely accepted that elderly women with hip fractures undergo surgery on the next available operating list, frequently 24 hours or

more after admission. We present a retrospective study relating this operative delay to the social circumstances of the patient three months after surgery.

Patients, methods, and results

We studied 205 consecutive women patients, who had been admitted over 18 months and had undergone hemiarthroplasty for a displaced subcapital fracture of the femoral neck. Of the 205 patients, 60 had intercurrent illness that was likely to interfere with postoperative rehabilitation and were excluded from the study.

The social circumstances of each patient were assessed on admission and three months after surgery as (a) independent if she lived without help from others; (b) sheltered if she received substantial help with daily activities; or (c) resident in hospital if she lived in an institution.¹ The time delay between injury and operation and typical reasons for delay were also recorded. At the review three months after surgery each patient was placed in one of two groups based on her social circumstances at that time. Group A consisted of patients whose circumstances were similar to those on admission, and group B consisted of patients whose circumstances had deteriorated and those who had died.

Statistical analysis was by Student's *t* test. Results for operative delay produced a skewed distribution and log_e (operative delay) was therefore used to correct the findings.

Of the 145 patients, 98 (68%) were admitted from an independent environment, 41 (28%) were sheltered, and 6 (4%) were resident in hospital. After three months similar percentages of each social category could be placed in group A (81% independent, 78% sheltered, and 83% resident in hospital). No patient improved after surgery; a patient stayed the same, became worse, or died.

Patients in group A had a median delay of 29 (range 6-184 h; mean 31.7 h) and group B patients a median delay of 57 (range 24-528 h; mean 74.7 h).

Department of Orthopaedic Surgery, Southampton General Hospital, Southampton, Hants

R N VILLAR, BSC, FRCS, orthopaedic registrar
S M ALLEN, MB, BS, orthopaedic senior house officer
S J BARNES, BM, orthopaedic senior house officer

Correspondence to: Mr R N Villar, Department of Orthopaedic Surgery, Addenbrooke's Hospital, Cambridge CB2 2QQ.

The difference between the mean operative delays in the two groups was therefore 43 hours (95% confidence interval 41.35 to 44.45 h; $t=6.0$, 143 df; $p<0.0001$). Despite a slight difference in the median ages for groups A and B (80 (range 47-96) and 82 (range 69-93), respectively) age distributions were similar. Typical reasons for delay included operator fatigue, delay in diagnosis, lack of theatre time, industrial action, and lack of patient's consent.

Discussion

This retrospective study suggests that the longer an otherwise fit patient has to wait for her hip fracture to be treated the less she will progress after discharge, regardless of her social circumstances. Many hospitals use trauma list systems, patients admitted one day being held over to the next before undergoing surgery. Previous work has shown that delays of 13-48 hours are not necessarily detrimental to patients' wellbeing.² Such work, however, emphasises the delay between admission and surgery rather than between injury and operation, the two observations often being widely different. Some of our patients had languished at home for several days before being admitted to hospital; thus in these cases

there was a short interval between admission and surgery but an unacceptable delay between injury and surgery.

Loss of independence is perhaps the most important social problem relating to patients with fractures of the femoral neck, resulting in blocked beds and overloading of social and primary care services. What then should be done to avoid this for the elderly patient who arrives in the accident and emergency department with a hip fracture? We believe that if she is otherwise fit every effort should be made for her to undergo surgery at the earliest opportunity. To include such patients in a standard trauma list system, however convenient it may be for surgeons, anaesthetists, and administrators, is to exacerbate a situation that is already out of control.

References

- 1 Hall G, Ainscow DAP. Comparison of nail-plate fixation and Ender's nailing for intertrochanteric fractures. *J Bone Joint Surg [Br]* 1981;63:24-8.
- 2 Aldrete JA, Davis HS, Hingson RA. Anesthesia factors in the surgical management of hip fractures. *J Trauma* 1967;7:818-26.

(Accepted 20 August 1986)

Trial of early nifedipine in acute myocardial infarction: the Trent study

R G WILCOX, J R HAMPTON, D C BANKS, J S BIRKHEAD, I A B BROOKSBY, C J BURNS-COX, M J HAYES, M D JOY, A D MALCOLM, H G MATHER, J M ROWLEY

Abstract

Over 30 months 9292 consecutive patients admitted to nine coronary care units with suspected myocardial infarction were considered for admission to a randomised double blind study comparing the effect on mortality of nifedipine 10 mg four times a day with that of placebo. Among the 4801 patients excluded from the study the overall one month fatality rate was 18.2% and the one month fatality rate in those with definite myocardial infarction 26.8%. A total of 4491 patients fulfilled the entry criteria and were randomly allocated to nifedipine or placebo immediately after assessment in the coronary care unit. Roughly 64% of patients in both treatment groups sustained an acute myocardial infarction. The overall one month fatality rates were 6.3% in the placebo treated group and 6.7% in the nifedipine treated group. Most of the deaths occurred in patients with an in hospital

diagnosis of myocardial infarction, and their one month fatality rates were 9.3% for the placebo group and 10.2% for the nifedipine group. These differences were not statistically significant. Subgroup analysis also did not suggest any particular group of patients with suspected acute myocardial infarction who might benefit from early nifedipine treatment in the dose studied.

Introduction

Nifedipine is a substituted dihydropyridine with calcium channel blocking properties.¹ Compared with verapamil it has very little cardiac electrophysiological effect.² In experimental myocardial infarction in animals pretreatment with nifedipine in a dose carefully regulated to avoid a large fall in blood pressure and reflex tachycardia causes an increase in coronary blood flow in both normally perfused and ischaemic areas of the heart, delays the release of cytoplasmic enzymes and the intracellular accumulation of calcium, preserves intracellular stores of adenosine triphosphate, and reduces infarct size.³⁻⁶ Nifedipine is active during periods of ischaemia and also during subsequent reperfusion.³ This has led to speculation that the drug may have a "cardioprotective" action in man.

In angina nifedipine has been shown to increase coronary perfusion and decrease afterload with minimal decrease in contractility^{7,8} and is thereby thought to stabilise the imbalance between oxygen supply and oxygen demand. Roberts and coworkers have shown that nifedipine produces similar haemodynamic effects in patients with acute infarction, suggesting that in this condition also the drug may be capable of improving a myocardial oxygen deficiency.⁹ Nifedipine also inhibits coronary artery spasm^{10,11} and exerts a mild antiaggregatory effect on platelets,¹² both effects that have been implicated in myocardial infarction.¹³

The study by Roberts *et al* also suggested that treatment with

TRENT Study Group

Coordinator: Dr R G Wilcox. *Chairman:* Professor J R Hampton (University Hospital, Nottingham). *Principal investigators:* Dr D C Banks (City Hospital, Nottingham); Dr J S Birkhead (General Hospital, Northampton); Dr I A B Brooksbey (Norfolk and Norwich Hospital, Norwich); Dr C J Burns-Cox (Frenchay Hospital, Bristol); Dr M J Hayes (Princess Margaret Hospital, Swindon); Dr M D Joy (St Peter's Hospital, Chertsey); Dr A D Malcolm (Airedale General Hospital, Keighley); Dr H G Mather (Southmead Hospital, Bristol); Dr J M Rowley (University Hospital, Nottingham); Dr R G Wilcox (University Hospital, Nottingham). *Data analysis:* Dr A M Skene, Pamela Skene, Professor A F M Smith (department of mathematics, University of Nottingham). *Ethical review committee:* Professor G A Rose, Professor M J Healy, Professor P Sever (London School of Hygiene and Tropical Medicine, St Mary's Hospital, London). *Bayer clinical research coordinators:* Dr G Macdonald, Dr K Young.

Correspondence to: Dr R G Wilcox, Department of Medicine, University Hospital, Queen's Medical Centre, Nottingham.