PRACTICE OBSERVED

Good Practice

What is a good GP?

SUSAN HOLMES

I was born in 1948—just three weeks before the birth of the National Health Service. My mother had a home confinement and was attended, I am told, by an ill tempered midwife and her own GP. Although technically a home confinement, I was not born in my own home but at the house of my mother's friend. She had a bathroom. We did not. The fact that this superior dwelling was some four miles outside our GP's area did not deter him from attending the birth. Nor was he perturbed by my mother's incongruous wish to have the Preface to Act One of *La Traviata* playing on the radio during the second stage of labour.

I have never heard my mother refer to Dr S as a "good GP." It was simply accepted that as the family doctor he was there when required by the family. He came at any hour of the day or night. He always remembered who we all were and recalled our various ailments without reference to notes. He was always there—like the milkman, the coalman, and the postman. As a child it never really crossed my mind that he might have any other patients. He appeared to be ours exclusively.

But those were the heady days of an infant Health Service. Forty years on, and middle aged in Maggie's Britain, things aren't quite the same. Now it seems you have to shop around to get the sort of service from a GP which used to be commonplace. More people, living longer, fewer resources, perhaps.

Variety

Since 1948 I have been ministered to by nine different general practitioners. Following the births of my four children I have naturally made use of them rather more frequently. The GPs have

88 Basford Park Road, Basford, Newcastle, Staffs ST5 5PP
SUSAN HOLMES, journalist

varied in quality and attitude. One practised in a gloomy Camden Town basement, operated no appointment system, and made it a policy to keep patients waiting at least an hour. Another, with delusions of grandeur, operated from a Victorian mansion and kept no one waiting for more than five minutes. This was a pity, as the waiting area was a fine marbled hall with an elegant staircase and huge exotic plants. He eventually gave up his home and practice to open a private clinic. Yet another GP died, I am told, by his own hand.

Only one of the nine has in every respect come up to my naive, rose coloured view of how an old fashioned family doctor—a "good GP"—should perform. Treating me and mine with an unperturbed attitude and with total respect. He also had another overwhelming virtue: he employed one solitary receptionist who called all the patients by their first name and recognised them instantly on sight. She also had a wonderful reputation among the patients of "always being able to tell you in" however busy the day's surgery.

I believe that many people judge a doctor by the initial response that they get from the receptionist—and these women have become a bit of an obsession with me of late. Often, they are not so much there to help and advise the patient as to play the part of doctors' minders. They are the medical equivalent of bouncers at the Palais. There to weed out the unsuitable, the awkward, the emotional—and often the ill. Some even demand to know symptoms and attempt diagnosis. Anything to fend off the attacking hordes and give the doctor a peaceful life.

They do the GPs a grave disservice. Many people judge a GP by the quality and attitude of the receptionist—or receptionists. Many operate as an army behind glass, and they are always wrong when they have comrades to back them up. I have heard old men shouted at and humiliated in front of a waiting room full of people because they were not a prescription card or mistaken an appointment time. I have seen a woman reduced to tears by being told publicly, by a receptionist, details of her husband's low sperm count and the subsequent tests the couple would need to undergo for their infertility problem.

Results

A total of 748 subjects were approached, of whom five men and five women refused and 18 returned incomplete questionnaires. Thus 700 (96%) subjects, 186 men and 514 women, were included in the analysis (table 1). Social class was defined according to Goldthorpe's scheme (table 1). EAT scores—Seventy one (13.3%) women and five (2.7%) men scored above the threshold (19.20). High scorers were more often women (diff=10.9%, 95% confidence limits (CI)=6.14, $p=0.001$), were older (diff=1.2, $p=0.001$) and between 21 and 25 years of age (diff=1.2%, 95% CI=0.22, $p=0.001$), $p<0.05$.

Diagnostic group high scorers—Sixty nine (91%) subjects agreed to be interviewed and were allocated to one of six diagnostic groups on the basis of the eating interview (table 1). Bulimia nervosa was diagnosed according to Russell's criteria.¹¹ All the subjects with bulimia were of normal weight (table 1), but three subjects, including the man, had a past history, on

TABLE 1—Demographic characteristics of screened population

	No.	(%)
Sex		
Men	186	26
Women	514	74
Age (years)		
<25	162	23
25-30	230	33
31-35	168	24
36-40	121	17
41+	9	1
Social class		
I	34	5
II	156	22
III	202	29
IV	27	4
V	79	11
VI, VII	19	3
Unemployed	14	2
Unemployed	14	2
Unemployed	14	2
Unemployed	14	2
Unemployed	14	2

*Only one of the six previous employment was given and was not used in the analysis. The other five subjects were working when screened for this study.

TABLE 11—Diagnostic groups and comparison low scorers

	Men	Women	Mean EAT score
Bulimia nervosa	1	6	35.7
Partial syndrome	1	13	32.1
Obsessive compulsive disorder	1	13	28.3
Normal eating	4	23	25.0
Obsessive	1	4	35.5
Disordered	1	4	35.5
Comparison group	8	32	25.7

EAT=eating attitudes test.

weight and menstrual and hormonal criteria, that suggested anorexia nervosa. Two of the seven were of West Indian origin, born in the United Kingdom. Five subjects with partial syndrome had a full syndrome in the past, and four of the obsessional dieters had at least a partial syndrome previously. Obsessional dieters were preoccupied with diet and weight but not pathologically. The three groups of women were caught in a cycle of dieting and overeating. The term disturbed was used to describe four women who, although having little concern about diet or weight, had chronic psychosocial lives and indicated their distress by responding non-specifically to the EAT as if it were a more general psychiatric questionnaire. Thus 11% of women and 0.5% of men had bulimia nervosa, and, in addition, 2.8% of women had a partial syndrome eating disorder. These high scorers were compared with low scorers of 10 men and 35 women selected randomly, eight men and 32 women (89%) were finally interviewed. No subjects with bulimia or partial syndrome were found.

Function of the EAT—At the recommended cutoff of 19.20, in detecting subjects with bulimia or partial syndrome the EAT functioned with a sensitivity of 0.1, a specificity of 0.91, and a positive predictive value of 0.33.

In view of the small number of high scorers among the men on the EAT the remaining analysis concerns women only. Weight—Reported weight was within 4% of actual weight at interview, and thus most measures could be placed on reported weights. Therefore, taking the screened population as a whole, high scorers on the EAT weighed more than low scorers (mean diff=15.7 lb (7.1 kg); 95% CI=7.7-23.7; $t=3.829$, $df=416$, $p<0.001$), although the difference was not statistically significant when the difference was calculated for the comparison group (mean diff=10.9 lb (4.9 kg); 95% CI=0.22-21.6; $t=2.24$, $df=62$, $p<0.05$). Fifteen of the women who were high scorers on the EAT, all of whom were within the normal weight range, considered themselves to be very fit, whereas none of the women in the comparison group did (diff=25%, 95% CI=9-41; Yates' $\chi^2=7.65$, $df=1$, $p<0.01$). Excessive exercising was not a common method of weight control in any of the diagnostic groups.

TABLE 111—Physical characteristics of each diagnostic group (women)

	Mean height (in)	Mean weight (lb)	Mean BMI ²	Mean difference between high and low scorers (lb)
Bulimia	159.7	64.9	22.1	1.20
Partial syndrome	158.1	64.9	25.3	1.20
Obsessive compulsive disorder	158.1	64.9	25.3	1.20
Normal eating	157.9	64.9	25.3	1.20
Obsessive	158.1	64.9	25.3	1.20
Disordered	158.1	64.9	25.3	1.20
All high scorers on EAT	158.1	64.9	25.3	1.20
Comparison group	158.1	64.9	25.3	1.20

*BMI=body mass index = weight (kg)/height (m)².

Based on 11 subjects who were within the normal weight range in the past.

Comparison group = 10 men and 35 women.

Menstrual function—A high score on the EAT was not associated with current menstrual cycle irregularity. More subjects with bulimia and partial syndrome (6/21) reported amenorrhoea associated with weight loss in the past than other high scorers on the EAT (3/43) (diff=22%, 95% CI=4-40; $\chi^2=5.458$, $df=1$, $p<0.05$), although they did not differ significantly from the comparison group (4/32).

Past history and family psychiatric history—There was a trend for high scorers on the EAT to report more "nervous trouble" or psychiatric contact in the past than low scorers. Although subjects with bulimia and partial syndrome, were no more likely to have had past nervous trouble than other high scorers, there was a trend for more of those with bulimia to report that first degree relatives had received psychiatric treatment.

Current psychiatric status—Subjects with bulimia scored highly on the symptom rating test and the clinical interview schedule (table IV). Subjects with bulimia and partial syndrome scored higher on the clinical interview schedule than comparison subjects with bulimia (diff=7.15, $t=5.17$, $df=36$, $p<0.001$ and mean diff=8.8, 95% CI=2.8-15, $t=5.28$, $df=45$, $p<0.001$) or normal dieters (mean diff=8.5, 95% CI=2.15, $t=5.28$, $df=27$, $p<0.05$ and mean diff=8.3, 95% CI=2.15, $t=5.28$, $df=36$, $p<0.05$). The disturbed group, however, were the highest scorers, indicating their level of psychological distress. High scorers on the EAT were significantly more likely than comparison subjects to score over 12 on the clinical interview schedule, a cutoff used for research purposes, or over 17, a level of "caseness" seen in psychiatric outpatients (table V). Twenty five of the 64 high scorers were assigned a score according to the International Classification of Diseases,¹² in addition to the eating disorder where present.

TABLE IV—Mean scores on the three tests (women)

Diagnosis	EAT	Clinical interview schedule	Symptom rating test
Bulimia	35.7	16.5	41.0
Partial syndrome	32.1	14.3	34.3
Obsessive compulsive disorder	28.3	14.2	31.3
Normal eating	25.0	14.2	31.3
Obsessive	35.5	14.2	31.3
Disordered	35.5	14.2	31.3
Comparison group	25.7	12.3	28.7

Minimum score on symptom rating test=30.

Receptionists are in the front line. More of them should be encouraged to do that line.

Back and call

"What is a good GP?" Discuss, referring to personal experiences. A good GP doesn't talk down to all his patients as a person. He doesn't adopt a lowest common denominator policy and treat every patient at that level. He doesn't stand on a pedestal with his patients clustered below him on the plinth. He talks to his patients—and he listens. He doesn't do over prescriptions in lieu of thoughtful advice. He does ask questions and give answers, and he does know you when he sees you.

We all have demands which doctors might consider unreasonable, of course. What we all want, particularly people like me—senior, part time depressive mums—are doctors who are at our beck and call, and that of our children, and who really do perform above and beyond the call of duty. From my personal experience, such doctors do exist.

But on the whole do we expect too much of our GPs? Certainly we expect, and should get, a more personal service than we get from specialists. One of these, an elderly paediatrician, repeatedly

referred to my naked baby daughter as "the" throughout a consultation. We went out of the house not to be like that but to be part of the family and to do for us what we set fit.

An acquaintance of mine begged for a home confinement and was rejected by her GP because of her age. She batted throughout the pregnancy to find some other GP to take on the delivery. She was still battling when she went into early labour, at home with her husband as stand in for both GP and midwife. She had a healthy daughter.

But although we perhaps don't get such a personal service as we might like, many general practices are now extremely worthy—well in line I should think for good practice allowances. On the face of it they have exceptionally high standards. At some practices nurses, health visitors, and midwives all operate under the same roof as the doctor. You can get your child vaccinated; attend ante-natal classes; have smear tests and breast checks; get advice on giving up smoking or losing weight; and take part in small research projects and medical questionnaires.

You can also be at the mercy of medical referral agencies at night; you can be kept away from your own practice; you can perhaps see him occasionally if your illness descends at a convenient time. It's computerised, high tech, general practice. It's good practice—but who for?

Practice Research

Eating disorders in general practice

MICHAEL B KING

Abstract

A total of 748 patients who attended four south London group practices were screened using the eating attitudes test; 1% of women had bulimia nervosa and a further 3% a partial syndrome eating disorder. Eating and weight control behaviour and psychiatric indicators for an eating disorder were analysed. Patients with bulimia nervosa and partial syndromes were remarkably similar. They were mainly women, from the middle to upper classes, in the normal weight range but having had considerable weight fluctuation in the past, more likely to have had a history of menstrual irregularity, often psychologically troubled, and tended to have more family psychopathology.

Introduction

The epidemiology of eating disorders has suffered from a problem common to many other psychiatric disorders, whereby concepts of "caseness" appropriate to specialist settings have been applied to general populations. This has obscured our understanding of the range of the disorders and their natural history.¹

Little is known of the role of the general practitioner in recognising and managing patients with eating disorders. Despite the growing number of publications on anorexia and bulimia

nervosa, most have concentrated on hospital studies or special populations.² In the community distinctions between those who are affected and those who are not affected become less clear, but it seems that partial syndromes are much more prevalent than the "classical" cases seen in hospital clinics.³ Partial syndromes share features typical of anorexia and bulimia nervosa,⁴ but the degree of weight loss or amenorrhoea is less pronounced.⁵ Intervention at an earlier stage might be more successful than that reported from hospital clinics.

The aims of this study were to: (a) estimate the prevalence and range of eating disorders among general practice attenders and (b) investigate the clinical profile of eating disorders in the community.

Method

Consecutive patients aged 16 to 35 who attended four south London group practices were asked to complete the 26 item eating attitudes test (EAT)⁶ and supply demographic information, including details concerning height and weight. Women were also asked about their menstrual cycle.

All high scorers on the EAT (cutoff=19) and a random selection of low scorers were asked to take part in a semistructured interview consisting of: (a) the clinical interview schedule, which was constructed for the assessment of non-psychotic disorders in the community;⁷ and (b) the eating interview, which is a detailed format to establish the clinical diagnosis of an eating disorder.⁸ Subjects then completed the symptom rating test, which is a 30 item questionnaire concerning psychological symptoms in the preceding week.⁹ All subjects were weighed at interview. All high scorers were followed up after 12 months to assess clinical change and intervention by the general practitioner.

in contrast to only two of the 32 subjects in the comparison group (mean diff=13%, 95% CI=1.5-24.5; Yates' $\chi^2=9.8$, $df=1$, $p<0.01$). Diagnoses were mainly neurotic depression (which in the classification includes mixed states of anxiety and depression) or depressive reaction.

TABLE V—Psychiatric caseness on the clinical interview schedule (women)

	No.	Cutoff=12	Cutoff=17
Bulimia	6	4 (66%)	3 (50%)
Partial syndrome	13	8 (61%)	6 (46%)
Obsessive compulsive disorder	13	5 (38%)	1 (8%)
Normal eating	23	4 (17%)	3 (13%)
Obsessive	4	1 (25%)	1 (25%)
Disordered	4	1 (25%)	1 (25%)
Comparison group	32	2 (6%)	1 (3%)
All high scorers on EAT	66	27 (41%)	21 (31%)

*High scorers on EAT: diff=36%, 95% CI=17.35-54.3, $p<0.001$.

Comparison group: diff=14.16%, 95% CI=4.16-24.16, $p<0.01$.

Discussion

Four London group practices cannot be considered to be representative of primary care throughout the United Kingdom nor can patients who attend these practices be seen to be representative of all practice populations or of the community. The high proportion of women who attended is in keeping with general practice attenders in this age range.¹⁰

The EAT has been shown to be sensitive and specific in this population of attenders at general practice, albeit at the expense of a large proportion of false positives. The 13.3% of women who scored above the cutoff for the EAT is higher than that reported in other UK studies.¹¹ Six women and one man had bulimia nervosa of sufficient severity for outpatient treatment. A female prevalence of 1.1% for bulimia and 2.8% for partial syndrome agrees with the results in studies of schoolgirl populations¹² and a self report community study in the UK in which there was a point prevalence of 1.9% for bulimia nervosa.¹³ Thus these disorders are uncommon but by no means rare.

In the only other study reported from general practice screening was restricted to women aged 18 to 22 years, and a relatively poor response rate to both screening and subsequent interview was obtained.¹⁴ One case of bulimia nervosa was found directly from screening, and although partial syndromes were reported, few details of their prevalence or clinical presentation were given.

All subjects with bulimia and partial syndrome were of middle to high social class. This has been widely reported for Britain¹⁵ and North America.¹⁶

There appears to be considerable flux in these disorders. As many as three of the seven subjects with bulimia may have had anorexia nervosa in the past, a slightly higher proportion than that in other reports.¹⁷ In addition, five of the subjects with partial syndrome appeared to have suffered the full syndrome in the past, and up to a third of the obsessional dieters may previously have had at least a partial syndrome disorder. In other studies it has been shown that even subjects with a clear diagnosis of anorexia nervosa may show considerable improvement without intervention in as little as 12 months.¹⁸

Although of normal weight, two thirds of the subjects with bulimia and partial syndrome considered themselves to be overweight. Despite this, their target weights were not below population norms, and thus they were not seeking extreme thinness. Menstrual irregularities were not a helpful guide in the two groups, although these have been reported for bulimia nervosa.¹⁹ There was a link, however, with past amenorrhoea associated with weight loss, suggesting that menstrual function might be most closely related to change of weight rather than to disordered eating habits per se.

A high score on the EAT has been linked with the reporting of neurotic symptoms,²⁰ but, as shown here by the small group of "disturbed" subjects, this may be unrelated to a primary concern about diet or weight. Subjects with bulimia and partial syndrome

and obsessional dieters were all more likely to have psychological problems, which has also been reported for 2.8% of partial syndrome eating disorder. The considerable changes in the severity of the disorders suggest that intervention by the general practitioner would be most appropriate when a patient begins obsessional dieting or at the partial syndrome stage when symptoms may be less entrenched.

"Caseness" on the clinical interview schedule or a diagnosis based on the International Classification of Diseases was associated with at least a partial syndrome eating disorder. The role of psychological symptoms in the range or evolution of eating disorders is not clear, although most of the subjects with bulimia and partial syndrome were viewed by their distress as secondary to their eating behaviour.

Five (2.6%) of the 184 men screened scored above the cutoff of the EAT, considerably more than in other UK series.¹¹ Only one (0.5%) man, however, had an eating disorder—a similar rate to that in a self report study in the UK¹³ but unfortunately little can be concluded from the clinical profile of one individual.

Of the subjects with bulimia, one woman and one man were of Afro-Caribbean racial origin, which adds to the evidence that eating disorders appear in first or second generation offspring of immigrants to first world countries.²¹ It may be that eating disorders in non-white races are less readily recognised and are under-represented in hospital series.

The findings showed that 1.1% of women who attended their general practitioners had bulimia nervosa and 2.8% a partial syndrome eating disorder. The considerable changes in the severity of the disorders suggest that intervention by the general practitioner would be most appropriate when a patient begins obsessional dieting or at the partial syndrome stage when symptoms may be less entrenched.

I thank Dr Paul Williams and Professor Michael Shepherd, Institute of Psychiatry, for their helpful criticism and advice, and particularly Dr Deborah Sharp, Institute of Psychiatry, for suggestions and support.

References

- Williams P, Topley H, et al. A case of anorexia nervosa: a psychiatric epidemiological survey. *Psychol Med* 1980;10:161-164.
- Smolkin G. The epidemiology of anorexia nervosa and bulimia. *J Psychol Res* 1982;19:143-53.
- Russell G. The changing nature of anorexia nervosa: an introduction to the conference. *J Psychol Res* 1982;19:1-11.
- Burns EJ, Whitman A. Subclinical anorexia nervosa. *Psychol Med* 1981;11:309-16.
- Smolkin G. Prevalence and prevalence in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.
- Smolkin G, Whitman A, Wood R, Moore R, Davies A, Smolkin G. Screening for anorexia nervosa and bulimia in a population of English schoolgirls. *Report of the International Study on Anorexia Nervosa and Bulimia*. London: Taylor & Francis, 1981:17-38.

(Accepted 17 September 1986)

General Practice Research Unit, Institute of Psychiatry, Denmark Hill, London SE5 8AF.
MICHAEL B KING, MRCP, MRCPsych, honorary lecturer