transmission by any type of penetrative sex. The results of this study may be used, however, to help assess the degree of risk for those who find celibacy unacceptable.

Heterosexuals may be less efficient at transmitting HIV, but the virus can certainly pass from men to women and from women to men during vaginal intercourse.³ And whatever their sexual orientation, those who have multiple sexual partners not only increase their chances of encountering HIV but also are more likely to acquire other sexually transmitted diseases, which may facilitate HIV transmission.

Despite worries of a heterosexual epidemic there is little evidence of this happening yet outside Africa. In the USA 96% of cases of AIDS belong to recognised risk groups—a figure that has remained constant for years.⁴ In Britain the Public Health Laboratory Service in its national surveillance of HIV antibody testing has not noted an outbreak in those without risk factors (McAvoy M, personal communication), and the low prevalence of antibodies to HIV in blood donors is further evidence against an impending epidemic. One explanation for these observations is that people choose sexual partners from their own social circle with very little mixing. This sexual insularity will tend to slow the advance of HIV.

HIV may well become endemic outside the current risk groups, but this could take years and might never be extensive. The question thus arises whether large sums of money earmarked for an advertising campaign might be better spent on improving treatment facilities or on basic research. An education campaign aimed at preventing the spread of HIV is, of course, to be welcomed provided that unnecessary fear is not provoked and that this rare disease is kept in perspective. Certainly, one could argue for more emphasis on the very low infectivity of the virus under normal circumstances and on the many ways by which it cannot be spread.

The gay community is well informed, and homosexual lifestyle has already changed.⁵ Many of those who are infected or at high risk practise the ideal of monogamy with "safe sex," which implies "dry kissing" and either mutual masturbation or the careful and consistent use of condoms. Education needs to be aimed at highly sexually active heterosexuals with a message neither dogmatic nor moralistic. Those outside high risk groups should be advised to use condoms with all casual partners and to reduce the number of partners as far as possible.

Evaluation of these educational measures may be difficult, but an alteration in sexual lifestyle should have a beneficial effect on other important sexually associated conditions such as tubal infertility and cervical cancer, which are far more common than AIDS.

CAROLINE BRADBEER

Senior Registrar,
Department of Genitourinary Medicine,
St Thomas's Hospital,
London SE1 7EH

- 1 Lifson AR, Ancell RA, Brunet JB, et al. The epidemiology of AIDS worldwide. *Clinics in Immunology and Allergy* 1986;6.
- 2 Evans BA, Dawson SG, McLean KA, et al. Sexual lifestyle and clinical findings related to HTLV III/LAV status in homosexual men. *Genitourinary Medicine* 1986;62:386-91.

- 3 Calabres LH, Gopalakrishna KV. Transmission of HTLV-III infection from man to woman to man. *N Engl J Med* 1986;318:987.
- 4 Anonymous. Heterosexual transmission of human T-cell lymphotropic virus type III/lymphadenopathy-associated virus. *MMWR* 1985;34:561-3.
- 5 Burton SW, Burn SB, Harvey D, Mason M, McKerrow GM. AIDS information. *Lancet* 1986;ii:1040-1.

AIDS: a doctor's duty

Caring for the sick has always been a hazardous occupation. Even as recently as the first 40 years of this century doctors and nurses working with patients with diseases such as typhus or tuberculosis knew they risked infection which could be fatal. Pathologists risked death if they cut themselves in the necropsy room, and until the 1980s—with the introduction of an effective vaccine—medical and nursing staff regularly caught hepatitis B and sometimes were killed by it.

Against that background, how should doctors and nurses be responding to the rising numbers of patients infected with the human immunodeficiency virus (HIV)? All the evidence points to the risks being extremely small. HIV seems much less infectious than the hepatitis B virus, for example, and no health professional anywhere in the world is known to have contracted the acquired immune deficiency syndrome (AIDS) nosocomially (though a very few have developed antibodies to HIV after needlestick injuries).¹

Those doctors who are calling for patients to have antibody tests before they undergo surgical procedures—and those who refuse to treat patients who have antibodies—are contributing to current hysteria about the disease. The antibiotic era has changed public attitudes to infections; this has been evident in the overreaction by the popular media to "new" diseases such as legionnaires' disease and Lassa fever.² Doctors ought to have led public opinion on those occasions, but—sadly—some doctors contributed to the atmosphere of fear and isolationism. The same pattern is now being seen with AIDS; anyone who has kept abreast of reports in even a single journal should be familiar with the risks and the reasonable precautions, but some doctors are still making comments suggesting that they know little about the disease.

In simple terms AIDS and HIV carriage do not represent a substantial health risk for doctors; if they did there would have been reports of medical infection from the United States, where around two million people have antibodies to HIV. The Royal College of Nursing has had the courage to recommend the disciplining of any nurse who refuses to care for a patient infected with HIV.³ Should not the General Medical Council do the same?

TONY SMITH

Deputy Editor, *BMJ*

- 1 Geddes AM. Risks of AIDS to health care workers. *Br Med J* 1986;292:711-2.
- 2 Tilzey AJ, Webster M, Banatvala JE. Patients with suspected Lassa fever in London during 1984: problems in their management at St Thomas's Hospital. *Br Med J* 1985;291:1554-5.
- 3 Royal College of Nursing Working Party on AIDS. *Nursing guidelines on the management of patients in hospital and the community suffering from AIDS*. London: Royal College of Nursing, 1986.