could find no other major social or environmental change that occurred in the population we served which might have explained these striking findings.

We agree with Dr Platt that it would have been advantageous to compare our experience with that of similar districts in England. Despite strenuous efforts, however, we could not obtain comparable data of this type. The experience in the Edinburgh poisons unit has apparently been different from our own, but several factors may explain this. In contrast to our own policy during the period of study, we understand that in Edinburgh, particularly during the latter years of the review described by Dr Platt, the admission policy has changed in that only some of the patients reaching the accident and emergency department are admitted to the poisoning centre. Also, from the data presented, it is clear that the percentage of Edinburgh patients who took alcohol with their overdose before the change in the licensing laws was more than twice that in our own series. This may well have obscured and at least reduced any impact which the change in the law may have caused in the Edinburgh admission rate. As far as poisoning is concerned, it is well established that patients from capital cities and other dense conurbations behave differently from those in a mixed urban and rural community.

We would suggest, therefore, that the data presented by Dr Platt are not necessarily strictly comparable with our own. Clearly it is difficult to provide any proof of the association which we suggested and we would be most interested to learn of others' experience, particularly from comparable district general hospitals.

A LAWSON
JOHN McMURRAY
D B NORTHIDGE

Milesmark Hospital,
Rumblingwell,
Dunfermline KY12 9NR

simply consist of part trained consultants ill supported by resources (secretary, office, junior staff) who will be expected to do the real consultants' night duty for them?

(7) Has anyone considered the wisdom of a system which effectively abolishes all competition above SHO grade? Future consultants will be chosen within two or three years of graduation, long before their suitability for such posts can be assessed.

(8) Does the BMA understand the problems implicit in a reduction in clinical research? The distinction drawn between training in research methods and in clinical research is nonsense. To suggest that a research registrar should be drawn from the overall quota of acute service registrars is also nonsense—unless there are to be more registrars allocated than there is clinical work to be done.

In summary, we are deeply disturbed about both the fundamental principles and the details of this report. We believe the solutions proposed for the problems of the hospital medical career structure are ill considered. We suggest, moreover, that the whole problem has been approached in the wrong way. Matching the number of registrar posts to the expected number of consultant vacancies is a very blinkered way of planning medical staffing of the health service. How much better to begin by matching the numbers of the different grades of staff to the work to be done. Then we might know whether the system could support a pyramidal career structure without undue wastage or whether a wholesale change in this historic pattern might be better.

Achieving a Balance may solve some of the problems of medical career structure but as a package deal we believe it has so many flaws that it would be best to reject it completely and to have no deal at all.

A MORGAN and W JEFFCOATE
and 104 other doctors

City, General, and University Hospitals,
Nottingham

received any steroids. This cannot therefore be said to constitute a trial of steroid therapy in an unselected series of patients.

The other papers³⁻⁵ quoted by Mr Thomas and others and by Dr C E Fadul and others (p 1015) concern randomisation of patients who had survived major surgery for two to four weeks and who were "fit enough" to withstand radiotherapy or chemotherapy. The criteria for major surgery were not given and we do not know what happened to those who did not undergo major resection. The patients admitted to these surgical trials were, as Dr Fadul and colleagues acknowledge, a different cohort from those assessed in our clinical audit. Each trial showed that radiotherapy did prolong life in these selected cases and that any additional effect of chemotherapy was marginal. Indeed, each paper pleaded for a more effective management regimen.

We would emphasise again that our report was a clinical audit of an unselected group of patients presenting consecutively to different consultants in a regional neurosciences unit. Many of the patients would not have been considered suitable for entry into the surgical series discussed by your correspondents. This does not mean that we refuse to pursue an aggressive management policy in those patients in whom this is appropriate. Unfortunately such patients represent an all too small proportion of the population presenting to us. We will, as we have in the past, continue to join multicentre trials, usually in conjunction with our colleagues in radiotherapy and oncology.

P M FOY
M D M SHAW
I R WILLIAMS
D W CHADWICK

Mersey Regional Department of
Medical and Surgical Neurology,
Walton Hospital,
Liverpool L9 1AE

Manpower

SIR,—We recently received the communication from the chairmen of the Central Committee for Hospital Medical Services and the Hospital Junior Staff Committee about hospital medical staffing. We are worried about the implications of the report *Achieving a Balance* and feel that agreements have been reached on our behalf with which we are not in accord. The consultation process adopted by the BMA leadership has not successfully informed the leaders of what their members think.

The circular attempting to explain *Achieving a Balance* leaves many questions unanswered.

(1) Why is there such a need to rush into restructuring junior hospital doctors' jobs?

(2) How can expansion of the consultant grade be accommodated "within existing facilities"? How can a surgeon operate without theatre space or without an anaesthetist? How can a physician run a clinic when there is no clinic space?

(3) What is the good of simply shifting the bottleneck in junior posts down to senior house officer grade? The competition for hospital SHO posts and for vocational training scheme posts is already bad enough.

(4) The BMA has not explained how a reduction in quality of patient care can be avoided in those districts in which there is a reduction in numbers not only of registrars but also of SHOs and senior registrars. In one paragraph the crucial service role of the registrar is acknowledged, yet the basis of this report is that the numbers of registrar posts should be reduced.

(5) How is it possible to avoid creating two types of registrar who provide different levels of care? District registrar posts will be essential for service needs, but is it right for the National Health Service to depend on a supply of foreign graduates on short term contracts?

(6) Will not the new intermediate service grade

Neurological and neurosurgical approaches in the management of malignant brain tumours

SIR,—In our original paper (18 October, p 1015) we perhaps did not make it clear that the 18 cases in the text referred only to those of a proved glioma (excluding ependymoma) for the diagnosis of malignant glioma was certainly not proved in the remaining cases, which were also recorded in table V. Certainly the great majority of patients submitted to radiotherapy had undergone surgery, but in most cases this was a burr hole biopsy alone. There is no evidence to support the contention that patients who did not undergo surgery included a greater proportion with low grade tumours or no tumours than the group who underwent biopsy. Indeed, as judged by the other criteria the groups were broadly similar.

Mr D G T Thomas and others (10 January, p 123) correctly quote from the discussion in Salcman's paper,¹ but the data in that paper showed that the median survival in glioblastoma multiforme was 4 months with operation alone and 9.25 months when radiotherapy was added. Whether this represents an improvement other than marginal, once time has been allowed for treatment and its effects, is a matter of opinion.

Mr Thomas and his colleagues quote the trial of Walker *et al*² as one in which steroids alone were used in one arm. Ninety five per cent of patients admitted to that trial had undergone a major resection, and all had to have a projected survival of at least two months postoperatively and to have received steroids for less than one month before they could be admitted to one of the four arms of that trial. Only 76% of the patients actually

- 1 Salcman M. Survival in glioblastoma multiforme: historical perspective. *Neurosurgery* 1980;7:435-9.
- 2 Walker MD, Alexander E Jr, Hunt WE, *et al*. Evaluation of BCNU and/or radiotherapy in the treatment of anaplastic gliomas. *J Neurosurg* 1978;49:333-43.
- 3 Walker MD, Green SB, Byar DP, *et al*. Randomized comparisons of radiotherapy and nitrosoureas for the treatment of malignant glioma after surgery. *N Engl J Med* 1980;303:1323-9.
- 4 Kristiansen K, Hagen S, Kollevold T, *et al*. Combined modality therapy of operated astrocytomas grade III and IV. Confirmation of the value of postoperative irradiation and lack of potentiation of bleomycin on survival time: a prospective multicenter trial of the Scandinavian Glioblastoma Study Group. *Cancer* 1981;47:649-52.
- 5 Green SB, Byar DP, Walker MD, *et al*. Comparisons of carmustine, procarbazine and high-dose methylprednisolone as additions to surgery and radiotherapy for the treatment of malignant glioma. *Cancer Treat Rep* 1983;67:121-32.

* This correspondence is now closed.—ED, BMJ.

Dialysis arthropathy: amyloid or iron?

SIR,—We investigated 10 patients undergoing regular haemodialysis treatment (duration 4-18 years) who underwent a carpal tunnel release operation (bilateral in two patients). The synovial tissue was histologically examined for iron, aluminium, amyloid, and β_2 microglobulin. Of these eight had amyloid and their synovial tissue stained positively for β_2 microglobulin using an immunoperoxidase method. Aluminium was not detected in any of these specimens, and only one patient had deposits of haemosiderin but her serum ferritin concentration was 39 $\mu\text{g/l}$. Another patient who was anephric had received 130 units of blood transfusions; her serum ferritin concentration was 8030 $\mu\text{g/l}$ and she showed no evidence of iron in the synovial tissue. However, in this patient excess iron was present at the osteoid mineralisation