- surveillance of AIDS in health care workers. *JAMA* 1986; 256:3231-4.
- 11 Fultz PN. Components of saliva inactivate human immunodeficiency virus. *Lancet* 1986;ii:1215.
 - 12 Groopman JE, Salahuddin SZ, Sarngadharan MG, et al. HTLV-III in saliva of people with AIDS related complex and healthy homosexual men at risk for AIDS. *Science* 1984;226: 447-9.
 - 13 Ho DD, Byington RE, Schooley RT, et al. Infrequency of isolation of HTLV-III virus from saliva in AIDS. *N Engl J Med* 1985;1606.
 - 14 Zagury D, Bernard J, Leibowitch J, et al. HTLV-III in cells cultured from semen of two patients with AIDS. *Science* 1984;222:449-51.
 - 15 Ho DD, Schooley RT, Rota TR. HTLV-III in the semen and blood of a healthy homosexual man. *Science* 1984; 226: 451-3.
 - 16 Lifson AR, Ancelle RA, Brunet JB, Curran JW. The epidemiology of AIDS worldwide. In: Pinching AJ, ed. *Clinics in immunology and allergy*. Vol 6. London: W B Saunders, 1986:441-65.

State of the public health

SIR,—The suspension of Dr David Josephs by South Bedfordshire Health Authority for his remarks about the action needed to contain the epidemic of the acquired immune deficiency syndrome contrasts sharply with the freedom of the chief constable of Manchester to expound on the same subject. Taken together, these two examples provide a dramatic insight into the state of public health in Britain.

In the 1840s and 1850s there was widespread debate in Europe about the need for "medical police" to control public health hazards. Such an authoritarian style was never imposed, but in Britain we developed a systematic approach to the protection of public health based on technical skill and scientific knowledge rather than popular superstition about disease. This skill was housed in

the office of the medical officer of health in the local authority. Starting with Duncan of Liverpool in 1847, it came to be accepted that the medical officer of health could not be sacked for speaking out on contentious matters that might upset politicians or vested interests; he could be sacked only for professional incompetence. His powers were reinforced by the Public Health Act 1848.

Since the 1974 NHS and local government reorganisations this has changed, and the independent advocacy function of public health has been all but lost. Sadly, this has often occurred with the apparent collusion of community physicians, who have inherited the mantle of the medical officer. In Britain today the public health function is going by default; those who should be making their voices heard on behalf of the public health are all too often keeping their heads down in self imposed silence. Those few who are brave enough to speak out find themselves reprimanded or, like Dr Josephs, actually suspended.

So while a self proclaimed prophet trained as a policeman is free to offer public advice on public health problems a doctor with technical skill in public health who voices a legitimate opinion finds his livelihood threatened. David Player already looks set to be sacrificed for speaking out against the pedlars of ill health. We are certainly back in Victorian Britain with a vengeance (pre-1848) and increasingly seem simultaneously stuck in 1984; rationality seems to have bowed down to superstition on the one hand and intolerance and authoritarianism on the other. Surely it is time to stand up and be counted.

J ASHTON

Department of Community Health,
University of Liverpool,
Liverpool L69 3BX

Points

An assessment of the preregistration year experience

Dr R F Gledhill (Department of Internal Medicine, Kalafong Hospital, Pretoria 0001, South Africa) writes: Drs J E Elizabeth and S Hughes (13 December, p 1559) may have been too ready to accept the veracity of the self assessments provided by their Liverpool preregistration graduates. The known limitations of this evaluation method^{1,2} were strikingly illustrated in a report from the Mount Sinai School of Medicine,³ in which 78% of subjects (third and fourth year students and residents) felt confident in their ability to perform cardiopulmonary resuscitation, but only 2.9% could execute this procedure correctly on examination. In contrast to the figure of more than 95% at Liverpool, an average of only 37% to 75% of Cambridge University graduands felt competent to manage a seizure and acute presentations of left ventricular failure, asthma, and gastrointestinal bleeding.⁴ Furthermore, whereas two thirds of the Liverpool house officers considered themselves competent in performing a lumbar puncture and abdominal paracentesis, the figures for Cambridge were 30% and 12% respectively.⁴ With respect to medical emergencies the findings reported by Wakeford and Roberts⁴ accord with those obtained in surveys of graduates from McMaster University⁵ and interns from the University of Stellenbosch.⁶ Liverpool University Medical School may well conduct highly effective undergraduate and postgraduate programmes but there would also seem good reason to question whether the results of the survey are especially impressive because some of their respondents tended to overestimate their abilities.

- 1 Kaisen A, Kjsets GA, Lie RK, et al. Interns' evaluation of their preparation for general practice: a comparison between the University of Tromsø and the University of Bergen. *Med Educ* 1984;18:349-54.
- 2 Scabassi SE, Woelfel SK. Development of self assessment skills in medical students. *Med Educ* 1984;18:226-31.

- 3 Nelson M. Evaluation of CPR performance among medical students, residents and attendings at Mount Sinai School of Medicine. *Mt Sinai J Med* 1981;48:89-94.
- 4 Wakeford R, Roberts S. An evaluation of medical students' practical experience upon qualification. *Medical Teacher* 1982; 4:140-3.
- 5 Woodward CA, Ferrier BM. The content of the medical curriculum at McMaster University: graduates' evaluation of their preparation for postgraduate training. *Med Educ* 1983; 17:54-60.
- 6 Gledhill RF. Undergraduate clinical neurosciences programme development: A consumer based evaluation. *J R Soc Med* (in press).

Man bites dog

Mr K HASHEMI (Accident and Emergency Service, Mayday Hospital, Thornton Heath, Surrey CR4 7YE) writes: I agree with Drs B Kirkpatrick and R Wise (13 December, p 1522) that potentially the most serious form of human bites and the ones most likely to cause complications are clenched fist injuries. These injuries are more prone to complications because doctors often fail to recognise the injury as caused by a human bite. This, coupled with patients' own reluctance to give a clear history of the cause of their injury, leads to delay in providing appropriate treatment. The only sure way of avoiding this problem is to treat every "knuckle" injury as a clenched fist injury. Although an aggressive approach is necessary, I do not, however, agree that inpatient treatment is usually needed. In my own practice exploration and irrigation of the wound with any necessary debridement are carried out at the time of first presentation. Swabs and irrigation samples are sent for bacteriological culture. The open wound is then dressed and the injured hand raised in a high arm sling. The patient is started on a course of antibiotics and my own personal choice is Augmentin (amoxycillin and clavulanic acid). Every patient is reviewed the next day, and those who show evidence of spreading cellulitis are given a volar plaster cast immobilising the injured hand from the finger tip to the point of their

elbow, and the hand is again raised in a high arm sling, and these patients are reviewed daily. Those who do not show evidence of spreading cellulitis are reviewed at lengthened intervals. I have found the above regimen extremely effective, and since its introduction more than three years ago there has been no case of serious complications or inpatient treatment.

Attitudes to prescribing iron supplements in general practice

Dr J J BANNISTER (Department of Surgery, Royal Hallamshire Hospital, Sheffield S10 2JF) writes: In the survey of Drs D G Waller and A G Smith most GPs usually prescribed iron on the basis of the full blood count (93%) or a blood film suggesting iron deficiency (81%), apparently without further investigation. The study does not delve into the reasons for the subjects being iron deficient. It should be emphasised that iron deficiency anaemia is not a disease in its own right, to be treated with iron supplements, but merely a sign of an imbalance between, on the one hand, the dietary intake and absorption of iron and, on the other hand, the body's losses of iron. In all patients, with the possible exception of women with heavy menstrual loss or the loss of iron associated with pregnancy, the finding of a microcytic and hypochromic anaemia should alert the attending doctor to the possibility of chronic bleeding from the gastrointestinal tract. Testing for the presence of occult blood in the faeces may be useful in confirming bleeding, but a full and prompt radiographic or endoscopic study of the upper and lower bowel is the only way in which symptomless and possibly remediable gastrointestinal lesions, such as caecal carcinoma or colonic polyps, will be detected.

Self help groups

Dr ELIZABETH BRYAN and Ms MARGARET HARRISON (Home-Start Consultancy, Leicester LE1 7JL) write: Many readers working with self help groups will have been delighted to see Dr Stephen Lock's valuable article about them (20-27 December, p 1596). But does not their role in preventive medicine deserve as much emphasis as that of supporting patients (and their families) suffering disease or disability? If an overstressed mother can be prevented from abusing her child or needing antidepressant treatment through the committed support of another mother there can be little doubt of the cost effectiveness, let alone human benefit. In recent years this has been clearly demonstrated by Home-Start, a voluntary scheme in which volunteers offer practical support and friendship to families with children under 5 in their own homes. There are now over 60 local Home-Start schemes. The role of the volunteer is, of course, complementary to that of the professional worker, and the organiser of each scheme always has close links with the statutory services.

Depression after stroke

Dr DAVID AMES (Academic Department of Psychiatry, Royal Free Hospital, London NW3 2QG) writes: Dr Allan House's otherwise excellent review (10 January, p 76) contains a common but incorrect assumption about the uses to which Hamilton's depression rating scale¹ may justifiably be put. Hamilton's depression rating scale is intended not as a diagnostic instrument, as Dr House seems to believe, but specifically to rate the level of depression in patients already diagnosed as depressed. Hamilton makes this point clear in his original paper. For patients over 65 with stroke the most useful standardised diagnostic interview is probably the geriatric mental state of Copeland *et al.*² This schedule is better suited to the diagnostic requirements of an elderly population than is the present state examination.

- 1 Hamilton MA. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56-62.
- 2 Copeland JRM, Kelleher MJ, Kellett JM, et al. A semi-structured clinical interview for the assessment of diagnosis and mental state in the elderly: the geriatric mental state schedule. *Psychol Med* 1976;6:433-49, 51-9.