another large study as 1 in 175 patient years.¹⁶ For children and young adult patients this may be considered a high enough risk to justify antireflux surgery in the hope that the risk is thereby reduced, but it is not established that the benefits of surgery outweigh its hazards or that regular endoscopic surveillance allows usefully early recognition of carcinoma. These issues can and should be addressed in prospective trials. For now the evidence justifies a more conservative policy directed at relieving symptoms and healing oesophageal ulcers.

R C HEADING

Senior Lecturer,
University of Edinburgh,
Department of Medicine,
Royal Infirmary,
Edinburgh EH3 9YW

- 1 Barrett NR. Chronic peptic ulcer of the oesophagus and oesophagitis. *Br J Surg* 1950;38:175-82.
- 2 Spechler SJ, Goyal RK. Barrett's esophagus. *N Engl J Med* 1986;315:362-71.
- 3 Hassall E, Weinstein WM, Ament ME. Barrett's esophagus in childhood. *Gastroenterology* 1985;89:1331-7.
- 4 Paul A, Trier JS, Dalton MD, Camp RC, Loeb P, Goyal RK. The histologic spectrum of Barrett's esophagus. *N Engl J Med* 1976;295:476-80.
- 5 Peuchmaur M, Potet F, Goldfain D. Mucin histochemistry of the columnar epithelium of the oesophagus (Barrett's oesophagus): a prospective biopsy study. *J Clin Pathol* 1984;37:607-10.
- 6 Lee RG. Mucins in Barrett's esophagus: a histochemical study. *Am J Clin Pathol* 1984;81:500-3.
- 7 Herlihy KJ, Orlando RC, Bryson JC, Bozymski EM, Carney CN, Powell DW. Barrett's esophagus: clinical, endoscopic, histologic, manometric and electrical potential difference characteristics. *Gastroenterology* 1984;86:436-43.
- 8 Naef AP, Savary M, Ozzello L. Columnar-lined lower esophagus: an acquired lesion with malignant predisposition. *J Thorac Cardiovasc Surg* 1975;70:826-34.
- 9 Starnes VA, Adkins RB, Ballinger JF, Sawyers JL. Barrett's esophagus. A surgical entity. *Arch Surg* 1984;119:563-6.
- 10 Schnell T, Sontag S, Wanner J, et al. Endoscopic screening for Barrett's esophagus (BE), esophageal adenocarcinoma (Ad Ca) and other mucosal changes in ambulatory subjects with symptomatic gastroesophageal reflux. *Gastroenterology* 1985;88:1576.
- 11 Sarr MG, Hamilton SR, Marrone GC, Cameron JL. Barrett's esophagus: its prevalence and association with adenocarcinoma in patients with symptoms of gastroesophageal reflux. *Am J Surg* 1985;149:187-92.
- 12 Kerlin P, D'Mellow G, Van Deth A. Barrett's esophagus: clinical endoscopic and histologic spectrum in fifty patients. *Aust NZ J Med* 1986;16:198-205.
- 13 Wesdorp ICE, Bartelsman J, Schipper MEI, Tytgat GN. Effect of long-term treatment with cimetidine and antacids in Barrett's oesophagus. *Gut* 1981;22:724-7.
- 14 Kothari T, Mangla JC, Kalra TMS. Barrett's ulcer and treatment with cimetidine. *Arch Intern Med* 1980;140:475-7.
- 15 Skinner DB, Walther BC, Riddell RH, Schmidt H, Jascone C, De Meester TR. Barrett's esophagus. Comparison of benign and malignant cases. *Ann Surg* 1983;198:544-65.
- 16 Spechler SJ, Robbins AH, Rubins HA, et al. Adenocarcinoma and Barrett's esophagus. An overrated risk? *Gastroenterology* 1984;87:927-33.
- 17 Cameron AJ, Ott BJ, Payne WS. The incidence of adenocarcinoma in a columnar-lined (Barrett's) esophagus. *N Engl J Med* 1985;313:857-9.
- 18 Brand DL, Ylvisaker JT, Gelfand M, Pope CE. Regression of columnar esophageal (Barrett's) epithelium after anti-reflux surgery. *N Engl J Med* 1980;302:844-8.

Depression in children

Until recently most doctors have ignored or denied the possibility of depressive disorder in prepubertal children.^{1 2} Those who acknowledged it were either overinclusive¹ or suggested that it was masked by other symptoms,³ thus enabling society and its psychiatrists to continue to disbelieve that children could be depressed. The same happened with infantile sexuality. In the past 10 years research has shown that prepubertal depression is a reality and may continue to become an adult depressive disorder.

Two factors have made these advances possible; firstly, developmental psychopathology has led to an understanding of normal and deviant development of affective expression⁴; and, secondly, operational definitions of psychiatric disorders have been introduced.^{5 6} A depressive disorder may be diagnosed if the child has had a depressed mood for at least four weeks and if he or she has two or more of the following symptoms: anxiety, sleep disturbance, irritability, suicidal thoughts, eating disturbances, school refusal, phobias, ali-

mentary disorders, obsessions, and hypochondriasis. The symptoms must be severe enough to interfere with the child's social life, cognitive function, or both, and represent a change from his or her previous state.^{5 6}

Using these criteria Pearce found that 15-20% of children referred for psychiatric help had a depressive disorder.⁶ General population studies using strict criteria across all age groups are not yet available, but misery and unhappiness are common in childhood (occurring in 10% of 10 year olds in one study).⁷ The same workers found that 1.4% of 10 year olds and 4% of 14 year olds had a depressive disorder.^{7 8} Girls seem more vulnerable than boys at all ages, and depression seems to be more common in preschool children and adolescents than in primary school children. Depression commonly coexists with other syndromes such as conduct disorder and hyperkinesis.

The aetiology of depression in childhood is not yet established and is controversial. Neuroendocrine abnormalities have been found,⁹ and genetic factors are important in manic depressive disorders, which rarely manifest themselves prepubertally, but may also play a part in unipolar depression.¹⁰ Having a depressed mother affects young children's behaviour and learning.¹¹ Adverse life events—particularly loss of a parent—are associated with depression in later life^{12 13} but have not been clearly shown to produce a specific syndrome in prepubertal children. Rather they may be associated with a wide range of behavioural and emotional disorders, including depression.¹⁴ The concept of "learned helplessness"¹⁵ is useful in childhood depression and may apply, for example, in children who have been sexually abused.¹⁶

But neither the "learned helplessness" nor the "adverse life events" hypothesis are compatible with depression being commoner in preschool children and adolescents than in primary school children. The most reasonable hypothesis is one that postulates a three factor model of depression: provoking agents—for example, psychological factors (losses) or physical factors (especially viral infections)—in the presence of vulnerability factors—for example, social adversity—produce depression if the child is predisposed by early experience, especially parental death or genetic or acquired biological factors—for example, brain injury or handicap.¹⁷

Doctors need to distinguish between normal sadness in response to adversity, which should remit and can be helped by counselling, and illness, which requires treatment. Suicidal thoughts and feelings are often present in children but are rarely acted on until puberty. The child may need admission, although most can be managed at home. Psychological methods of treatment—including family therapy, environmental changes, and individual psychotherapy—produce improvement in many cases. Antidepressants are best reserved for severe depression, for where there are clear biological symptoms (sleep and appetite disturbance, diurnal variation in mood, or psychomotor retardation), or for where there is a family history of depression which has responded to drugs. Electroconvulsive therapy is rarely needed in children but can occasionally be lifesaving. Cognitive therapies may be helpful in the older child. Few of these treatments have yet been evaluated in children, but in view of the complex aetiology a pragmatic approach—attempting to promote normal mourning, improve family and school functioning, increase adaptive behaviour, and improve self esteem and using physical treatments where indicated—would appear to be good practice.⁶

Childhood depression may be prevented to some extent by minimising bond disruption and promoting healthy mourning. Young children of depressed mothers may need to be cared for by others while their mothers are receiving energetic treatment, and doctors and others should help children of all ages who are at high risk for depression by eliminating or reducing avoidable provoking and vulnerability factors and strengthening the children's coping behaviours. The psychotherapeutic work of the Tavistock Clinic with severely deprived children appears promising and urgently needs evaluating—as do other approaches.¹⁸

DORA BLACK

Consultant Child and Adolescent Psychiatrist,
Royal Free Hospital,
London NW3 2QG

- 1 Frommer E. Depressive illness in childhood. In: Coppen A, Walk A, eds. *Recent advances in affective disorders*. Br J Psychiatry 1968;117:36. (Special publication No 2.)
- 2 Graham P. Depression in prepubertal children. Dev Med Child Neurol 1974;16:340-9.
- 3 Glaser K. Masked depression in children and adolescents. Am J Psychiatry 1967;21:565-74.
- 4 Rutter M. The developmental psychopathology of depression: issues and perspectives. In: Rutter M, Izard CE, Read PB, eds. *Depression in young people*. New York: Guildford Press, 1986:3-30.
- 5 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders DSM III*. 3rd ed. Washington DC: American Psychiatric Association, 1980.
- 6 Pearce J. Depressive disorder in childhood. J Child Psychol Psychiatry 1977;18:79-84.
- 7 Rutter M, Tizard J, Whitmore K, eds. *Education, health and behaviour*. London: Longman, 1970.
- 8 Rutter M, Graham P, Chadwick O, Yule W. Adolescent turmoil fact or fiction? J Child Psychol Psychiatry 1976;17:35-6.
- 9 Puig-Antich J. Psychological markers: effects of age and puberty. In: Rutter M, Izard CE, Read PB, eds. *Depression in young people*. New York: Guildford Press, 1986:341-81.
- 10 Cytryn L, McKnew DH, Zahn-Waxler C, Gershon ES. In: Rutter M, Izard CE, Read PB, eds. *Depression in young people*. New York: Guildford Press, 1986:163-88.
- 11 Cogill SR, Caplan HI, Alexandra H, Robson KM, Kumar R. Impact of maternal depression on cognitive development of young children. Br Med J 1986;292:1165-7.
- 12 Birchneil J. Early parent death and psychiatric diagnosis. Social Psychiatry 1982;7:202-10.
- 13 Brown GW, Harris TO, Copeland JR. Depression and loss. Br J Psychiatry 1977;130:18.
- 14 Rutter M. *Children of sick parents: an environmental and psychiatric study*. London: Oxford University Press, 1966.
- 15 Seligman MEP, Peterson C. A learned helplessness perspective on childhood depression: theory and research. In: Rutter M, Izard CE, Read PB, eds. *Depression in young people*. New York: Guildford Press, 1986:223-50.
- 16 Anonymous. Childhood depression and sexual abuse [Editorial]. Lancet 1986;i:196.
- 17 Brown GW, Harris TO, Bifulco A. Long term effects of early loss of parent. In: Rutter M, Izard CE, Read PB, eds. *Depression in young people*. New York: Guildford Press, 1986:251-96.
- 18 Boston M, Szur R, eds. *Psychotherapy with severely deprived children*. London: Routledge, 1983.

Bone marrow transplantation in precocious osteopetrosis

Osteopetrosis is a rare disease of bone characterised by a densely sclerotic skeleton and was first described by a radiologist, Albers-Schönberg.¹ Since then many sporadic cases have been reported and collected in useful reviews.^{2,4} Osteopetrosis also occurs in animals, and rodents carrying the gene for osteopetrosis have been invaluable in elucidating the origin and function of the osteoclast, the cell that is deficient in this disease. Indeed, study of experimental osteopetrosis has been responsible for important therapeutic advances in man.⁵

The condition occurs in two main forms, although intermediate varieties have been described. The benign autosomal dominant type, referred to as Albers-Schönberg's or marble bone disease, is usually a chance finding on radiological examination. Life expectancy is normal, and no treatment is needed. The autosomal recessive type is a lethal disease recognisable at or soon after birth, and until recently no affected child had survived the first decade. Not only are the medullary cavities of shaft bones obliterated by unresorbed juvenile bone but they are also misshapen, stunting growth. Encroachment on neural foraminae by bone causes blindness,

deafness, and facial paralysis. Anaemia is severe and compensated for by hepatosplenomegaly. Death occurs from anaemia, haemorrhage, or infection. The intermediate forms of osteopetrosis are associated with rickets, renal tubular acidosis, and deficiency of the enzyme carbonic anhydrase II. The latter is important because it offers a means for genetic counselling. Accurate diagnosis is important not only because specific treatment is now available but also because the battered child syndrome has been mistaken for a form of bone dysplasia.⁶

As osteopetrosis is caused by a failure of osteoclastopoiesis a hormonal defect was suspected. Yet giving parathyroid hormone, vitamin D metabolites, and adrenal corticosteroids and manipulating calcium metabolism were all without much effect. Haematological features were temporarily improved by blood transfusion, splenectomy, and treatment with iron. Then Reeves *et al* in 1979 correctly postulated a generalised inherited abnormality of phagocytes and even osteoclasts, which was unlikely to be influenced by such treatments.⁷

Osteopetrosis also occurs widely in animals, and Walker was the first to use transplantation—parabiosis and bone marrow—in osteopetrotic rodents to show failed bony resorption because of a stem cell and not a hormonal defect.⁸ Skeletal sclerosis in the treated mutants disappeared, showing that an osteoclastic stem cell or precursor in the haematopoietic tissue was conveyed by the blood stream to sites needing bony resorption.⁸ Once the stem cell deficiency was confirmed⁹ it was a small step to clinical application. Further research with osteopetrotic rodents carrying cellular markers showed that in syngeneic bone marrow grafts (comparable to grafts from one identical twin to another) donation of a few stem cells was enough to maintain bony resorption. Immunosuppression was needed if there were greater antigenic differences between donors and recipients.¹⁰

The first report of human osteopetrosis treated with cellular infusions without immunosuppression produced results that might have been predicted from the experimental findings.¹¹ Of four children, one benefited by a marrow graft from her HLA identical sister, but the others did not because the graft failed to take. Thymic cells were also given in the belief that the thymus was important in bone resorption, but this has since been disproved.^{12,13} Three reports have described successful treatment of human autosomal recessive osteopetrosis with bone marrow transplants from matched sibling donors and with varying immunosuppressive regimens.¹⁴⁻¹⁶ A review of 14 children shows that six have been successfully treated, a considerable achievement in a disease that used to be always fatal.¹⁷

Study of osteopetrosis in animals and man has helped our understanding of osteogenesis¹⁸ and led to the disease being included among those that may be helped by bone marrow transplantation.¹⁹ The search for the primordial stem cell for the osteoclast continues.²⁰

N W NISBET

Retired Worker, supported by the
Medical Research Council,
Charles Salt Research Centre,
Robert Jones and Agnes Hunt Orthopaedic Hospital,
Oswestry,
Shropshire SY10 7AG

I am obliged to Dr John Loutit FRS and Dr Michael Marshall for their collaboration and help with the manuscript and to Miss Claire McLoughlin for secretarial help.

- 1 Albers-Schönberg H. Röntgenbilder einer seltenen Knochenkrankung. Münchener Medizinische Wochenschrift 1904;51:365-9.
- 2 Beighton P, Horan F, Hamersma A. A review of the osteopetroses. Postgrad Med J 1977;53:507-15.