solely or more intensely in the five days preceding the onset of menstruation. In other words, "contamination" by psychological symptoms of the premenstrual syndrome was specifically avoided.

Finally, Ms Warner doubts whether the findings about consultation rates have clear implications for general practice. The basis for her doubt is that we studied "a small total number of consultations." Nevertheless, in a random sample of more than 500 women the proportions of women consulting their general practitioner were as high as 24% for gynaecological and 17% for psychiatric complaints over 12 months. Moreover, the detailed findings do seem to have implications for general practice. Thus, of 127 women who consulted their doctor for gynaecological complaints, 70 had menstrual symptoms, of whom 12 (17.1%) were present state examination cases. Of the remaining 57 women, only two (3.5%) were present state examination cases. Among the 91 patients who had consulted their doctor for psychiatric problems 53 (58%) had menstrual symptoms at interview, of whom 19 (34%) were present state examination cases; by contrast, only four patients had non-menstrual gynaecological complaints, and none of them were present state examination cases. For these reasons we believe that the data about consultation histories, together with the overall pattern of association between gynaecological symptoms and mental state, support our recommendation that, "if a woman presents in the surgery with a gynaecological complaint, particularly with a menstrual complaint, it is advisable to look for any associated psychiatric disorder."

DENNIS GATH
SUSAN ILES
ANN DAY

Department of Psychiatry,
University of Oxford,
Oxford OX3 7JX

Are we drinking our neurones away?

SIR,—The quantitative histology of cerebral atrophy in chronic alcoholism is a controversial topic, so it is unfortunate that the article by Professor Clive Harper and colleagues (28 February, p 534) lacks certain critical details, casting doubt on the accuracy of some of the findings.

Our curiosity was aroused by the inconsistency of the cell counts in the alcoholic brain. It is particularly remarkable that the superior frontal gyrus showed a considerable loss of neurones without change in the glial cell population of the underlying white matter. Axonal degeneration in the white matter generally accompanies cortical neurone degeneration, and so a deficit of glia would be expected. A reduction in glial cells might also be predicted from the reduced brain density of chronic alcoholic patients shown by computed tomography.¹

We note that cell counts per unit volume were used and that Professor Harper and coworkers do not mention any correction for tissue atrophy. Cell counts in atrophic brain are comparable with control values only when allowance is made for alterations in tissue volume. If the cerebral cortex shrinks during cell loss the surviving cells pack down and their density (count per unit volume) may remain unchanged; indeed, the cellular density may actually increase if there is selective loss of the largest cell type. This pattern of change is found, for example, in the cerebral cortex of patients with Alzheimer's disease, which shows a selective loss of large neurones with an increase in the density but not the absolute numbers of small neurones and glial cells.^{2,3}

If a correction for tissue atrophy has been omitted, a rough estimate of the cell content in

Mean cell content of brains of alcoholic patients in relation to that of brains of controls

	Alcoholic patients		
	Controls	Cells/mm ³	Estimated absolute cell content
Frontal region:			
Neurones	100	77.8	69.6
Cortical glia	100	112.7	106.2
White matter glia	100	100.6	90.0
Motor region:			
Neurones	100	107.4	101.1
Cortical glia	100	106.9	100.7
White matter glia	100	111.4	99.7

absolute terms may be obtained by multiplying the values for tissue volume with those for cellular density. In the table we have expressed the cell counts per unit volume and also the estimated absolute counts from the alcoholic brains as quotients of the control values. This calculation suggests that the neurone deficit from the frontal cortex was underestimated and that there was indeed a loss of glial cells from the frontal white matter. On the other hand, the cell content of the motor region seems to be almost exactly the same in alcoholic patients as in controls.

A second criticism of the presentation is that we have to assume that a reduction in mean neurone area represents cellular shrinkage. In the frontal cortex, at least, however, the reduction could be due to selective loss of the largest cells as neurones of all sizes were measured together. Convincing evidence of neurone shrinkage requires measurement of the number and mean area of neurones sorted into three or more size categories.

J M ANDERSON
BETHAN M HUBBARD

Department of Pathology,
Ninewells Hospital and Medical School,
Dundee DD1 9SY

- 1 Golden CJ, Graber B, Blose I, *et al.* Difference in brain densities between chronic alcoholic and normal control patients. *Science* 1981;211:508-10.
- 2 Mountjoy CQ, Roth M, Evans NJR, Evans MH. Cortical neuronal counts in normal elderly controls and demented patients. *Neurobiol Aging* 1983;4:1-11.
- 3 Hubbard BM, Anderson JM. Age related variations in the neuron content of the cerebral cortex in senile dementia of Alzheimer type. *Neuropathol Appl Neurobiol* 1985;11:369-82.

AUTHORS' REPLY—With regard to the first point made by Drs Anderson and Hubbard—namely, that the glial cell population in the frontal white matter did not seem to differ in the control and alcoholic groups—there are several explanations and confounding facts to be considered. Firstly, as they point out, the actual shrinkage of the white matter, which is an important component of brain shrinkage in alcoholic patients, may cause a compaction of the glial cells and hence mask an actual reduction in the glial cell density. In a significant proportion of alcoholic patients (particularly the younger ones) the white matter shrinkage seems to revert towards normal after long periods of abstinence. Though the pathophysiological mechanisms underlying this reversible change have not yet been elucidated, it is probably not associated with changes in numbers of glial cells but related to some structural change in a component such as myelin. We are unaware of other studies in which white matter glial cell counts have been performed in patients with a known loss of neurones, such as those with Alzheimer's disease. Histologically, there is no obvious change in the cellularity of the white matter in such cases.

Despite the fact that computed tomography

has suggested reduced brain density in chronic alcoholic patients our studies of the specific gravity of grey and white matter in controls and alcoholic patients have shown a consistent, though not statistically significant, reduction in specific gravity in the alcoholic patients.

Neuronal counts per unit volume were not corrected for tissue atrophy because our previous studies have shown that cortical (grey matter) shrinkage is not significant in the alcoholic group and because it is difficult to calculate the shrinkage factor owing to substantial variation in brain volume. In any case, as Drs Anderson and Hubbard point out, such a correction would increase the difference in cell counts between the control and alcoholic groups.

With regard to neuronal shrinkage, further studies are in hand to examine the selective vulnerability of specific neuronal populations. It is conceivable that, as in Alzheimer's disease, large neurones might be more susceptible than small neurones. As there was shrinkage but no loss of cortical neurones in the motor cortex, however, it seems more likely that there is also a non-selective shrinkage of neurones in the superior frontal cortex.

CLIVE HARPER
JILLIAN KRIL

Department of Pathology,
University of Sydney,
Sydney 2006

JOHN DALY

Department of Neuropathology,
Royal Perth Hospital,
Perth

New contract for pharmacists

SIR,—The comments by Scrutator (14 March, p 715) may give rise to misconceptions about the new contract for National Health Service pharmaceutical services and its interaction with the Clothier scheme for dispensing in rural areas. The Clothier scheme aims not at preserving the status quo in rural areas but at ensuring that any changes are brought about in a controlled way. One of the aims of the new contract is to achieve a more rational distribution of pharmaceutical services in both urban and rural areas. Though these two aims are different, they are not necessarily opposed and could even be considered to be complementary.

The NHS regulations and Department of Health and Social Security guidance on the new contract make it clear that an application for the opening of a new pharmacy in a rural area may be considered by the pharmacy practices subcommittee of the family practitioner committee only if it already has the approval of the rural dispensing committee.^{1,2} The criterion used by the rural dispensing committee is whether the opening of a pharmacy would be prejudicial to the proper provision of general medical or pharmaceutical services in the area, while the pharmacy practices subcommittee will be required to decide whether or not the proposed pharmacy is necessary or desirable for the adequate provision of NHS pharmaceutical services. In other words, a pharmacist wishing to open a pharmacy in a rural area will have to prove, firstly, to the rural dispensing committee that his proposal will not damage existing medical or pharmaceutical services and, secondly, to the pharmacy practices subcommittee that his proposal will improve the pharmaceutical services.

No doubt there will be pharmacists who will resent this double barrier to their freedom to practise their profession in a location of their choice and doctors who will see it as a threat to the additional income that they obtain from dis-

pending. It is to be hoped, however, that most members of both professions will welcome the new contract as a means of achieving a more rational distribution of pharmacies and ensuring a more comprehensive pharmaceutical service for all our patients, both rural and urban.

IAN G SIMPSON

Oxfordshire Local Pharmaceutical Committee,
Oxford OX3 0JU

1 National Health Service, England and Wales. *National Health Service (General Medical and Pharmaceutical Services) Amendment (No 2) Regulations 1987*. London: HMSO, 1987. (Statutory Instrument No 401.)

2 Department of Health and Social Security. *Health notice HC(FP)(87)2*. London: HMSO, 1987.

LMSSA: a back door entry into medicine?

SIR,—I was dismayed to read Mr Richard Wakeford's article (4 April, p 890), with its many inaccuracies and innuendos. The Society of Apothecaries examines candidates by authority of the Medical Act 1886 passed by parliament and not by virtue of a charter granted by King James I. This authority is the same as that for every other examining body.

Most of our examiners are consultants from teaching hospitals, who are also examiners for the MB examination. It is generally agreed that the standard of the examination is the same as that of the MB, and a close marking system is adopted so that although a marginal failure in any one section of the examination may be compensated for by extra marks in another section, a severe fail cannot be compensated for. The examiners are all external as far as the candidates are concerned, which ensures an independent assessment, unbiased by any previous knowledge of the candidate's performance during his or her student years. This system is adopted in all postgraduate examinations and is considered to be fairer to the candidate.

Mr Wakeford referred to 10 candidates who failed the Cambridge MB in December 1982. Altogether 13 Cambridge students entered for the LMSSA in 1983, all of whom had been approved by the dean of the medical school. Six completed the LMSSA at the first attempt, four completed it after more than one attempt, and three who failed the LMSSA subsequently passed the Cambridge MB. It is unreasonable to conclude that the society's examination is a "second rate qualification" on the basis of these figures.

The regulations ensure that candidates must pass all sections of the examination within two years and do not allow more than four attempts at any one section. In practice, however, it is unusual for candidates to complete the examination if more than two attempts are required to pass any one part. Overseas candidates are accepted only after graduating from medical schools approved by the General Medical Council and are required to have passed the Professional and Linguistic Association Board examination or been exempted from it.

Mr Wakeford refers to some universities that allow students only two attempts at the final examination. It is accepted that the selection of medical students ensures that only those of the highest intellectual ability may begin the course, and it is surely a great waste of public money if such students are denied the chance of qualifying after a long period of study. Above all, regular inspection of all bodies by the General Medical Council ensures the maintenance of high standards, and the Society of Apothecaries' examinations committee has always been ready to adapt to the council's suggestions.

The Society of Apothecaries has provided a diploma of a high standard since 1815, and we intend to maintain the service that we have

hitherto provided for the profession and the public. Criticism of our examination is particularly resented when it comes from a medically unqualified and uninformed source who has never attended one of the society's examinations.

W F W SOUTHWOOD

Worshipful Society of Apothecaries of London,
London EC4V 6EJ

Points

Perforation of nasal septum due to button battery lodging in nose

Mr M DAVENPORT (Children's Hospital, Sheffield S10 2TH) writes: Dr P Fernando's report of an alkaline button battery lodged in the nasal cavity (21 March, p 742) may cause misunderstanding about the management of the more common problem of batteries being swallowed. If the battery has passed beyond the oesophagus and is progressing through the gastrointestinal tract it is safest to await its passage.¹ The mechanism of perforation in the case described by Dr Fernando seems somewhat speculative. A battery's alkaline contents would leak only if its metal casing had disintegrated—surely an improbable event after only 24 hours in the nasal cavity. It has been suggested that a rise in pH occurs at the anode side because of the current produced by an undischarged cell and that it is this that causes local tissue necrosis, rather than the battery's contents.

1 David JJ, Ferguson AP. Management of children who have swallowed button batteries. *Arch Dis Child* 1986;61:321-2.

Dr D W SKINNER (Queen's Medical Centre, Nottingham NG7 2UH) writes: I should like to emphasise the concern about tissue necrosis and other long term consequences of button batteries as foreign bodies not only in the nose, as described by Dr P Fernando (21 March, p 742), but also in the ear. Any delay in the removal of button batteries may have very severe consequences. Recently, I reported on a 12 year old child with a button battery in the nose, who suffered extensive destruction of the inferior turbinate and adjacent septum, followed by a persistent local atrophic rhinitis, and will probably develop intranasal adhesions as a consequence. A case of a 5 year old child with a button battery in the external auditory canal has also been described, in which complete destruction of the tympanic membrane and ossicular chain occurred, with subsequent pronounced conductive hearing loss and great difficulties in reconstructing the middle ear structures.¹

1 Skinner DW, Chui P. The hazards of "button-sized" batteries as foreign bodies in the nose and ear. *J Laryngol Otol* 1986;100:1315-8.

Identity cards for patients infected with HIV?

Dr GIUSEPPE ENRICO BIGNARDI (Pathology Department, Stepping Hill Hospital, Stockport SK2 7JE) writes: Dr N West (21 March, p 772) pointed out the consequences of breaches of confidentiality about patients with the acquired immune deficiency syndrome (AIDS). The dispatch of pathological specimens is unfortunately the best way to spread information on the identity of AIDS patients. According to DHSS guidelines, all specimens sent to a laboratory for whatever test must be accompanied by a form indicating knowledge or suspicion of HIV infection and a senior laboratory staff member must be informed in advance. This is unjustified. The risk of AIDS to health service staff is minimal and certainly lower than that posed by specimens positive for hepatitis B surface antigen.¹² While specimens are rightly tagged with a label indicating risk of infection, there is no reason for the card to carry other than a general label: possibly only one kind of label—"Risk of viral infection"—should be used for both hepatitis and HIV specimens with no need for any further indication. When samples are sent for HIV antibody

testing the identity of the patient could be protected by assigning the sample a number rather than a name, as in genitourinary medicine clinics.

- 1 McCray E. Occupational risk of the acquired immunodeficiency syndrome among health care workers. *N Engl J Med* 1986;314:1127-32.
- 2 Geberding JL, Hopewell PC, Kaminsky LS, Sande MA. Transmission of hepatitis B without transmission of AIDS by accidental needlestick. *N Engl J Med* 1985;312:56.

Future of the pathologist in an era of technological change and cost containment

Dr J K JOHNSON (Dumfries and Galloway Royal Infirmary, Dumfries DG1 4AP) writes: I am concerned that greater emphasis on patient care in chemical pathology, as suggested by Drs M H Labib and L R Ranganath (7 March, p 642), may lead to neglect of the pathological role of the specialty. Advertisements for service registrar posts in chemical pathology are now attracting very high calibre applicants. Though it is not a requirement, the competition ensures that most of those who are appointed have either a scientific qualification as well as their medical qualification or a higher medical qualification with proved aptitude for laboratory work. These are the doctors who are most likely to become chemical pathologists. The potential, in laboratory medicine, for an applicant with the MRCP but no scientific background is questionable.

Overuse of monitoring of blood concentrations of antiepileptic drugs

Dr E M R CRITCHLEY (Department of Neurology, Royal Preston Hospital, Preston, Lancashire PR2 4HT) writes: As Dr D W Chadwick suggests (21 March, p 723), concentration on drug monitoring may delay the search for the most suitable anti-convulsant for a particular patient. If an anti-convulsant does not reduce the number of fits when it is introduced it is unlikely to be successful if used in a higher dosage. If a patient's epilepsy becomes uncontrollable the possibility of an underlying lesion or of new trigger factors should be the doctor's first consideration. One man who was admitted to hospital repeatedly had his drug dosage increased on each admission. He was, in fact, an active homosexual who would have a fit whenever he got into trouble. Another patient, who weighed less than 44.5 kg and had miliary tuberculosis and meningitis, had a focal fit. He was given phenytoin but the serum concentration was monitored and the dosage increased to 400 mg with no account being taken of hepatic induction, which affected the blood concentrations of the antituberculous drugs that he was receiving.

Amenorrhoea

Dr G LIVINGSTON (Academic Department of Psychiatry, Royal Free Hospital, London NW3) writes: Dr S Franks mentions that the most common cause of secondary amenorrhoea, other than pregnancy, is weight loss (28 March, p 815) and that "the distinction between these patients and those with anorexia nervosa is not clear cut." As anorexia nervosa may be defined as "the mental symptoms of fear of fatness and distortion of body size, the behavioural signs of self starvation and weight loss and the pathophysiological state marked by amenorrhoea,"¹ many of these women may have anorexia nervosa. Anorexia nervosa is a positive diagnosis made clinically, which would be missed on this algorithm. It does not require endocrine investigations. In patients with amenorrhoea and weight loss the first investigation should be a history and identification of abnormal attitudes to eating. This would allow diagnosis and early treatment of anorexia nervosa, thus decreasing mortality and morbidity and saving the expense of unnecessary investigations.

- 1 Folstein MF, Wakeling A, DeSouza V. Analogue scale measurement of the symptoms of patients suffering from anorexia nervosa. In: Vigersky RA, ed. *Anorexia nervosa*. New York: Raven Press, 1977.