birth. No meconium was found in the airways. A chest x ray picture at 40 minutes showed an infiltrate in the right lower lobe. After extubation her condition remained stable, and she was followed in the neonatal unit for four days. On discharge at 5 days of age the girl was healthy and the chest x ray picture was normal.

The bronchial aspirate was analysed by thin layer chromatography, which showed an unidentified substance that was also present in the vaginal tablet. No prostaglandin E₂ was shown, however, with this technique. The concentration of prostaglandin E₂ in the aspirate, as determined by specific radioimmunoassay (antiserum from Pasteur Institute, Paris),³ was 1.5 ng/ml, which fell within the range of 0.3-4.7 ng/ml in normal neonates (n=4).

Comment

We believe that this is the first reported case of neonatal aspiration syndrome caused by undissolved remnants of a vaginal tablet containing prostaglandin. Results of thin layer chromatography suggested that the granular material suctioned from the airways contained the vehicle substance of the tablet. The clinical signs were the result of mechanical obstruction of the airways and were not caused by prostaglandin E₂. This conclusion is supported by the normal concentration of prostaglandin E₂ in the aspirate. Thus the tablet had probably released its prostaglandin into the vagina, and this apparently promoted the labour. Remnants of the vehicle substance were then transferred to the mouth of the baby while the head was passing through the vagina. The subsequent aspiration, which occurred during the baby's first breaths, was promptly treated by intubation and suction of the airways.

Aspiration syndrome is one of the most common emergencies in the delivery room. This potentially life threatening condition may also be iatrogenic, as shown in this case report. Obstetricians and paediatricians should be aware of the risk of using prostaglandin tablets, a risk that may be avoided by replacing prostaglandin tablets with prostaglandin viscous gel, which disappears from the vagina more quickly than tablets.⁴

- 1 MacKenzie IZ, Embrey MP. Cervical ripening with intravaginal PGE₂ gel. *Br Med J* 1977;ii:1369.
- 2 Wilson PD. A comparison of four methods of ripening the unfavourable cervix. *Br J Obstet Gynaecol* 1978;85:941.
- 3 Embrey MP, Graham NB, McNeill ME. Induction of labour with a sustained-release prostaglandin E₂ vaginal pessary. *Br Med J* 1980;281:901-2.
- 4 Ulmsten U, Wingerup L, Ekman G. Local application of prostaglandin E₂ for cervical ripening or induction of term labor. *Clin Obstet Gynecol* 1983;26:95-105.
- 5 Seppälä E, Pora O, Metsä-Ketelä T. A modified method for extraction and purification of prostaglandins with resin XAD-2. *Prostaglandins, Leukotrienes Med* 1984;14:235-41.

(Accepted 26 March 1987)

Children's Hospital, Helsinki, Finland

STURE ANDERSSON, MD, registrar in paediatrics
RITVA PAETAU, MD, registrar in paediatrics
MIKKO HALLMAN, MD, chief of neonatal unit

Department of Biomedical Sciences, University of Tampere, Tampere, Finland

HEIKKI VAPAATALO, MD, professor of pharmacology

I and II Department of Gynaecology and Obstetrics, University Central Hospital, Helsinki, Finland

OLAVI YLIKORKALA, MD, associate professor

Correspondence to: Dr Sture Andersson, Children's Hospital, Stenbäckinkatu 11, 00290 Helsinki, Finland.

Treatment of sensorineural deafness associated with ulcerative colitis

There is evidence that the pathogenesis of inflammatory bowel disease and its extraintestinal manifestations depend on immunological mechanisms. Recently, sensorineural deafness was added to the list of extraintestinal complications of ulcerative colitis, and as the disease is also associated with rheumatoid arthritis and polyarteritis nodosa it may have an immune basis.

In two of three cases the hearing loss associated with ulcerative colitis improved with steroids either alone¹ or in combination with cyclophosphamide.² The precise relation between improved hearing and drug treatment, however, was not studied in detail. As the hearing loss may result in serious morbidity prompt and appropriate treatment of this complication is essential. We describe the relation between hearing loss in ulcerative colitis and treatment with prednisolone and sulphasalazine either alone or in combination with azathioprine.

Case report

A 42 year old man who presented with a 17 year history of "colitis" and with symptoms of diarrhoea and rectal bleeding was found to have total ulcerative colitis of moderate severity on colonoscopy, which was confirmed by biopsies. His symptoms and macroscopic disease responded promptly to a short course of prednisolone and sulphasalazine. Four months after presentation he developed vertigo, nystagmus, and ataxia of sudden onset followed by complete deafness in the right ear and progressive deafness in the left. Neurological investigations, including a lumbar puncture and computed tomography of the brain, yielded normal results, but audiometry showed sensorineural deafness with complete loss of hearing in the right ear and 15 decibel loss in the left ear for most of the speech frequencies. His hearing remained stable at this level, and he was discharged taking sulphasalazine, the ulcerative colitis being quiescent at that time.

Three months later his hearing suddenly deteriorated. His ulcerative colitis remained in remission, autoantibody levels were normal, and immune complex titres were undetectable. On the assumption that his hearing loss was an extraintestinal manifestation of ulcerative colitis, prednisolone 60 mg/day was started. His hearing rapidly improved with this regimen but deteriorated whenever the dose was reduced below 40 mg/day (figure). The addition of azathioprine (2 mg/kg/day) allowed the steroid treatment to be slowly reduced and then stopped without appreciable deterioration in hearing (figure). Sulphasalazine was continued throughout in a dose of 1 g thrice daily. The hearing loss remained stable for over nine months despite an exacerbation of ulcerative colitis.

Prednisolone dosage and loss of hearing before and during treatment with azathioprine.

Comment

This report provides a further example of sensorineural hearing loss associated with ulcerative colitis. It shows dissociation between the activity of the colitis and impairment of hearing and the responsiveness of the disorder to prednisolone and azathioprine. In a young patient sudden loss of hearing results in severe disability and salvaging of residual auditory function is vital. On the basis of this experience we would recommend immediate treatment of such sensorineural deafness with a combination of azathioprine and prednisolone.

- 1 Weber RS, Herman AJ, Newton JC. Sensorineural hearing loss associated with ulcerative colitis. *Arch Otolaryngol* 1984;110:810-2.
- 2 Summers RW, Harker L. Ulcerative colitis and sensorineural hearing loss: is there a relationship? *J Clin Gastroenterol* 1982;4:251-2.

(Accepted 2 April 1987)

Department of Gastroenterology, Hope Hospital, University of Manchester School of Medicine, Salford

A DOWD, BM, MRCP, medical registrar
W D W REES, MD, MRCP, consultant physician and gastroenterologist

Correspondence to: Dr Dowd.