in such a group is not straightforward, but the cumulative incidence of Parkinson's disease, disease of the spinal cord, cerebellar syndromes, and dementia seemed much higher than could be expected.

These ideas may have important implications for those investigating the causes of neurodegenerative disorders. Firstly, a better understanding of regional vulnerability in the central nervous system to particular types of toxins might allow prediction of the sort of environmental agents that could be responsible for particular diseases and so provide clues for the epidemiologist. Secondly, attention should be directed at environmental exposures that occurred many years before the first symptoms. Thirdly, investigation of factors that influence the rate of cell loss with increasing age, in particular the possibility that an already depleted neuronal population loses cells faster than normal, could lead to treatments that would prevent the progression of subclinical damage to overt disease.

CHRISTOPHER N MARTYN

Clinical Scientist,
MRC Environmental Epidemiology Unit,
Southampton General Hospital,
Southampton SO9 4XY

- 1 Spencer PS, Nunn PB, Hugon J, Ludolph AC, Ross SM, Roy DW, Robertson RC. Linkage of Guam amyotrophic lateral sclerosis/parkinsonism-dementia to a plant excitant neurotoxin? *Science* 1987;237:517-20.
- 2 Spencer PS. Guam ALS/parkinsonism-dementia: a long-latency neurotoxic disorder caused by "slow toxin(s)" in food? *Can J Neurol Sci* 1987;14:347-57.
- 3 Denny-Brown D. Neurological conditions resulting from prolonged and severe dietary restriction. *Medicine (Baltimore)* 1947;26:41-113.
- 4 Spencer PS, Roy DW, Ludolph A, Hugon J, Duived MP, Schaumburg HH. Lathyrism: evidence for the role of the neuroexcitatory amino acid BOAA. *Lancet* 1986;ii:1066-7.
- 5 Meldrum B. Possible therapeutic applications of antagonists of excitatory amino acid neurotransmitters. *Clinical Science* 1985;68:113-22.
- 6 Langston JW. MPTP and Parkinson's disease. *Trends in Neurosciences* 1985;8:79-83.
- 7 Ministry of Health, Mozambique. Mantakassa: an epidemic of spastic paraparesis associated with chronic cyanide intoxication in a cassava staple area of Mozambique. 1. Epidemiology and clinical and laboratory findings in patients. *Bull WHO* 1984;62:477-84.
- 8 Ministry of Health, Mozambique. An epidemic of spastic paraparesis associated with chronic cyanide intoxication in a cassava staple area of Mozambique. 2. Nutritional factors and hydrocyanic content of cassava products. *Bull WHO* 1984;62:485-92.
- 9 Calne DB, Langston JW. Aetiology of Parkinson's disease. *Lancet* 1983;ii:1457-9.
- 10 Calne DB, Langston JW, Martin WR, et al. Positron emission tomography after MPTP: observations relating to the cause of Parkinson's disease. *Nature* 1985;317:246-8.
- 11 Calne DB, Eisen A, McGeer E, Spencer P. Alzheimer's disease, Parkinson's disease, and motoneurone disease: abiotrophic interaction between ageing and environment? *Lancet* 1986;ii:1067-70.
- 12 Gibberd FB, Simmonds JP. Neurological disease in ex-far-east prisoners of war. *Lancet* 1980;ii:135-7.

Unconventional viruses and the central nervous system

For several diseases a causative agent has still to be identified. Among these agents are those causing progressive degeneration of the central nervous system such as Creutzfeld-Jakob disease and kuru in man and scrapie in sheep and goats. These are referred to as unconventional viruses but there is no certainty about their composition or physical organisation. Indeed, are they viruses? This question was the focus for much of the discussion at a recent symposium held at the Ciba Foundation.*

Slow virus disease has generated considerable interest not only because of the enigmatic nature of the agents but also because of the similar lesions seen in Alzheimer's disease and the increasing suspicion that this disease too may be caused by a similar type of agent (though Alzheimer's disease has not

yet been transmitted to animals). Besides being linked by their similar pathological lesions Creutzfeld-Jakob disease, kuru, and scrapie all have a prolonged incubation period, sometimes lasting several years. The first two can be transmitted to apes and monkeys and scrapie can be transmitted to hamsters and mice but even in animals the incubation period is prolonged, making assay extremely slow and inaccurate. This in turn has made identification of the agents much more difficult than usual.

Nevertheless, the presence of about one billion infectious units per gram of brain from a mouse infected with scrapie would appear to someone outside the subject to provide ample material for purification studies. Why, then, has progress been so slow? Apart from the time required to assay the agent the other main reason is the starting material itself. Brain tissue is particularly difficult material to work with and the research worker can never be sure whether the effect of a particular reagent is being influenced by the host material. False suppositions in the early days led to many false starts, with the result that in the past 30 years there have been numerous hypotheses on the nature of the scrapie agent. Recently, however, attention has been focused on three of them.

The first hypothesis is that the scrapie agent is a virus after all, being identical with the scrapie associated fibrils seen on electron microscopy in preparations from infected animals. The virus hypothesis is supported by studies on the kinetics of inactivation of infectivity with different chemical and physical agents. The proponents of this view conclude that the agent contains a nucleic acid with a molecular weight of about one million. Although this is small it is not exceptionally so. Moreover, there are well defined strains of scrapie causing different signs and having different incubation times, which is a clear indication that the agent has its own independently replicating genome. Nevertheless, no scrapie specific nucleic acid has been identified.

The second hypothesis postulates the existence of a virino—a low molecular weight nucleic acid together with a protein derived from the host. In this model the nucleic acid is scrapie specific but does not encode any protein, and the nucleic acid of the agent would induce disease by disrupting the regulatory functions of the nucleic acids of the host. Moreover, given that the protein is derived from the host, the absence of an immunological response in infected animals would be explained.

The third and most provocative hypothesis is that the agent is a self replicating protein. The name prion (proteinaceous infectious particle) has been coined by the proponents of this view, who based it to the observation that, although the infectivity of brain homogenates was resistant to various treatments which should destroy nucleic acid, it was destroyed by several protein specific reagents. Moreover, workers have failed to detect any scrapie specific nucleic acid in highly purified infectious preparations of the agent. The major component of these preparations is a protein with a molecular weight of about 30 000, but the gene coding for this protein also turns out to be present in normal brain as well as other normal tissues. Thus apparently the prion protein is encoded in the host genome before infection. The most decisive information about the role of the prion would come from expression of the gene coding for it. If the protein alone is infectious then the product obtained by expressing the gene should also be infectious. So far this has not been shown.

One suggestion is that the prion protein is a component of

*The proceedings of the symposium will be published by John Wiley and Sons early in 1988 under the title *Novel Infectious Agents in the CNS*.

the amyloid plaques and fibrillar structures found in scrapie infected brain and it is clearly necessary to determine the relationship between them. An electron microscopic study of the fibrils found in infected brain extracts, reported at the Ciba meeting, has shown particles similar to those of tobacco mosaic virus from which part of the encapsulating protein has been removed. Are the exposed tails in fact the scrapie specific nucleic acid? It should be possible to answer this question by precise biochemical analysis.

Despite the differences of opinion and interpretation of data the clear message from the Ciba meeting was that with the powerful analytical techniques available the problem of the nature of the agent could be solved within the next few years. We should then have the springboard for attempts to control the diseases caused by this group of agents.

FRED BROWN

Head of Virology Research,
Wellcome Biotechnology Ltd,
Woking,
Surrey GU24 0NQ

Acute stress erosions: can they be prevented?

Stress erosions are superficial mucosal lesions found mainly in the fundus of the stomach. They occur in most if not all critically ill patients.¹ Acute gastric ulcers may give rise to bleeding if they erode arterioles in the gastric submucosa, or to perforation (as of the fundus and duodenum in patients with severe burns—the so called Curling's ulcer²). Both shock and hypotension probably have important roles in the formation of stress lesions,³ and many studies have been undertaken to assess the value of antacids and histamine H₂ receptor antagonists in preventing them. Do such agents work and is one type better than the other?

Such studies are difficult to analyse because of the heterogeneous nature of the lesions and because the researchers have not always included an untreated control group. Many studies have relied on the presence of occult blood in nasogastric aspirates as evidence of bleeding rather than on endoscopic inspection, although such blood may simply be caused by the trauma. Nevertheless, the balance of evidence has suggested that treatment with H₂ antagonists is less efficacious than the frequent administration of antacids.^{4,5} An exception seems to occur with the bleeding from erosions in fulminant hepatic failure, which is reduced by cimetidine, but with no effect on the overall mortality.⁶

Poleski and Spanier have compared the efficacy of prophylactic antacid with cimetidine in preventing stress erosions in critically ill patients, using gastroscopy to evaluate effectiveness.⁷ Some 44 patients at risk were randomised to receive cimetidine (300 mg six hourly intravenously initially to a maximum of 400 mg four hourly) or antacid (Mylanta II 30-90 ml/hour), the minimum dose being used to maintain hourly gastric pH around 4. Both treatments were equally effective in preventing stress ulcers despite the control of pH being inferior in the cimetidine group. The authors suggested that measurements of gastric pH might not fully reflect the beneficial effects of cimetidine, since this is only a concentration measurement and does not reflect volume of acid secreted. Cimetidine might protect the mucosa by decreasing

the acid load and also by an intrinsic effect.⁸ They confirmed that appreciable bleeding was rare, for none of their patients developed this despite 13 patients having several fundal erosions on endoscopy.

Possibly also the frequency of stress erosions and bleeding is decreasing because of improved supportive care,^{9,10} so might it not be wise to identify these critically ill patients who are at the maximum risk and give them prophylaxis? Given the side effects of the H₂ receptor antagonists,¹¹⁻¹⁴ it seems reasonable to restrict the latter to patients unable to retain antacids. Even the alkalinisation of gastric secretions in the critically ill poses the additional risk of reducing the indigenous flora in the stomach with the overgrowth of potentially pathogenic aerobic bacteria, which may infect vulnerable organs such as the lungs.¹⁵

Thus the doctor may reasonably restrict the use of antacids and H₂ receptor antagonists to those patients in the intensive therapy unit who are at serious risk of haemorrhage secondary to stress erosions. Such patients include those with severe uncontrolled sepsis, acid base disturbances, major organ failure, and extensive trauma including burns and head injuries. Patients having prolonged ventilation for respiratory failure appear to be particularly at risk.¹⁶ Antacids are preferable to H₂ antagonists, which should be used when antacids cannot be given. Whether ranitidine will prove to be safer than cimetidine is uncertain, and only time will tell.¹⁷ Methyl substituted prostaglandins will shortly be introduced on to the market and one of their selling points will undoubtedly be cytoprotection. Prostaglandins are concerned in the production of mucus, local secretion of bicarbonate, and gastric mucosal blood flow.¹⁸ How important these factors will be in preventing drug and stress induced gastrointestinal haemorrhage remains to be determined by studies now in progress.

GILLIAN C HANSON

Consultant Physician,
Whipps Cross Hospital,
London E11 1NR

B G GAZZARD

Consultant Physician,
Westminster Hospital,
London SW1P 2AP

- 1 Czaja AJ, McAlhany JC, Pruitt BA Jr. Acute gastroduodenal disease after thermal injury: an endoscopic evaluation of incidence of natural history. *N Engl J Med* 1974;291:925-9.
- 2 Pruitt BA, Goodwin CW. Stress ulcer disease in the burned patient. *World J Surg* 1981;5:209-22.
- 3 Moody FG, Zalesky CA, Larsen KG. Cytoprotection of the gastric epithelium. *World J Surg* 1981;5:153-63.
- 4 Stothert JC, Simonowitz DA, Dellinger EP, et al. Randomised prospective evaluation of cimetidine and antacid control of gastric pH in the critically ill. *Ann Surg* 1980;192:169-74.
- 5 Weigelt IA, Awbakken M, Geruetz BL, Snyder WH. Cimetidine versus antacid in prophylaxis for stress ulceration. *Arch Surg* 1981;116:597-602.
- 6 MacDougall BRD, Bailey AJ, Williams R. H₂ antagonists and antacids in the prevention of acute gastrointestinal haemorrhage in fulminant hepatic failure. *Lancet* 1977;i:617-8.
- 7 Poleski MH, Spanier AH. Cimetidine versus antacids in the prevention of stress erosions in critically ill patients. *Am J Gastroenterol* 1986;81:107-11.
- 8 Guth PH, Awes D, Paulsen G. Topical aspirin plus HCL gastric lesions in the rat: cytoprotective effect of prostaglandin, cimetidine and probanthine. *Gastroenterology* 1979;76:88-93.
- 9 Skillmann JJ, Bushnell LS, Goldman H, Silen W. Respiratory failure, hypotension, sepsis and jaundice: a clinical syndrome associated with lethal haemorrhage from acute stress ulceration of the stomach. *Am J Surg* 1969;117:523.
- 10 Hastings PR, Skillman JJ, Bushnell LS, Silen W. Antacid titration in the prevention of acute gastrointestinal bleeding. *N Engl J Med* 1978;298:1041-5.
- 11 Zeldis JB, Friedman LS, Isselbacher KJ. Ranitidine: a new H₂ receptor antagonist. *N Engl J Med* 1983;309:1368-73.
- 12 Anonymous. Ranitidine (Zantac). *Med Lett Drugs Ther* 1981;24:111-3.
- 13 Dundee JW, McGowan WA, Elwood RJ, Hildebrand PJ. Plasma benzodiazepine concentrations following oral administration with and without H₂ receptor blockers. *Ir J Med Sci* 1982;151:413-4.
- 14 Spahn H, Mutschler E, Kirch W, Ohnhaus EE, Janisch HD. Influence of ranitidine on plasma metoprolol and atenolol concentrations. *Br Med J* 1983;286:1546-7.
- 15 Van Saene HKF, Stoutenbeck Ch P, Miranda DR, Zandstra DF. A novel approach to infection control in the intensive care unit. *Acta Anaesthesiol Belg* 1983;3:193-208.
- 16 Schuster DP, Rowley H, Feinstein S, McGue MK, Zuckerman GR. Prospective evaluation of the risk of upper gastrointestinal bleeding after admission to a medical intensive care unit. *Am J Med* 1984;76:623-30.
- 17 McCarthy DM. Ranitidine or cimetidine. *Ann Intern Med* 1983;99:551-3.
- 18 Harvey CJ, Rampton DS. Prostaglandins and the gastrointestinal mucosa are they important in its function, disease or treatment? *Gastroenterology* 1985;89:1162-88.