CORRESPONDENCE

Postoperative thromboembolic disease and oral contraceptives N Puttick, FFARCS, and Melanie Jones, FFARCS	387	Prevention of recurrent acute cystitis by methenamine hippurate W Brumfitt, FRCPATH, and J M T Hamilton-Miller; S Cronberg, MD, and others.....	390	Secrecy and distinction awards S Bourne, FRCPsych, and P Bruggen, FRCPsych	393
Pyomyositis associated with hepatitis Lindsay Dow, MRCP, and others	387	Converting enzyme inhibition and kidney function in normotensive diabetic patients with persistent microalbuminuria P H Winocour, MRCP, and others	391	Points Telling patients about their medicines (C W Koenigsberger); Evaluation of portable haemoglobinometer in general practice (R Quinn); Lymphoedema of the arm (Helen J Stewart, J D E Knox).....	393
Delivery after caesarean section P Maouris, MB, and P Macrow, MB; R P Gleeson, MRCOG, and others	388	Loin pain and haematuria syndrome S M W Hutchison, MRCP; Y T Kontinen, MD, and others.....	391	Elective surgery in rural east Africa (T G Harland); Exploring therapeutic risks (G Venning); Mortality from myocardial infarction in different types of hospitals (W James and S Dharmavaratha); Resuscitating resuscitation (J F Searle); Diagnosing pulmonary thromboembolism (S W Watkin and C C Evans); Measles: who pays the cost? (Allison Streetly); Contamination of dropper bottles with tear fluid in an ophthalmic outpatient clinic (B W Cromie); Preventing alcohol and tobacco abuse (A S Amsden); Alcohol consumption and the risk of alcohol related cirrhosis (K G Green).....	394
Randomised comparison of early versus late induction of labour in post-term pregnancy P T Mohide, FRCS; P Bergsjø, MD	388	AIDS: a faltering step I Mohammed, FRCP, and others	392		
Extracranial-intracranial bypass, one; clinical trials, nil H A F Dudley, FRCS	389	After safe sex, safe surgery? K Lafferty, FRCS, and A P Wyatt, FRCS	392		
Microdiets M J Hall, MRCP; J Marks, FRCP	389	Halothane hepatitis in children J F Mayhew, MD	392		
Nuclear magnetic resonance imaging C E Bennett, FRCP, and R D Hoare, FRCP	390	The cost of nursing K W M Scott, FRCPATH.....	393		
		References for junior doctors P G P Lawler, FFARCS	393		

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

Postoperative thromboembolic disease and oral contraceptives

SIR,—The relation between postoperative thromboembolic disease and the combined oral contraceptive pill is now well established.¹⁻⁵ Guidelines have been suggested by Guillebaud¹ and incorporated into the *British National Formulary* in order to correct misunderstandings and minimise the risk of unwanted pregnancy.⁶ In addition, the subject has been extensively discussed in medical correspondence.⁷⁻⁹

Many anaesthetists, however, have noted with concern that there is still considerable confusion about this matter. Women admitted for major surgery continue to arrive in hospital still using combined oral contraception; conversely, many other women have their minor surgery unnecessarily postponed, often for three months. The anaesthetist is often expected to resolve this problem at a preoperative visit.

Both circumstances are avoidable. It is clearly the responsibility of the surgeon to elicit the relevant history in the outpatient department. When major surgery is planned the patient should be counselled and given at least one month's notice of admission. This should not be impossible in view of the fairly small proportion of admissions accounted for by this population. In the case of minor surgery the patient can be reassured that any increase in risk is minimal in comparison with that of pregnancy. Whatever advice is given should also be included in correspondence to the patient's general practitioner. Junior surgical staff can be instructed on this topic in a matter of minutes, and such tuition would need repeating only every six months.

Why, then, is there still a problem? We believe

that the answer is threefold: firstly, lack of knowledge; secondly, failure to organise an efficient scheme for detecting, advising, and admitting these patients; and, thirdly, apprehension engendered by a case reported in the Medical Defence Union's annual report for 1986.¹⁰ This has been widely misinterpreted as implicating a contraceptive pill in the occurrence of a brainstem thrombosis under anaesthesia. Careful reading of the report, however, shows that no such thrombosis occurred, and when we contacted the Medical Defence Union for further details we were told that the settlement made no reference to the contraceptive pill.

Postoperative thromboembolic disease is primarily a complication of surgery; we suggest that the onus of counselling the patient, or cancelling the operation, should no longer be placed on the anaesthetist. The evening before the list is not the time to discuss such a matter; the patient suffers needless anxiety, and no useful contribution to her management can be made.

NIGEL PUTTICK
MELANIE JONES

Department of Anaesthetics,
University of Wales College of Medicine,
Cardiff CF4 4XW

- Guillebaud J. Surgery and the pill. *Br Med J* 1985;291:498-9.
- Stadel BV. Oral contraceptives and cardiovascular disease. *N Engl J Med* 1981;305:612-8.
- Stadel BV. Oral contraceptives and cardiovascular disease. *N Engl J Med* 1981;305:672-7.
- Sagar S, Stamatakis JD, Thomas DP, Kakkar VV. Oral contraceptives, antithrombin III activity, and postoperative deep vein thrombosis. *Lancet* 1976;i:509-11.
- Vessey MP, Doll R, Fairbairn AS, et al. Postoperative thrombo-

embolism and the use of oral contraceptives. *Br Med J* 1970;iii:123-6.

- Canter RJ, Pryce J. Risk of pregnancy while waiting for an operation. *Br Med J* 1985;291:516.
- Waldram M. Surgery and the pill. *Br Med J* 1985;291:742.
- Dyson PHP. Surgery and the pill. *Br Med J* 1985;291:899.
- Guillebaud J. Surgery and the pill. *Br Med J* 1985;291:1201.
- Medical Defence Union. Annual report. London: Medical Defence Union, 1986:19.

Pyomyositis associated with hepatitis

SIR,—Dr R A Watts and colleagues (13 June, p 1524) describe a case of pyomyositis in a patient with antibodies to the human immunodeficiency virus; we recently saw a patient with pyomyositis associated with hepatitis A infection, which has also not previously been reported.

A 21 year old heterosexual naval engineer was admitted with fever, hepatosplenomegaly, and painful swelling of the right thigh. He had just returned from a four month voyage to the near east, during which six crew members had fallen ill with infectious hepatitis. Five of them made uncomplicated recoveries, but our patient, two weeks after the onset of jaundice, developed bilateral thigh pain and swelling. The left thigh improved spontaneously, but the right thigh became more painful, and four weeks after the onset of symptoms he could not bear weight on the right leg. There was no history of trauma, skin lesions, or intramuscular injections. On examination he was feverish (38.5°C) and had moderate hepatosplenomegaly and a swollen, woody, and hard right thigh. A test for hepatitis A IgM yielded