Proceedings of the American Society of Bariatric Surgery (in press).

- 6 McCarthy HB, Rucker RD, Chan EK, *et al*. Gastritis after gastric bypass surgery. *Surgery* 1985;98:68-71.
- 7 Morson BC, Sobin LH, Grundmann E, *et al*. Precancerous conditions and epithelial dysplasia in the stomach. *J Clin Pathol* 1980;33:711-21.
- 8 Park HK, Sinar DR, Sloss RR, *et al*. Histologic and endoscopic studies before and after gastric bypass surgery. *Arch Path Lab Med* 1986;110:1164-7.

AUTHORS' REPLY.—Mr Owen and colleagues are correct that we did not directly refer to the shortened lifespan in gross obesity because no operation has yet been shown to reverse this trend successfully. We did not discuss the Scopinaro biliopancreatic bypass because only its originator has substantial experience of it.

We were perhaps a little uncritical in quoting reports of dysplasia and gastritis in some of the gastric procedures. But no other reports on the Alden procedure have been published so far that refute the assertions of McCarthy *et al*. We remain unconvinced, however, that gastric operations for gross obesity are a satisfactory solution; the small proximal pouch (50 ml) impairs normal eating for the rest of the patient's life. In a masterly review Näsund compared gastric bypass surgery and gastroplasty and pointed out that little is known about the physiology of these operations.¹ The several published modifications to these surgical techniques point to the difficulty of obtaining consistently good results.

We take issue with Mr Owen and colleagues over the ease of reversing gastric bypass. In our experience great difficulty has occurred, with not insignificant morbidity.² Unfortunately, five years' follow up is not enough to draw firm conclusions. Jejunoileal bypass remains a very successful operation even after 10 years, and only after that time was it abandoned. So once again we advise patience before drawing conclusions.

J-C GAZET
T R E PILKINGTON

St George's Hospital and Medical School,
London SW17 0RE

- 1 Näsund I. Gastric bypass versus gastroplasty. *Acta Chir Scand* 1987;suppl:536.
- 2 Gazet J-C. Surgery for morbid obesity. *Current Opinion in Gastroenterology* 1986;3:896-8.

Type A behaviour and coronary heart disease

SIR.—Dr Tom Sensky's cautious optimism about type A behaviour (11 July, p 69) has been confirmed by a recent meta-analysis that examined all studies relating type A behaviour to coronary heart disease published between 1950 and 1985.¹

The authors showed that the effect size (Pearson's *r*) for all measures of type A behaviour and all coronary heart disease groups was 0.136 ($p < 0.0001$). This means that type A behaviour accounts for about 2% of the variance in coronary heart disease. Although this is a small effect size, the reliability of the finding and the number of people affected make it non-trivial. The more detailed results of the study were instructive and included highly significant effect sizes on coronary heart disease for the following personality indices: Jenkins activity survey, 0.067; type A structured interview, 0.221; extraversion, 0.071; anxiety, 0.124; and depression, 0.205.

Other results from the same study were relevant to Dr Sensky's concerns about the relation between the structured interview and self report measures: the "hard driving" component of the Jenkins activity survey had an effect size of 0.153, larger than either the type A (all measures) or the total Jenkins activity survey score effect sizes. Further, "job involvement" on the Jenkins activity survey was not related to coronary heart disease

while "speed and impatience" were weakly related, leading the authors to conclude that this finding seemed to contradict the popular notion of the workaholic who develops coronary heart disease.

Both Dr Sensky's and Booth-Kewley's and Friedman's studies fail to emphasise that those who have developed the type A concept have been reinventing the wheel. As Eysenck said, "proponents of the concept often argue and write as if well established models of personality did not exist and fail to relate their concepts to these dimensions."² Fortunately, this is beginning to change and there are now reports that type A behaviour is related to neuroticism, trait tension, anxiety, depression, and extraversion.^{2,3} If these obvious empirical investigations had been done earlier we might have discovered before now that a single component of the Jenkins activity survey predicts coronary heart disease better than the whole scale.

Sensky comments that the type A concept is not a unitary one, and, indeed, this conclusion has occurred to psychologists concerned with a scale which, superficially, appears to describe a "type."^{2,4} Eysenck, for example, concluded, "Clearly type A behaviour is not unitary; certain aspects of it are closely related to neuroticism, others to extraversion and the two groups of type A behaviour traits are themselves almost unrelated. . . . In this general description, it will be noted, there is no trace remaining of the concept of type A behaviour as such; the concept has been shown to be a chimera, stemming from perfectly correct observations of the originators of the concept, followed by psychometrically inappropriate analysis, and disregard of much better established personality dimensions."²

In a discipline that now accepts psychology as one of its basic sciences these are embarrassingly elementary lessons. The finding that personality predispositions relate to coronary heart disease is important, and its investigation requires an awareness of the appropriate methods and skills.

IAN J DEARY

Department of Psychology,
University of Edinburgh,
Edinburgh EH8 9JZ

- 1 Booth-Kewley S, Friedman HS. Psychological predictors of heart disease: a quantitative review. *Psych Bull* 1987;101:343-62.
- 2 Eysenck HJ, Fulker D. The components of type A behaviour and its genetic determinants. *Personality and Individual Differences* 1983;4:499-505.
- 3 Langeluddecke PM, Tennant CC. Psychological correlates of the type A behaviour pattern in coronary angiography patients. *Br J Med Psychol* 1986;59:141-8.
- 4 Ray JJ. Confusions in defining 'A-B' personality type: a rejoinder to Jenkins and Zyzanski. *Br J Med Psychol* 1984;57:385.

Managing psychopathic offenders

SIR.—Dr Derek Chiswick does not appear to have addressed the fundamental question of whether psychopaths are suffering from some form of mental abnormality that actually requires treatment—however ineffective current treatment may be (18 July, p 159). He is not correct when he states that psychopaths are not treated with electroconvulsive therapy, as a preliminary analysis of data on 131 psychopaths (men and women) in Broadmoor and Rampton Hospitals found that 29% had received electroconvulsive therapy at some time in their lives, with 55% previously admitted to ordinary psychiatric hospitals, not infrequently on a hospital order, but invariably under the legal category of mental illness (J W Coid, Royal College of Psychiatrist's conference, 1987). One in four women and one in five men had suffered a schizophrenic breakdown (usually short lasting) and 61% had had a depressive illness during their lifetime. These patients tended to

receive diagnoses of personality disorder (DSM-III multiple axis 2). Their histories also showed a surprisingly high incidence of early cerebral trauma, emotional and physical deprivation, and depression and personality disorder in their first degree relatives.

Some of the issues seem confused, with one argument made against another problem. For example, Dr Chiswick is right to point out the unsatisfactory situation of mental health review tribunals discharging dangerous psychopaths back into the community, but this is not directly relevant to their need for treatment in a therapeutic setting. Risk to the public from future offending may well be high when psychopaths were compared with mentally ill subjects in the studies quoted, but none compared a similar group of personality disordered subjects released from prison without treatment. It may be premature to conclude from the statistical associations of the study by Dell *et al*¹ that the discharge of psychopaths relates merely to the seriousness of the offence while that of schizophrenics relates to the chronicity of illness. It is also unfair to describe the endeavours of the small group of special hospital staff who are actively concerned with psychopaths as mere "time fillers."

Certainly many legal aspects of the detention of psychopaths outlined by Dr Chiswick are not entirely satisfactory, and treatment programmes for these patients are sometimes patchy and poorly organised. However, careful unbiased research into their treatment and its effectiveness is badly needed. Perhaps more importantly, research is needed into what is actually being treated. Reducing the numbers of psychopaths detained in psychiatric hospitals does not relate to their need for treatment but to how staff feel about attempting treatment. In the case of the state hospital at Carstairs it most probably related to the terrible events of the past.² Dr Chiswick is right to describe the management of psychopathic disorders as a problem that will not go away. Excluding them from hospital merely dumps highly disturbed individuals into an already overcrowded prison system with no hope of treatment before their return to the community.

JEREMY COID

Maudsley Hospital,
London SE5 8AZ

- 1 Dell S, Robertson G, Parker E. Detention in Broadmoor: factors in length of stay. *Br J Psychiatry* 1987;150:824-7.
- 2 Scottish Home and Health Department. *State hospital, Carstairs. Report of public local inquiry into circumstances surrounding the escape of two patients on 30 November 1976 and into security and other arrangements at the hospital.* Edinburgh: HMSO, 1977.

Prediction of resources needed for treatment of renal failure

SIR.—We are aware of the project of Ms Ruth Davies and Mr Huw Davies (25 July, p 271) and hope that their model will produce precise predictions of value. Unfortunately, no such predictions are available and no comparisons may be drawn. The superiority of the model they propose awaits corroboration other than that provided by references to their own work. Furthermore, there is debate about the relative merits of mathematical models. We chose to use a steady state approach to avoid the difficulties that arise if too detailed an attempt is made to give year by year numbers and costs for a renal replacement programme. We are satisfied that our approach provides a minimal target of need as we have used the most favourable implications for outcome in every case—for example, with regard to transplant availability, the proportion of patients suitable for independent dialysis, and the costs for different methods of treatment including continuous ambulatory peritoneal dialysis.

It is important that debate about modelling does not obscure the practical implications of the predictions on which all are agreed: a considerable increase in resources for the treatment of renal failure will be required over the next decade solely to keep pace with the numbers of patients predicted from the target of 40 new patients per million population per year. Overall survival is now so good that the stock of patients kept alive with treatment will not level out before the end of the century. In addition, acceptance rates are expected to continue to rise. By 1985 numbers accepted already exceeded 40 per million population in 17 other European countries and were over 50 in eight of them and over 60 in two.¹

Since our prediction of dialysis requirements is minimised by assuming free availability of kidneys for grafting, failure in the supply of donor organs will hold up attainment of steady state and shift more patients into (more expensive) dialysis modes of treatment.

N P MALLICK

Manchester Royal Infirmary,
Manchester

I T WOOD

Crewe and Alsager College of Higher Education,
Crewe

MARY MCGEOWN

Belfast City Hospital,
Belfast BT9 7AB

A J WING

St Thomas's Hospital,
London

1 Broyer M, Brunner FP, Brynger H, et al. EDTA registry centre survey, 1985. *Nephrology Dialysis and Transplants* (in press).

Treatment of recurrent spontaneous abortion

SIR,—Dr Willem Vlaanderen and Professor Pieter E Treffers reported the outcome of fourth and later pregnancies in 24 women with a history of three consecutive miscarriages and concluded that no specific treatment should be offered to such couples (11 July, p 92). The spontaneous success rate of 80% is surprising in view of the data reported in other (larger) series.^{1,2} Habitual aborters are a heterogeneous group, and those women who have simply been unlucky may have been over-represented in this small group. The fact that either three or all four (it was not made clear) of the women miscarried in their fourth pregnancies supports that suggestion. Be that as it may, the authors' advice about treatment should be received with caution.

Women experiencing repeated pregnancy loss commonly complain that they are simply told that they should just try again because there is a good chance they will be luckier next time. While such a statement may be true, the patient sometimes considers that it reflects a lack of understanding, and it does nothing to alleviate depression or apprehension about future pregnancies. In this context any treatment may be considered better than nothing so long as it is not harmful. Tender loving care or its extension, psychotherapy,³ is both helpful and harmless and should be practised irrespective of other treatment. Immunotherapy by infusion or vaccination of blood components is more controversial and undoubtedly carries some risk, but encouraging results have been reported by several independent groups.^{4,6}

We have been treating selected women with this problem with two injections of their husbands' mononuclear cells, given before pregnancy and shortly after conception. The risks are small and probably acceptable. This treatment may be no more effective than tender loving care, but it is of obvious psychological benefit to the patients.

Moreover, all six of our patients who have completed the treatment either have delivered or are at an advanced stage of pregnancy. Possibly all idiopathic aborters will succeed eventually, but any treatment which might reduce the number of antecedent failures deserves consideration.

D C KILPATRICK

Blood Transfusion Centre,

W A LISTON

Gynaecology Pavilion,
Royal Infirmary,
Edinburgh EH3 9HB

- 1 Poland BJ, Miller JR, Jones DC, Trimble BK. Reproductive counselling in patients who have had a spontaneous abortion. *Am J Obstet Gynecol* 1977;199:685-91.
- 2 Beer AE, Semprini AE, Xiaoyu Z, Quebbeman JF. Pregnancy outcome in human couples with recurrent spontaneous abortions: HLA antigen profiles; HLA antigen sharing; female serum MLR blocking factors; and paternal leukocyte immunisation. *Experimental Clinical Immunogenetics* 1985;2:137-53.
- 3 Stray-Pedersen B, Stray-Pedersen S. Etiologic factors and subsequent reproductive performance in 195 couples with a prior history of habitual abortion. *Am J Obstet Gynecol* 1984;148:140-6.
- 4 Mowbray JF, Liddell H, Underwood JL, Gibbings C, Reginald PW, Beard RW. Controlled trial of treatment of recurrent spontaneous abortion by immunisation with paternal cells. *Lancet* 1985;i:941-3.
- 5 Taylor CG, Faulk WP, McIntyre JA. Prevention of recurrent spontaneous abortions by leukocyte transfusions. *J R Soc Med* 1985;78:623-7.
- 6 Unander AM, Lindholm A. Transfusions of leukocyte-rich erythrocyte concentrates: a successful treatment in selected cases of habitual abortion. *Am J Obstet Gynecol* 1986;154:516-20.

Fatal bronchospasm after topical lignocaine before bronchoscopy

SIR,—The short report by Dr S P Ruffles and Dr J G Ayres (27 June, p 1658) reveals several important errors of management. It is possible, as the authors suspected, that their patient's bronchospasm was a hypersensitivity reaction to lignocaine, but the fundamental reasons for his death were (a) that he was being subjected to an unnecessary and, in his case, potentially dangerous investigation (fiberoptic bronchoscopy), (b) that no effort was apparently made to correct fairly severe hypoxaemia before the procedure was started, and (c) that the management of his cardiorespiratory arrest was inadequate.

The indication for bronchoscopy was stated to be weight loss suggesting a diagnosis of lung cancer. A chest radiograph had, however, shown "no suggestion of a carcinoma." It was thus highly improbable that the weight loss could have been due to intrathoracic malignant disease. Even if there had been radiological evidence of tumour, fiberoptic bronchoscopy, with its known risk of aggravating hypoxaemia,¹ would have been hazardous in a patient with a forced expiratory volume of 0.5 l, a PaO₂ of 7.6 kPa, and a PaCO₂ of 6.4 kPa. It is also questionable whether a patient with such a degree of ventilatory insufficiency and respiratory failure would have derived any real benefit from treatment for bronchial carcinoma had that improbable diagnosis been confirmed.

Although there is some ambiguity about the sequence of electrocardiographic changes, it would appear that he first developed hypoxic cardiac arrest in asystole, ventricular fibrillation occurring only as an agonal event after the administration of adrenaline and calcium chloride. In these circumstances cardioversion is futile, and unless oxygenation can be rapidly restored by intubation and ventilation with 100% oxygen death is inevitable. Manual ventilation was difficult, no doubt because of severe bronchospasm, and the patient's by then profound hypoxaemia was presumably never even partially corrected. Gross pulmonary hyperinflation caused by a combination

of bronchospasm and emphysema would also have prevented external cardiac massage from achieving an adequate cardiac output. Only open cardiac massage might conceivably have averted a fatal outcome, although such a drastic procedure could hardly have been justified in a man who must already have been a respiratory cripple.

Finally, I refer the authors to the following passage in a leading article I contributed to the *BMJ* less than a year ago²: "Perhaps too many high risk patients—the elderly, the frail, and those with advanced chronic obstructive airways disease—are now being subjected to an invasive investigative procedure that is unlikely to influence management. Respiratory physicians should not forget that bronchoscopy in these patients may be dangerous as well as pointless." The only amendment I might now want to make to that passage would be to omit the word "perhaps."

IAN W B GRANT

Kirkcubbin,
West Lothian EH27 8EA

- 1 Salisbury BG, Metzger LF, Altose MD, Stanley NN, Cherniak NS. Effect of fiberoptic bronchoscopy on respiratory performance in patients with chronic airways obstruction. *Thorax* 1975;30:441-6.
- 2 Grant IWB. Hazards of bronchoscopy. *Br Med J* 1986;293:286-7.

AUTHORS' REPLY.—Dr Grant raises several points worthy of comment, though they are somewhat peripheral to the main reason for our reporting this case. We are glad that Dr Grant believes that the patient's death was a hypersensitivity reaction to the lignocaine.

Although this man had a low forced expiratory volume in one second he had premedication with serial doses of nebulised salbutamol and was given oxygen on the ward, although this was not made clear in our short report. The reason for bronchoscopy was weight loss and increasing breathlessness coupled with the fact that he was very anxious to find the cause. It is well recognised that a central small carcinoma of the bronchus can be invisible in a plain chest radiograph and can cause marked symptoms of increasing breathlessness by partially occluding the main bronchus. Increasing dyspnoea often has an anorectic effect because of the sheer effort involved in mastication and deglutition. For these reasons we contemplated bronchoscopy, as radiotherapy to such a lesion could relieve the breathlessness. In the past we have performed bronchoscopy in many patients with a forced expiratory volume of below 1 litre, this, in our opinion, not being an absolute contraindication to bronchoscopy where future management may be affected. As far as the management of his cardiac arrest is concerned, clearly it is difficult for a man with such lung disease to recover, but this should not stop our attempts to resuscitate him.

Dr Grant's original final paragraph in his leading article on the hazards of bronchoscopy need not be altered. The indications for performing bronchoscopy in this patient were clear cut. The possibility of a central tumour, though rare, was real and if one had been found radiotherapy would have conferred symptomatic benefit in the patient's final months of life.

JON G AYRES
S P RUFFLES

Department of Respiratory Medicine,
East Birmingham Hospital,
Birmingham B9 5ST

SIR,—Dr Said Abdallah's dogma that patients whose forced expiratory volume in one second is less than 1 litre are unfit for bronchoscopy (15 August, p 444) should not go unchallenged. I am sure there are many fiberoptic bronchoscopists