eventually resolved completely. Even the most persistent proteinuria resolved, despite most patients having various recognisable glomerular lesions. The message appears reassuring and simple: on finding proteinuria stop gold treatment, give a high protein diet with diuretics if needed, and wait. Systemic steroids are not required routinely. They are indicated only when the patient has life threatening proteinuria; they are probably best given as pulsed doses of intravenous methylprednisolone followed by a short course of oral methylprednisolone at a dose of 80-100 mg on alternate days until the proteinuria is controlled.

Proteinuria after gold treatment is less common than skin and haematological toxicity, but its possible implications have been a worry to doctors and patients. These new findings, which suggest a "sitting on hands approach" to

proteinuria caused by gold while nature effects a cure, are reassuring.

A J COLLINS

Senior Lecturer in Pharmacology,
University of Bath, and
Royal National Hospital for Rheumatic Diseases,
Bath BA1 1RL

- 1 Empire Rheumatism Council. Gold therapy in rheumatoid arthritis. Report of a multicentre controlled trial. *Ann Rheum Dis* 1960;19:95-119.
- 2 Co-operating Clinics Committee of the American Rheumatism Association. A controlled trial of gold salt therapy in rheumatoid arthritis. *Arthritis Rheum* 1973;16:353-8.
- 3 Eigler JW, Bluhm GB, Duncan H, Sharp JT, Ensign DC, McCrum WR. Gold salts in the treatment of rheumatoid arthritis. A double-blind study. *Ann Intern Med* 1974;80:21-6.
- 4 Luukkainen R, Kajander A, Isomaki M. Effect of gold in progression of erosions in rheumatoid arthritis. Better results with early treatment. *Scand J Rheumatol* 1977;6:189-92.
- 5 Blodgett RC, Heuer MA, Pietrusic RG. Auranofin: a unique oral chrysotherapeutic agent. *Semin Arthritis Rheum* 1984;13:255-73.
- 6 Girdwood RH. Death after taking medicaments. *Br Med J* 1974;ii:501-4.
- 7 Silverberg DS, Kidd EG, Shintka TH, Olan RA. Gold nephropathy: a clinical and pathological study. *Arthritis Rheum* 1970;13:812-25.

Action on alcohol—at last

Those worried by Britain's alcohol problems have had little to cheer them for a long time. But last week's announcement that a committee of ministers is to consider all aspects of alcohol abuse is very good news indeed. Such a committee was recommended in May 1979 by the suppressed (and thereafter repeatedly leaked) "think tank" report.¹ The report recognised that when 16 government departments had an interest in alcohol forging a coherent policy on alcohol would be impossible without interdepartmental liaison. Many people thought that more than any other this recommendation caused the report to be suppressed: the government wanted to have its cake and eat it, with some departments interested in boosting alcohol production, the Treasury gratefully filling its coffers with the proceeds of alcohol taxes, and the Home Office and the Department of Health and Social Security wringing their hands and bemoaning the crime and sickness associated with alcohol. An interdepartmental committee would have highlighted these conflicts, and the new committee will have to acknowledge them.

The ministerial committee might well begin its work by dusting off the think tank report. No other government report has analysed the problem anything like so clearly. And one thing that the committee will discover from the report is that it will be unproductive to think simply in terms of "the

minority who abuse alcohol." The Home Secretary, when he announced the setting up of the committee last week, talked of "misuse of alcohol by a minority [having] serious consequences for the health and social services." But most of the evidence shows that the hundreds of problems associated with alcohol cannot be controlled by concentrating on a few hooligans, criminals, and drunken drivers.^{2,4} Much of the harm done by alcohol is the result of patterns of drinking that are widespread among the population, and an effective policy on alcohol would have an impact on most of us. Virtually all of the many groups that have looked at responding to alcohol problems have recognised that a wide range of responses is necessary and that most important are measures to contain and probably reduce national alcohol consumption.^{1,2,5} Raising the price of alcohol is the best way to do this. We will look to next spring's budget to see how seriously the interministerial committee is taking Britain's alcohol problems.

RICHARD SMITH

Assistant editor, *BMJ*

- 1 Brunn K. *Alcohol policies in the United Kingdom*. Stockholm: Sociologiska Institutionen, 1982.
- 2 Special Committee of the Royal College of Psychiatrists. *Alcohol: our favourite drug*. London: Tavistock Publications, 1986.
- 3 Kreitman N. Alcohol consumption and the preventive paradox. *Br J Addict* 1986;81:353-63.
- 4 Kendall RE. Drinking sensibly. *Br J Addict* (in press).
- 5 Scottish Health Education Co-ordinating Committee. *Health education in the prevention of alcohol related problems*. Edinburgh: Scottish Home and Health Department, 1985.