- 12 Wakamatsu E, Sissons HA. The cancellous bone of the iliac crest. *Calcified Tissue Research* 1969;4:147-61.
- 13 Parfitt AM, Mathews CHE, Villanueva AR, Kleerekoper M, Frame B, Rao DS. Relationships between surface, volume, and thickness of iliac trabecular bone in aging and in osteoporosis. Implications for the microanatomic and cellular mechanisms of bone loss. *J Clin Invest* 1983;72:1396-409.
- 14 Reeve J. A stochastic analysis of iliac trabecular bone dynamics. *Clinical Orthopaedics and Related Research* 1986;213:264-78.
- 15 Lips P, Courpron P, Meunier PJ. Mean wall thickness of trabecular bone packets in the human iliac crest: changes with age. *Calcified Tissue Research* 1978;26:13-7.
- 16 Eriksen EF, Melsen F, Mosekilde L. Reconstruction of the resorption site in iliac trabecular bone: a kinetic model of bone resorption in 20 normal individuals. *Metab Bone Dis Relat Res* 1984;5:235-42.
- 17 Courpron P, Lepine P, Arlot M, Lips P, Meunier PJ. Mechanism underlying the reduction with age of the mean wall thickness of trabecular basic structure unit (BSU) in human iliac bone. *Metab Bone Dis Relat Res* 1980; suppl 2s:323-9.
- 18 Reeve J, Hesp R, Wootton R. A new tracer method for the calculation of rates of bone formation and breakdown in osteoporosis and other generalised skeletal disorders. *Calcified Tissue Research* 1976;22:191-206.
- 19 Deacon AC, Hulme P, Hesp R, Green JR, Tellez M, Rinsler MG, Reeve J. The estimation of whole body bone resorption rate biochemically: a comparison of urinary total hydroxyproline excretion with two radioisotopic tracer methods in patients with osteoporosis. *Clin Chim Acta* 1987;166:297-306.
- 20 Ettinger B, Genant HK, Cann CE. Postmenopausal bone loss is prevented by low-dosage estrogen with calcium. *Ann Intern Med* 1987;106:40-5.
- 21 Genant HK, Cann CE, Boyd DP, Kolb FO, Ettinger B, Gordan GS. Quantitative computed tomography for vertebral mineral determination. In: Frame B, Potts JT, eds. *Clinical disorders of bone and mineral metabolism*. Amsterdam: Excerpta Medica, 1983:40-7.
- 22 Krølner B, Nielsen SP. Bone mineral content of the lumbar spine in normal and osteoporotic women: cross-sectional and longitudinal studies. *Clin Sci* 1982;62:329-36.
- 23 Lindquist O, Bengtsson C, Hansson T, Roos T. Bone mineral content in relation to age and menopause in middle aged women. A study of bone density in lumbar vertebrae by dual photon absorptiometry in a population sample of women. *Scand J Clin Lab Invest* 1981;41:215-23.
- 24 Riggs BL, Wahner HW, Dunn WL, Mazess RB, Offord KP, Melton LJ. Differential changes in bone mineral density of the appendicular and axial skeleton with aging. Relationship to spinal osteoporosis. *J Clin Invest* 1981;67:328-35.
- 25 Riggs BL, Wahner JW, Melton LJ, Richelson LS, Judd HL, Offord KP. Rate of bone loss in the appendicular and axial skeletons of women. Evidence of substantial vertebral bone loss before menopause. *J Clin Invest* 1986;77:1487-91.
- 26 Garn SM. *The earlier gain and the later loss of cortical bone in nutritional perspective*. Springfield, Illinois: Thomas, 1970.
- 27 Christiansen C, Rüs BJ, Rødbro P. Prediction of rapid bone loss in postmenopausal women. *Lancet* 1987;i:1105-8.

(Accepted 20 May 1987)

Lesson of the Week

Vitamin and energy injections in the Indian subcontinent

ELIZABETH H BOXALL

Though it is now common for general practitioners to request immunoglobulin for protection against hepatitis A for travellers visiting the Indian subcontinent, probably more should be done to protect patients against possible exposure to hepatitis B. This is a particular problem with Asian patients who go abroad for extended periods and are therefore more likely to require medical attention while there. Twice during the past year we have diagnosed acute hepatitis B in young women who had received injections while in Pakistan.

Case histories

Case 1—A 26 year old Asian woman normally resident in Britain visited Pakistan for six weeks and returned five weeks before admission. While in Pakistan she was reported to have had an "energy injection." Her two children were in hospital in Birmingham being treated for typhoid when she was taken ill with abdominal pain and fever and admitted with suspected typhoid. She was 16 weeks pregnant. Liver function tests yielded a serum aspartate transaminase activity of 4300 IU/l with a bilirubin concentration of 46 $\mu\text{mol/l}$, and hepatitis B surface antigen was present to a titre of $>1/8000$ by reverse passive haemagglutination (Hepatest, Wellcome Laboratories). Four days after admission to this hospital she was transferred to King's College Hospital, London, with a diagnosis of fulminant hepatic failure due to infection with hepatitis B virus, and she died four days later.

Case 2—A 20 year old Asian woman normally resident in Britain until her marriage in Pakistan in 1984 returned from a visit lasting 12 months in December 1986. During her stay she had suffered persistent diarrhoea which had resulted in weight loss for which she had had several "vitamin" injections from local doctors. She was admitted less than a month after her return with diarrhoea, vomiting, anorexia, and jaundice. On admission her serum aspartate transaminase activity was 1545 IU/l and bilirubin concentration 152 $\mu\text{mol/l}$. Hepatitis B surface antigen was present to a titre of 1/32 (Hepatest, Wellcome). She recovered well and was discharged two weeks later for outpatient follow up.

Regional Virus Laboratory, East Birmingham Hospital, Birmingham B9 5ST
ELIZABETH H BOXALL, PHD, MRCPATH, principal virologist and deputy director

Patients should be warned about the infectious risks of "traditional medicines" given by injection while abroad

Discussion

Both patients had IgM class antibody to hepatitis B core antigen, indicating a diagnosis of acute hepatitis B acquired within the previous two to six months. The most likely source of infection in both patients was the injections that they had received. Energy and vitamin injections are presumably not essential to life. Though the local population and family of the visitors may speak highly of such injections, these people have the advantage of probably being naturally immune to hepatitis B through previous exposure. The risk is probably not in the vitamin preparation itself but in the use of inadequately sterilised equipment, especially needles.

Hepatitis is not uncommon in Asians returning from the Indian subcontinent. In a study from East Birmingham Hospital of 50 consecutive cases of acute hepatitis, 16 were in Asians returning from the Indian subcontinent.¹ Three of these were diagnosed as hepatitis B, five were hepatitis A, and eight were non-A non-B.

Cases of hepatitis B such as those described here are avoidable. General practitioners should be especially vigilant in warning their patients of the dangers of accepting traditional medicines given by injection while they are abroad. The risks of infection with the human immunodeficiency virus by the same route may serve to emphasise this message more strongly.

Reference

- 1 Busher GL, Skidmore SJ, McKendrick MW, Geddes AM. Sporadic nonA-nonB hepatitis in Birmingham. *Journal of Infection* 1981;3:45-9.

(Accepted 10 July 1987)