Basic Molecular and Cell Biology

Cell reproduction

PAUL NURSE

The sequence of events and processes leading to cell reproduction is known as the cell cycle. During the cell cycle there is generally a duplication of cellular components followed by their partitioning into two daughter cells at cell division. For many components present in moderate to large amounts within the cell this duplication and partition need not be controlled exactly. The chromosomes, however, must be precisely replicated and segregated during each cell cycle to produce two viable daughter cells. Replication of the deoxyribonucleic acid (DNA) making up the chromosomes occurs during S phase, and segregation of the chromosomes occurs during mitosis. These are the two major events of all cell cycles. In most cell cycles there are also gaps between these events: a G1 gap after mitosis and before S phase and a G2 gap after S phase and before mitosis.

When the cells are not actively undergoing reproduction they exit the cell cycle and enter a quiescent state. This exit usually occurs from G1, and the quiescent state is called G0. Quiescent cells may be metabolically active, but they do not grow and do not undergo any of the events and processes of the cell cycle. The shift between the G0 quiescent state and the cell cycle plays an important part in the growth and maintenance of the organism. Regulated entry into the cell cycle is necessary for proper development and for processes such as wound healing and tissue replenishment. If regulation becomes defective it may result in uncontrolled cell reproduction, leading to cancer.

I describe here some recent progress in our understanding of how the transition from quiescence to the cell cycle is controlled. I also consider briefly two further controls over cell reproduction which act during the cell cycle, one in late G1 before S phase and a second at the transition from G2 to mitosis.

Growth factors and their receptors

Mammalian cells may be grown in cell culture on the surface of a culture dish bathed in a nutrient medium supplemented with serum. If the cells are deprived of serum they stop dividing and become quiescent. Components called growth factors are the active principle present in serum necessary to stimulate entry into the cell cycle. Large numbers of different growth factors have now been isolated from a variety of different tissues. They are generally proteins active at low concentrations—often less than one microgram of growth factor per millilitre can substitute for the serum. For maximum stimulation of cell growth, however, it may be necessary to add several different growth factors to the culture medium. Two commonly investigated growth factors are platelet derived growth factor and epidermal growth factor.

Growth factors stimulate cells to grow by binding to specific growth factor receptors located on the cell surface membrane. Many receptors have been found to be tyrosine protein kinases, phosphorylating tyrosine residues of substrate proteins. The binding of a growth factor to its receptor is thought to activate the protein kinase activity, and this leads to phosphorylation of its protein substrates. How this stimulating signal is transduced to the rest of the cell is not well understood. In some cases signalling may entail breakdown of membrane inositol phospholipids generating second messengers inositol triphosphate and diacylglycerol. Coupling between this breakdown and the receptor may be brought about by guanosine triphosphate binding proteins regulating a phospholipase C enzyme that hydrolyses inositol phospholipid. Regardless of the precise mechanism of signal transduction the consequence of growth factor binding is to generate a wide range of changes in the cell. Within minutes there are changes in concentrations of low molecular weight components such as hydrogen and calcium ions and cyclic adenosine monophosphate; later there are changes in specific transcript and protein concentrations. These changes result in an overall increase in the rates of cellular protein and ribonucleic acid synthesis, leading to cell growth and eventual entry into the cell cycle.

Cell spreading and anchorage dependence

Cell spreading and attachment to a surface also influences whether cells enter the cell cycle. Normal cells not attached to a surface round up and become quiescent. On attachment to a surface they spread out and enter the cell cycle. This requirement to be attached to a surface is called anchorage dependence. Study of cells grown on special islands in a culture dish that restrain the extent of cellular spreading has shown that there is a correlation between the amount of spreading and the frequency with which a cell divides. When the island is large cells continue dividing until they fill the island and form a confluent monolayer. Because the cells then have less room they round up and become quiescent. Wounding this monolayer by scraping away a strip of cells allows the cells at the edge of the wound to spread out and to re-enter the cell cycle.

Part of the explanation for these phenomena is likely to be that the growth factor receptors on the surface of spread out cells are more accessible to growth factors in the culture medium. Also there appears to be an absolute requirement for cells to have some attachment to a surface. This may be necessary to generate changes in the cell cytoskeleton, which are required before cells can enter the cell cycle.

Cell transformation and oncogenes

Some cell lines do not exhibit anchorage dependence and are not so dependent on the presence of growth factors in the culture medium. When grown on a culture dish surface they form layers of cells on top of each other instead of forming a confluent monolayer.

Such cell lines are called transformed; they can be generated in a variety of ways such as by infection with certain animal viruses or after prolonged growth in culture. After injection back into an animal transformed cells often form tumours, suggesting that they are defective in the normal controls regulating entry into the cell cycle.

A class of genes called oncogenes have been found to be responsible for cell transformation. They have been isolated from transforming viruses and from DNA extracted from tumour cells. Homologues to oncogenes have been found in normal untransformed cells and are called proto-oncogenes. Cloning and sequencing of these genes have established that some of them encode previously identified components of the growth factor signal transducing pathway—for example, the proto-oncogene *c-sis* encodes a subunit of platelet derived growth factor, *c-erbB* a receptor for epidermal growth factor, and *c-ras* a guanosine triphosphate binding protein. Mutants of these proto-oncogenes are thought to generate oncogenes by altering the signal transducing pathway and stimulating entry into the cell cycle without the usual requirements for growth factors and anchorage dependence.

Relevance for the organism

These studies of cells in culture have provided a picture of how cell reproduction may be controlled in the intact organism. Cells remain quiescent unless they are exposed to particular growth factors and have the appropriate receptors to respond. Finer control is possible if several growth factors have to act together to elicit maximal response. Contacts between cells and surfaces can modulate the stimulation and as a consequence may be important for maintaining tissue cohesiveness and for preventing cells reproducing when detached from their current position. In these ways cells are stimulated to enter the cell cycle at the appropriate time and place necessary for the correct development and maintenance of the organism. Failure of these controls leads to cell transformation and possibly to the formation of tumours.

Commitment to the cell cycle

Once a cell has been stimulated to enter the cell cycle it remains subject to various controls. A major point of regulation is located towards the end of G1, where commitment to the cell cycle takes place. In mammalian cells this is often known as the restriction point. Cells deprived of growth factors before this point will leave the cell cycle and enter the GO quiescent state. Once past this point deprivation of growth factors often has little effect on the cell cycle in progress. Only after entering the G1 of the next cell cycle can the cell become quiescent. Before the restriction point cells are also more sensitive to inhibition of protein synthesis. If protein synthesis is inhibited to about half the normal level in such cells they cannot proceed through to S phase and mitosis; once past the restriction point, however, inhibition to these levels has no effect on the ability of cells to complete that cell cycle.

A similar commitment control called start acts in the G1 phase of the unicellular eukaryotic yeasts. Several genes required for yeast cells to complete start have been identified. One of these, *cdc2/28*, has been shown to encode a threonine or serine protein kinase, indicating that protein phosphorylation plays an important part in committing yeasts to the cell cycle. Once past start the cells begin the programme of events which lead eventually to S phase and mitosis. For example, after start there is synthesis of specific components which will be required for DNA replication during S phase.

Transition from G2 to mitosis

A further point of control occurs at the initiation of mitosis. Fusion of mammalian cells at different stages of the cell cycle has revealed the existence of factors which can induce cells into mitosis.

These factors can be assayed directly by injecting cell extracts into frog oocytes arrested at meiotic nuclear division. If the extracts contain these factors then the oocyte is induced into meiosis. Such experiments have shown that the inducing factors are present in mitotic mammalian cells and vary in concentration during cleavage of the frog embryo, the concentration reaching a peak as the cell enters mitosis.

Genes which are important for initiating mitosis have also been identified in yeast. Interestingly, the same gene *cdc2/28* implicated in cell cycle commitment also plays a part in the initiation of mitosis. The *cdc2/28* protein kinase presumably phosphorylates certain key proteins, which leads to progress through the cell cycle. In G1 at start these activate processes leading to S phase, and in late G2 they activate processes leading to mitosis.

Similarities between yeast and mammals

Recently, human cells have been found to contain a *cdc2/28* gene homologue. The gene can substitute for the yeast gene and encodes a structurally closely related protein kinase. The conservation of a cell cycle control gene in yeasts and humans indicates that aspects of the control will probably be similar in all eukaryotic organisms. It also suggests that protein phosphorylation has a key role in mammalian cells, both in commitment to the cell cycle and in the initiation of mitosis. Rapid advances may now be expected in this area as yeast is convenient for such studies and may be used as a surrogate organism for investigating the mode of action and regulation of the human *cdc2/28* homologue.

Further reading

Several excellent chapters relevant to cell reproduction may be found in Bradshaw RA, Prentis S, eds. *Oncogenes and Growth Factors*. Amsterdam: Elsevier Science Publications, 1987: Bishop JM. Trends in oncogenes, 1-10; Weinberg RA. Cellular oncogenes, 11-6; Burgess A. Growth factors and oncogenes, 123-34; Hunter T. Oncogenes and growth control, 135-42; Majerus PW, Wilson DB, Connolly TM, Bross TE, Neufeld EJ. Phosphoinositide turnover provides a link in stimulus-response coupling, 269-74; Nurse P. Cell cycle control genes in yeast, 275-83. O'Neill CH, Jordan P, Ireland G. Evidence for two distinct mechanisms of anchorage stimulation in freshly explanted and 3T3 Swiss mouse fibroblasts. *Cell* 1986;44:489-96. Lee MG, Nurse P. Complementation used to clone a human homologue of the fission yeast cell cycle control gene *cdc2*. *Nature* 1987;327:31-5. Kirschner M, Newport J, Gerhart J. The timing of early developmental trends in *Xenopus*. *Trends in Genetics* 1985;1:41-7.

MATERIA NON MEDICA

Taking silk

England, December 1954. It was the season of peace and good will to all. Father Christmas's much awaited visit was drawing near.

On the morning of the appointed day I set out on my good will mission to the hospital wards. In one ward, after the exchange of customary greetings, I was detained—that is how I would put it—by the sister in charge, for a function that was non-medical and concerned the dining table. I was led by her to the middle of the ward where a food trolley was set. Someone in the meantime had tied an apron on me and put a chef's cap on my head. I was then asked to dish out Christmas dinner to the patients. Of course I had assistants. I carved the turkey with utmost care, so there were no complaints, and everybody got a fair deal. But I don't know if I succeeded in my effort (probably I made a mess of it). Everybody watched my action with interest and amusement. My consternation contributed, in a positive way, to the festive mood of a festive season. When at last I finished my assigned task I was promptly derobed—I mean the apron and cap were removed—and I was again led by the sister, this time to the Christmas tree in the ward. I found Father Christmas had not forgotten me while preparing for his annual visit. A small packet was brought down from the tree and was handed to me. While opening it I wondered what Santa Claus had brought for me from the Arctic—it was a silk tie, I found.

India, Independence Day 1987. The tie, though worn, is still in my wardrobe.—S K MAJUMDAR, Jodhpur, India.