

omeprazole has not been shown to cause any changes in endocrine cell density,<sup>16</sup> but further studies are needed to elucidate the possible consequences of prolonged acid inhibition in man.

We thank Mrs Rigmor Petersen and Mrs Grete Eskesen for secretarial help; the staff of our endoscopy units for their valued support; Mr Kjeld Clemmensen-Rotne, Astra-Gruppen A/S, Albertslund, Denmark, for skilfully monitoring the conduct of the study; and A B Hässle, Mölndal, Sweden, for the study drugs.

## References

- Gledhill T, Hunt RH. Oesophagitis and hiatus hernia. In: Misiewicz JJ, Pounder RE, Venables CW, eds. *Diseases of the gut and pancreas*. Oxford: Blackwell, 1987:137-51.
- Johansson K-E, Boeryd B, Johansson K, Tibbling L. Double-blind crossover study of ranitidine and placebo in gastro-oesophageal reflux disease. *Scand J Gastroenterol* 1986;21:769-78.
- Wesdorp E, Bartelsman J, Pape K, Dekker W, Tytgat GN. Oral cimetidine in reflux esophagitis: a double blind controlled trial. *Gastroenterology* 1978;74:821-4.
- Wesdorp ICE, Dekker W, Klinkenberg-Knol EC. Treatment of reflux oesophagitis with ranitidine. *Gut* 1983;24:921-4.

- Koelz HR, Birchler R, Bretholz A, et al. Healing and relapse of reflux esophagitis during treatment with ranitidine. *Gastroenterology* 1986;91:1198-205.
- Clissold SP, Campoli-Richards DM. Omeprazole. A preliminary review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in peptic ulcer disease and Zollinger-Ellison syndrome. *Drugs* 1986;32:15-47.
- Walt RP, Gomes M de FA, Word EC, Logan LH, Pounder RE. Effect of daily oral omeprazole on 24 hour intragastric acidity. *Br Med J* 1983;287:12-4.
- Lauritsen K, Rune SJ, Bytzer P, et al. Effect of omeprazole and cimetidine on duodenal ulcer: a double-blind comparative trial. *N Engl J Med* 1985;312:958-61.
- Lauritsen K, Rune SJ, Wulff HR, et al. Effect of omeprazole and cimetidine on prepyloric gastric ulcer: double blind comparative trial. *Gut* (in press).
- Bardhan KD, Bianchi Porro G, Bose K, et al. A comparison of two different doses of omeprazole versus ranitidine in treatment of duodenal ulcers. *J Clin Gastroenterol* 1986;8:408-13.
- Schaub N, Meyrick Thomas J, Misiewicz JJ, Lowell D, Trotman IF. Investigation of ranitidine 150 mg bd or 300 mg bd in the treatment of reflux disease. *Hepatogastroenterology* 1986;33:208-13.
- Fleiss JL. *Statistical methods for rates and proportions*. New York: Wiley, 1981.
- Klinkenberg-Knol EC, Jansen JMBJ, Festen HPM, Meuwissen SGM, Lamers CBHW. Double-blind multicentre comparison of omeprazole and ranitidine in the treatment of reflux oesophagitis. *Lancet* 1987;i:349-51.
- Sonnenberg A, Lepsien G, Müller-Lissner SA, Koelz HR, Siewert JR, Blum AL. When is esophagitis healed? Esophageal endoscopy, histology and function before and after cimetidine treatment. *Dig Dis Sci* 1982;27:297-301.
- Langman MJS. Antisecretory drugs and gastric cancer. *Br Med J* 1985;290:1850-2.
- Karvonen A-L, Keyrilainen O, Uusitalo A, et al. Effects of omeprazole in duodenal ulcer patients. *Scand J Gastroenterol* 1986;21:449-54.

(Accepted 27 October 1987)

# Upper gastrointestinal Kaposi's sarcoma in patients positive for HIV antibody without cutaneous disease

I G BARRISON, S FOSTER, J W HARRIS, A J PINCHING, J G WALKER

## Abstract

Six patients with antibodies to the human immunodeficiency virus (HIV) and with persistent gastrointestinal symptoms of HIV infection but without cutaneous lesions of Kaposi's sarcoma underwent endoscopy. Four also underwent barium meal examination. In all six cases small lesions were seen in the stomach at endoscopy, and histological examination of biopsy specimens taken from the lesions confirmed the diagnosis of Kaposi's sarcoma. The barium meal examinations were reported as normal in three patients and showed oesophageal candidiasis in the fourth.

These findings suggest that Kaposi's sarcoma of the upper gastrointestinal tract is common in patients positive for HIV antibody, even those without cutaneous lesions. Endoscopy, with biopsy of suspicious lesions, is necessary to make the diagnosis and is recommended in all HIV antibody positive patients with persistent upper gastrointestinal symptoms.

## Introduction

Infection with human immunodeficiency virus (HIV) results in a wide range of opportunistic infections and tumours, the commonest of which is Kaposi's sarcoma. Two studies have suggested that

gastrointestinal Kaposi's sarcoma is more likely to be present in this group of patients if there is extensive cutaneous disease.<sup>1,2</sup> We describe six patients positive for HIV antibody without cutaneous disease in whom upper gastrointestinal Kaposi's sarcoma was diagnosed after endoscopy for the investigation of persistent symptoms.

## Case reports and methods

In all six patients, all of them men, the presence of HIV infection was diagnosed by a positive competitive enzyme linked immunosorbent assay (ELISA) for the detection of anti-HIV antibodies (Wellcome) and confirmed by a competitive ELISA system (Abbott). Their associated conditions and reasons for performing endoscopy are shown in the table. Endoscopies were performed by IGB using an Olympus GIF-T<sub>10</sub> fully immersible endoscope. Two to four biopsy specimens were obtained from each lesion using the 5 mm Olympus biopsy forceps. Cutaneous Kaposi's sarcoma was excluded by at least two clinical examinations by independent observers.

Findings on endoscopy and the results of a barium meal examination (performed in four patients) are shown in the table, and typical endoscopic appearances are shown in the figure. Histological confirmation of the diagnosis of Kaposi's sarcoma was obtained in all six cases. Five of the six patients had oral candidiasis but only three had oesophageal candidiasis. The barium meal examinations were reported as normal in three patients and showed oesophageal candidiasis in the fourth.

## Discussion

These six cases show that upper gastrointestinal Kaposi's sarcoma frequently occurs in HIV antibody positive patients with persistent symptoms in the absence of cutaneous disease. At present the prevalence of gastric Kaposi's sarcoma among HIV antibody positive patients without cutaneous lesions is unknown. We have found lesions in six out of 25 symptomatic patients (26%) examined to date and in seven out of 13 patients with cutaneous Kaposi's

Departments of Gastroenterology, Clinical Immunology, and Genitourinary Medicine, St Mary's Hospital and Medical School, London W2 1NY

I G BARRISON, BSC, MRCP, senior medical registrar


S FOSTER, MRCP, senior registrar in genitourinary medicine

J W HARRIS, FRCP, consultant physician in genitourinary medicine

A J PINCHING, DPHIL, FRCP, senior lecturer in clinical immunology

J G WALKER, MD, FRCP, consultant physician and gastroenterologist

Correspondence to: Dr Barrison.



Endoscopic appearances (marked by arrows) in three patients with gastrointestinal Kaposi's sarcoma. Left (case 1): macular lesions in body of stomach; middle (case 4): polypoid lesion on greater curve of stomach; right (case 5): plaque-like lesion on lesser curve of stomach.

Clinical details of HIV antibody positive patients with gastric Kaposi's sarcoma

Case No	Age and sex	Barium meal	Other HIV associated disease	Indications for endoscopy	Findings on endoscopy
1	35 M		Chronic hepatitis B carrier, oral candidiasis, perianal herpes simplex	Postprandial epigastric pain	Two small plaques (<5 mm) in body of stomach
2	39 M	Normal	Oral candidiasis, persistent generalised lymphadenopathy	Intractable dysphagia	Oesophageal candidiasis, multiple small plaques throughout stomach
3	29 M	Oesophageal candidiasis	Oral candidiasis	Intractable dysphagia	Oesophageal candidiasis, several small (<5 mm) red plaques in gastric antrum
4	37 M		Oral candidiasis, persistent generalised lymphadenopathy	Chronic nausea and anorexia	Oesophageal candidiasis, polypoid lesions throughout body and antrum of stomach
5	35 M	Normal	Angiofollicular hyperplasia	Weight loss, persistent vomiting	Multiple small red plaques in stomach and duodenum
6	43 M	Normal	Oral candidiasis, perianal herpes simplex	Postprandial epigastric discomfort	Multiple small red plaques in oesophagus, stomach, and duodenum

sarcoma (54%). Friedman *et al* reported a prevalence of 40% in 50 patients with the acquired immune deficiency syndrome (AIDS) and cutaneous Kaposi's sarcoma<sup>2</sup> and Saltz *et al* a prevalence of 50% in 19 similar patients.<sup>1</sup>

In two of our four patients the lesions were less than 5 mm in diameter and could have been misinterpreted as patchy gastritis at endoscopy. Biopsy of all suspicious lesions is necessary in this group of patients. Barium examination of the stomach was reported as normal in three patients with gastric Kaposi's sarcoma, and, although contrast studies are helpful when the classical submucosal filling defects are present, the technique is not very sensitive in detecting macular lesions such as those seen in five of our six patients. There is still some concern about the safety of fiberoptic endoscopy in patients with HIV antibodies—which is unjustified—and some doubts about the diagnostic yield. One study in patients with upper gastrointestinal symptoms and oropharyngeal candidiasis suggested that endoscopy would merely confirm the presence of oesophageal candidiasis.<sup>3</sup> Five of our six patients had oral candidiasis and the diagnosis of Kaposi's sarcoma would not have been made without endoscopy.

Newer, more efficacious treatments for HIV related diseases are becoming available, and prompt diagnosis of gastrointestinal

Kaposi's sarcoma is clearly desirable from a prognostic and a therapeutic point of view. Symptoms may resolve after chemotherapy and the progression of the Kaposi sarcoma may be delayed. Specific anti-HIV treatments are now available—for instance, the reverse transcriptase inhibitor zidovudine (azidothymidine)—which may influence the clinical course of the disease and therefore increase the importance of making an early and accurate diagnosis.<sup>4</sup> We therefore recommend endoscopy in all patients with HIV antibodies who have persistent upper gastrointestinal symptoms.

References

1 Saltz RK, Kurtz RC, Lightdale CJ, *et al*. Kaposi's sarcoma: gastrointestinal involvement and correlation with skin findings and immunologic function. *Dig Dis Sci* 1984;29:817-23.

2 Friedman SL, Wright TL, Altman DF. Gastrointestinal Kaposi's sarcoma in patients with acquired immunodeficiency syndrome. *Gastroenterology* 1985;89:102-8.

3 Tavittian A, Raufman JP, Rosenthal LE. Oral candidiasis as a marker for oesophageal candidiasis in the acquired immunodeficiency syndrome. *Ann Intern Med* 1986;104:54-5.

4 Fischl MA, Richman D, Grieco MH, *et al*. The efficacy of azidothymidine (AZT) in the treatment of patients with AIDS and AIDS-related complex. *N Engl J Med* 1987;317:185-91.

(Accepted 26 October 1987)