the appropriateness of referrals—that is, the extent to which general practitioners', consultants', and patients' expectations are met.¹⁷ More studies are needed on qualitative aspects of referrals so that rates of referral may be interpreted in relation both to the benefits of referral—for example, improved health and well being—and to the risks of referral—for example, unnecessary investigation or treatment.

Many of the problems in measuring and interpreting rates of referral have yet to be solved. General practitioners should welcome meaningful data on their referrals as a means of improving patient care. The benefits that will result from discussion of referrals are likely to be educational ones and will arise from discussions between small groups of general practitioners and between general practitioners and district consultants to whom they refer. The DHSS intends to use information on rates of referral "to ensure quality of care in a cost effective way." Whether the accent will be on quality or cost remains to be seen.

MARTIN ROLAND

Director of Studies in General Practice,
Clinical School,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

- 1 Royal College of General Practitioners, Office of Population Censuses and Surveys, Department of Health and Social Security. *Morbidity statistics from general practice 1971-2. Second national survey.* London: HMSO, 1979. (Studies on medical and population subjects, No 36.)
- 2 Cummins RO, Jarman B, White PM. Do general practitioners have different "referral thresholds?" *Br Med J* 1981;282:1037-9.
- 3 Gillam DM. Referral to consultants—the National Health Service versus private practice. *J R Coll Gen Pract* 1985;35:15-8.
- 4 Wilkin D, Smith A. Variation in general practitioners' referrals to consultants. *J R Coll Gen Pract* 1987;37:350-3.
- 5 Secretaries of State for Social Services, Wales, Northern Ireland, and Scotland. *Promoting better health. The government's programme for improving primary health care.* London: HMSO, 1987. (Cmd 249.)
- 6 Royal College of General Practitioners, Office of Population Censuses and Surveys, Department of Health and Social Security. *Morbidity statistics from general practice 1981-82. Third national survey.* London: HMSO, 1986. (Series MB5.)
- 7 McBride M, Bligh J, Bradley P, Clarke G, Fleming D. *A descriptive survey of morbidity encountered in general practice in the Mersey Region.* Liverpool: Royal College of General Practitioners, Mersey Faculty, 1981.
- 8 Scott R, Gilmore M. The Edinburgh hospitals. In: McLachlan G, ed. *Problems and progress in medical care.* Oxford: Oxford University Press, 1966:5-41.
- 9 Wright HJ. *General practice in south west England.* London: Royal College of General Practitioners, 1968. (Report from general practice, No 7.)
- 10 Forsyth G, Logan RFL. *Gateway or dividing line? A study of hospital outpatients in the 1960s.* Oxford: Oxford University Press, 1968.
- 11 Robertson RC. Variations in referral pattern to the psychiatric services by general practitioners. *Psychol Med* 1979;9:355-64.
- 12 Dowie R. *General practitioners and consultants—a study of outpatient referrals.* London: King Edward's Hospital Fund for London, 1983.
- 13 Wilkin D, Smith A. Explaining variation in general practitioner referrals to hospital. *Fam Pract* 1987;4:160-9.
- 14 Morrell DC, Gage HG, Robinson A. Referral to hospital by general practitioners. *J R Coll Gen Pract* 1971;21:77-85.
- 15 Ross AK, Davis WA, Horn G, Williams R. General practitioner orthopaedic outpatient referrals in north Staffordshire. *Br Med J* 1983;287:1439-41.
- 16 Evans EO, McBride K. Hospital usage by a group practice. *J R Coll Gen Pract* 1968;16:294-306.
- 17 Grace JF, Armstrong D. Referral to hospital: perceptions of patients, general practitioners and consultants about necessity and suitability of referral. *Fam Pract* 1987;4:170-5.

Consensus on overtreating cancer

Too many patients overtreated

No doctor likes to have patients die, and for many it seems natural to keep on trying to achieve a cure until all hope is lost. But attitudes to death have changed in the past 20 years, and doctors can no longer justify such therapeutic persistence by arguing that the patient would feel abandoned if the search for a cure was halted. Heroic, massive surgery has already gone out of fashion, but too many patients with cancer still suffer from excessive treatment with radiotherapy or drugs, or both. Sadly, the public's image of treatment of cancer is heavily weighted by the minority of patients suffering from loss of hair, repeated vomiting, and other grim toxic effects of chemotherapy—and then going on to die. Giving a regimen of treatment designed to achieve cures in circumstances when the intent should only be palliation is bad medicine. The patient suffers unpleasant side effects, and the high cost of the treatment usually has to come from a health authority budget that may already be in the red.

These were the conclusions at the halfway point of a recent meeting of cancer specialists, health economists, and others concerned with cost-benefit analysis in oncology. Example after example of inappropriate management was quoted: investigations being done long after the cancer was disseminated and combinations of anticancer drugs being given to patients with tumours known *not* to respond to chemotherapy. And, the participants agreed, when curative chemo-

therapy was indicated the drugs chosen were often new, not fully evaluated, and much more expensive than the well tried alternatives. At least subconsciously the doctor often seems to be hoping that the latest drug will be superior to its analogues.

On p 471 the working group publishes its conclusions. Firstly, it argues that doctors should no longer claim an unfettered right to treat their patients with whatever treatment they might think best. This clinical freedom merges imperceptibly into clinical irresponsibility; doctors' exercise of autocratic rights may infringe both the patient's right to treatment that is of established worth and minimal toxicity and their colleagues' rights to their fair share of NHS resources.

Such a judgment implies that there is a consensus on managing patients with different types of cancer at various stages. In reality there are few types of cancer for which this is true, but a working consensus could probably be agreed for most others. The declaration calls for oncologists, radiotherapists, and others to come together and establish review panels to work out guidelines. Doctors competing for scarce resources seem more likely to get the money they need if they can show a rational, peer reviewed basis for their treatments. And their patients should benefit too.

TONY SMITH

Deputy Editor, *BMJ*