Writing a wrong

Kenneth Mullan

In 1983 the Court of Appeal, in the case of *Dwyer v Roderick and others*,¹ emphasised the serious consequences of negligently writing and dispensing medical prescriptions. Those consequences were re-emphasised in the recent High Court decision in the case of *Prendergast v Sam and Dee Ltd*.²

Wrong dosage

Dr Ian Roderick wrote a prescription for Mrs Joan Dwyer, who had complained to him of severe headaches, for a pain killing drug which was proved successful in the treatment of migraine. The drug, ergotamine tartrate (Migril), is extremely dangerous if not taken in proper doses. It can produce gangrene. Dr Roderick did not prescribe the drug in the proper doses. Mrs Dwyer took the prescription to the pharmacy of Cross Chemists (Banbury) Ltd. There she was given ergotamine tartrate in a container displaying the exact dosage as recommended by Dr Roderick.

Mrs Dwyer began to take the drug as directed and rapidly became very ill. During this time she was seen by a partner of Dr Roderick, Dr Jackson, who called to see Mrs Dwyer from his own home and therefore did not have her medical notes with him. He gave evidence that he was unaware that Mrs Dwyer was taking ergotamine tartrate. He stated that he had examined drugs that were on her bedside table but had not seen ergotamine tartrate. By the time that the mistake was discovered Mrs Dwyer was suffering from gangrene and her toes had to be amputated. As a result she became permanently crippled.

In the High Court Mr Justice Stuart-Smith noted that negligence was admitted by Dr Roderick who had written the prescription and by the pharmacy which had dispensed it. There were therefore two main issues for the judge to decide. Firstly, he had to consider whether any further liability lay with Dr Jackson. In an attempt to limit its liability the pharmacy had joined Dr Jackson as another defendant. The judge held that the overwhelming likelihood was that on Dr Jackson's first visit to Mrs Dwyer a bottle containing ergotamine tartrate was by her bedside. Dr Jackson had persuaded himself, during the eight years which it took for the case to come to trial, that he could not have known that the plaintiff was taking the drug.

Having concluded that Dr Jackson had also been negligent, the judge had to decide what the proper apportionment of liability should be. Accordingly, Mr Justice Stuart-Smith awarded damages of £100 000 against Dr Roderick, Dr Jackson, and the pharmacy to be apportioned as to 45% to Dr Roderick, 15% to Dr Jackson, and 40% to the pharmacy.

On appeal the Court of Appeal reversed the decision of the High Court in relation to the liability of Dr Jackson. Lord Justice May took the view that after the passage of eight years it was inevitable that however truthful a witness might be trying to be, at least part of his evidence would be inaccurate. By agreement, the pharmacy accepted liability for the 15% liability which had rested on Dr Jackson.

In the end, therefore, Dr Roderick's initial gross negligence only cost him 45% of the blame with the pharmacy accepting the remaining 55%.³

Dr Stuart Miller wrote a prescription for Mr Prendergast, who was asthmatic with a chest infection, prescribing three Ventolin (salbutamol) inhalers, 250 Phyllocontin (aminophylline) tablets, and 21 Amoxil (amoxycillin) tablets. It was accepted that this would

be a commonplace combination of drugs for a patient with asthma and a chest infection.

Mr Prendergast took the prescription to the pharmacy of Sam and Dee Ltd, where it was dispensed by a pharmacist, Mr Peter Kozary. Mr Kozary dispensed the inhalers and Phyllocontin correctly, but instead of the Amoxil he had dispensed Daonil (glibenclamide), a drug used for diabetes to reduce the sugar content in the body. Mr Prendergast was not a diabetic and as a result of taking a large dosage of the Daonil suffered permanent brain damage and symptoms of hypoglycaemia.

In his judgment Mr Justice Auld dealt firstly with the position of the pharmacist, Mr Kozary. Mr Kozary had argued in his defence that the word Amoxil on the prescription was unclear and was capable of being read as Daonil. The "A" could be mistaken for a lower case "d" and the "x" for "n." His lordship, therefore, found it necessary to consider Dr Miller's handwriting. While the question would always be one of general impression, in his view the word Amoxil on the prescription was capable of being read as Daonil.

Other indications

Assuming, however, that the writing was not clear, the judge was of the opinion that there were sufficient other indications to put Mr Kozary on inquiry that something was wrong. More particularly, it was known that Daonil was made only in 5 mg strengths, while the word Amoxil was followed by "250." Mr Kozary's defence that he thought that Dr Miller had mixed Daonil up with another diabetic drug which was taken in 250 mg was confirmation of the need for the prescription to be checked with the doctor.

Secondly, the dosage of 250 mg was normal for Amoxil, but high, and unusually high for Daonil. This fact combined with the assumed knowledge that the taking of Daonil was dangerous for non-diabetics should have put Mr Kozary on his guard. Finally, Mr Kozary should have noticed that Mr Prendergast had paid for the drugs on collection when it might have been expected that a diabetic would have been entitled to free drugs.

Accordingly, the judge concluded that had Mr Kozary been paying attention when he dispensed the drugs he should have known that there was something amiss with the contents of the prescription and should have checked these with the doctor. He had not been paying attention and therefore fell below the standard of care and skill expected of a professional pharmacist and had been negligent. There had been sufficient information on the prescription as a whole and in his dealings with the patient to put him on inquiry.

Mr Justice Auld then turned to the position of Dr Miller. He first made it clear that if Dr Miller owed a legal duty of care to his patient and had been in breach of that duty it would be no defence to his liability to rely on the already established negligence of Mr Kozary.

The judge indicated forcefully that a doctor did owe a duty of care to his patient to write a prescription clearly and of sufficient legibility to allow for possible mistakes by a busy pharmacist who might be distracted by other customers. Having already established that in his opinion the word Amoxil on the prescription could have been read as Daonil, Dr Miller had been in breach of his duty of care to write clearly and had been negligent. Such liability could not be excused by the argument that there had been sufficient information on

the prescription to put Mr Kozary on his guard. Dr Miller's negligence had contributed to the negligence of Mr Kozary, although the greater proportion of the responsibility lay with Mr Kozary.

Accordingly, Mr Justice Auld awarded damages of £119 302 plus interest against Mr Kozary, his pharmacy, and Dr Miller, the proper apportionment of which was that Dr Miller was 25% liable and Mr Kozary 75% liable.

The implications

Though the implications of the decision in these cases may seem obvious from the facts, their seriousness should not be underestimated by those in the medical and pharmaceutical professions. What is clear from the decision in *Prendergast* is that doctors are under a legal duty of care to write clearly—that is, with sufficient legibility to allow for mistakes by others. Both *Prendergast* and *Dwyer* show that liability need not stop when the prescription leaves the hands of the doctor, even when the doctor has been grossly negligent. It may extend into and be a cause of the negligent mistakes of others.

The pharmacist must be aware that he or she must not tacitly accept what they see, or perceive to see, on the written prescription before them. They are under a legal duty of care to draw on their skill and knowledge of drugs to inquire into the surrounding circumstances of the case. In this respect the finding of his lordship in *Prendergast* that the pharmacist should have noticed that the patient paid for the drugs is noteworthy. If there is any doubt in the pharmacist's mind then the prescription should be checked by the prescribing doctor.

When thousands of prescriptions are being written and dispensed every day it is inevitable that mistakes will creep in. The judgments in *Dwyer v Roderick* and others and in *Prendergast v Sam and Dee Ltd* have shown that the courts will take a less than tolerant attitude towards those who make careless but negligent mistakes which result in personal injury and will award sufficient damages accordingly.

1 [1983] 80 *Law Society Gazette* 3003.

2 *The Times* 1988; Mar 14:19.

3 As to the morality of such a finding, see Finch J. A costly oversight. *New Law Journal* 1982; Feb 25:176-7.

Cost versus benefit in non-surgical management of patients with cancer

A working group of cancer physicians, epidemiologists, and health economists met recently to consider the need for medical audit of the treatment of patients with cancer. The meeting was set up with the generous support of the Cancer Research Campaign because its participants believed that such an audit has become urgent, with the number of cancer patients rising (owing to the increase in the proportion of elderly people in the population) and with the resources available being limited. The first step should be the development of guidelines for optimal patient care and use of these resources. The group agreed that two key questions had to be answered:

- (1) Is unfettered clinical freedom appropriate in the management of patients with cancer?
- (2) Is consensus possible on what is appropriate and inappropriate care for patients with cancer?

On the first issue the group believes that unrestrained clinical freedom is inappropriate for management of patients with cancer as it is for other diseases. It results in wide variations in management without corresponding differences in outcome as measured by benefits to patients.¹ In the context of cancer care such variations are sometimes harmful to the patient and may also waste resources. In particular, while some investigations clearly influence patient management, other indiscriminate investigations usually contribute solely to the acquisition of data and are both irrelevant and wasteful. Examples include the routine use of skeletal surveys, radionuclide scans, and ultrasound in patients with early asymptomatic breast cancer or the search for disease at other sites in patients with documented metastases.²

Several controlled studies have shown that no treatment, less treatment, or shorter or cheaper treatment may sometimes be as good as, or better than, a more expensive option. Examples include one versus ten fractions of palliative radiotherapy for the relief of metastatic bone pain³ and a randomised trial of cytotoxic chemotherapy versus symptomatic care only for patients with advanced non-small cell lung cancer (C J Williams *et al*, paper to 4th European Conference

on Cancer and Cancer Nursing, 1987). Similarly it is not easy to justify (other than within the context of clinical research studies) the use of new and expensive cytotoxic or hormonal drugs where clear advantage over older and cheaper agents has not been established in controlled trials.

Whenever possible the optimum management for individual patients should be based on the outcome of carefully planned and well controlled clinical trials, but when such research studies are in apparent conflict or there is controversy about treatment decisions should be assisted by consensus-based guidelines.

Curative treatments may be costly, but they are often proportionally less expensive than palliative treatment when measured in terms of life years gained.⁴ When planning treatment clinicians need to distinguish clearly between those with curative and those with palliative intent; failure to do so may result in complex expensive treatments of unproved benefit being given to patients who would be better off with no treatment or a simpler, cheaper option (S E Vogl *et al*, paper to 13th International Congress of Chemotherapy, 1983). Failure to distinguish between palliative and curative intent may also lead to diversion of resources from areas where the same expenditure may procure substantially greater benefit for the same or other patients.

All cancer physicians must, we believe, consider not only the medical but also the social and financial consequences of their actions; if we do not recognise this responsibility then non-clinicians will take over decision making on the distribution of resources. For this reason the working group was concerned by the apparent lack of interest of many physicians in the issue of resource allocation and the apparent inability of clinicians to decide on a consensus for the management of common clinical conditions. The following recommendations were agreed and will form a basis for further action:

- (1) Guidelines defining appropriate and inappropriate care in cancer need to be set out by an expert review panel for each tumour site.

Adrian R Timothy,
coordinator,
Thurstan Brewin,
Jocelyn Chamberlain,
A Horwich, Bryan Jennett,
Paul Kind, Robert Maxwell,
James Raftery, Gareth Rees,
Peter Selby, Petr Skranabek,
M L Slevin, A J Smith,
Basil Stoll, Simon Sutcliffe,
James Till,
J M A Whitehouse

Correspondence to: Dr
Adrian R Timothy,
Department of
Radiotherapy and Oncology,
St Thomas's Hospital,
Lambeth Palace Road,
London SE1 7EH.