- sion in sexually transmitted human immunodeficiency virus infection. *Lancet* 1987;ii:589-93.
- 15 Fincher RME, De Silva M, Lobel S, *et al.* AIDS-related complex in a heterosexual man seven weeks after transfusion. *N Engl J Med* 1985;313:1226-7.
 - 16 Vittecoq D, Autran B, Bourstyn E, *et al.* Lymphadenopathy syndrome and seroconversion two months after single use of needle shared with an AIDS patient. *Lancet* 1986;ii:1280.
 - 17 Tucker J, Ludlam CA, Craig A, *et al.* HTLV-III infection associated with glandular-fever-like illness in a haemophilic. *Lancet* 1985;ii:585.
 - 18 Boiteux F, Vilmer E, Giro R, *et al.* Lymphadenopathy syndrome in two thalassemic patients after LAV contamination by blood transfusion. *N Engl J Med* 1985;312:648-9.
 - 19 L'age-Steir J, Schwartz A, Offermann G, *et al.* HTLV-III infection in kidney transplant recipients. *Lancet* 1985;ii:1361-2.
 - 20 White GC, Matthews TJ, Weinhold KJ, *et al.* HTLV-III seroconversion associated with heat-treated factor VIII concentrate. *Lancet* 1986;ii:611-2.
 - 21 Franceschi S, Tirelli U, Vaccher E, *et al.* Seroconversion to HTLV-III in haemophilic given heat-treated factor VIII concentrate. *Lancet* 1986;ii:803-4.
 - 22 Ruutu P, Suni J, Oksanen K, *et al.* Primary infection with HIV in a severely immunosuppressed patient with acute leukemia. *Scand J Infect Dis* 1987;19:369-72.
 - 23 Colebunders R, Greenberg AE, Francis H, *et al.* Acute HIV illness following blood transfusion in three African children. *AIDS* 1988;2:125-7.
 - 24 Carne CA, Tedder RS, Smith A, *et al.* Acute encephalopathy coincident with seroconversion for anti-HTLV-III. *Lancet* 1985;ii:1206-8.
 - 25 Ho DD, Sarngadharan MG, Resnick L, *et al.* Primary human T-lymphotropic virus type III infection. *Ann Intern Med* 1985;103:880-3.
 - 26 Piette AM, Tusseau F, Vignon D, *et al.* Acute neuropathy coincident with seroconversion for anti-LAV/HTLV-III. *Lancet* 1986;ii:852.
 - 27 Biggar RJ, Johnson BK, Musoke SS, *et al.* Severe illness associated with appearance of antibody to human immunodeficiency virus in an African. *Br Med J* 1986;293:1210-1.
 - 28 Hagberg L, Malmvall BE, Svennerholm L, *et al.* Guillain-Barré syndrome as an early manifestation of HIV central nervous system infection. *Scand J Infect Dis* 1986;18:591-2.
 - 29 Denning DW, Anderson J, Rudge P, *et al.* Acute myelopathy associated with primary infection with human immunodeficiency virus. *Br Med J* 1987;294:143.
 - 30 Wiselka MJ, Nicholson KG, Ward SC, *et al.* Acute infection with human immunodeficiency virus associated with facial nerve palsy and neuralgia. *J Infect* 1987;15:189-90.
 - 31 Abb J, Zachoval R, Zachoval V, *et al.* HIV antigen, HIV antibody and serum interferon in a patient with encephalopathy. *Infection* 1987;15:425-6.
 - 32 Calabrese LH, Proffitt MR, Levin KH, *et al.* Acute infection with the human immunodeficiency virus (HIV) associated with acute brachial neuritis and exanthematous rash. *Ann Intern Med* 1987;107:849-51.
 - 33 Gordon SM, Valentine FT, Holzman RS, *et al.* Acquired immune deficiency syndrome possibly related to transfusion in an adult without disease-risk factors. *J Infect Dis* 1984;149:1030-2.
 - 34 Feorino PM, Kalyanaraman VS, Haverkos HW, *et al.* Lymphadenopathy associated virus infection of a blood donor-recipient pair with acquired immunodeficiency syndrome. *Science* 1984;225:69-72.
 - 35 Goudsmit J. Persistent generalised lymphadenopathy and acyclovir. *Lancet* 1984;ii:1236.
 - 36 Gaines H, von Sydow M, Sönnernborg A, *et al.* Antibody response in primary human immunodeficiency virus infection. *Lancet* 1987;ii:1249-53.
 - 37 Gaines H, von Sydow M, Parry JV, *et al.* Detection of immunoglobulin M antibody in primary human immunodeficiency virus infection. *AIDS* 1988;2:11-5.
 - 38 von Sydow M, Gaines H, Sönnernborg A, *et al.* Antigen detection in primary HIV infection. *Br Med J* 1988;296:238-40.
 - 39 Albert J, Gaines H, Sönnernborg A, *et al.* Isolation of human immunodeficiency virus (HIV) from plasma during primary HIV infection. *J Med Virol* 1987;23:67-73.
 - 40 Greenblatt RM, Lukehart SA, Plummer FA, *et al.* Genital ulceration as a risk factor for human immunodeficiency virus infection. *AIDS* 1988;2:47-50.
 - 41 Kreiss J, Carael M, Mehus A. Role of sexually transmitted diseases in transmitting human immunodeficiency virus. *Genitourin Med* 1988;64:1-2.
 - 42 Kreiss JK, Koech D, Plummer FA, *et al.* AIDS virus infection in Nairobi prostitutes: spread of the epidemic to east Africa. *N Engl J Med* 1986;314:414-8.
 - 43 Tindall B, Barker S, Donovan B, *et al.* Characterization of the acute clinical illness associated with human immunodeficiency virus infection. *Arch Intern Med* 1988;148:945-9.
 - 44 Hoagland RJ. The clinical manifestations of infectious mononucleosis. A report of two hundred cases. *Am J Med Sci* 1960;240:55-63.
 - 45 Joncas J, Chiasson JP, Turcotte J, *et al.* Studies on infectious mononucleosis: III. Clinical data, serologic and epidemiologic findings. *Can Med Assoc J* 1968;98:848-54.
 - 46 Niederman JC, McCollum RW, Henle G, *et al.* Infectious mononucleosis. Clinical manifestations in relation to EB virus antibodies. *JAMA* 1968;203:139-43.

(Accepted 16 August 1988)

Pursuing efficiency in surgical practice

John H Wyllie, Iain G Kidson, David H Wyllie

Abstract

To examine fluctuations in numbers of patients on surgical wards the dates of admission and discharge of each of the 5556 patients admitted from 1 January 1985 to 31 December 1987 were examined during computerised audit of a single surgical firm. The numbers of patients under the care of the firm fluctuated widely, often exceeding the 38 beds nominally available. Duration of stay varied from two days or less (3062 admissions) to more than a month (163 admissions). One patient was in hospital for 278 days. The patients admitted for more than a month (2.9% of the total) filled 28% of the beds; not all these patients were elderly.

A further increase in throughput of patients undergoing elective operations might be achieved by always admitting patients on the day of operation, and perhaps by discharging patients even sooner than at present. Efficiency would increase but so would overall costs.

Introduction

An efficient surgical service should have a high throughput of patients and a low rate of cancellation. It is difficult to run: during a lull the beds and lists seem wastefully underused, and during a spate of emergencies elective operations have to be cancelled. These fluctuations have received scant attention. In 1984 we set up a computer system for audit on our general surgical firm, which has special interests in vascular and oesophageal surgery. We looked at records for 1985, 1986, and 1987 to see how the load on the wards fluctuated and to find where economies might be made

to preserve the service should financial stringency demand a cut in the number of beds.

Data collection

For all inpatients and for patients having operations on an outpatient or day case basis house surgeons recorded the following details on a microcomputer: the hospital number; surname; sex; age; consultant in charge; whether admission was emergency or elective; date of admission or transfer; date of discharge, transfer, or death; and up to four diagnoses, complications, and operations. These data were checked at the weekly meeting of the whole firm (two consultants, a senior registrar, a registrar, and three house surgeons) at which we review our work; diagnoses were modified as necessary each week until the patients were discharged. Programs to analyse the data were written.

This paper deals with patient episodes; any patient appears as many times as he or she was admitted during the study. Duration of stay was calculated as zero days if the patient was discharged on the date of admission, one day if discharge was on the day after admission, and so on. As this does not allow estimation of the number of beds required for day cases we assumed that for a stay of zero days the requirement was 0.5 bed day.

Results and discussion

NUMBERS OF ADMISSIONS AND BEDS USED

From 1 January 1985 to 31 December 1987 there were 5556 admissions. Table I shows that the number of emergencies rose by 18.8% and of elective opera-

Academic Unit of Surgery,
Whittington Hospital,
London N19 5NF
John H Wyllie, FRCS,
honorary consultant surgeon
Iain G Kidson, FRCS,
consultant surgeon

University College and
Middlesex School of
Medicine, University
College London, London
WC1E 6BD
David H Wyllie, medical
student

Correspondence to:
Professor J H Wyllie.

TABLE 1—Numbers of admissions, mean length of stay, and use of beds

	1985	1986	1987	Total
No of admissions*	1698	1882	1976	5556
Emergencies	743	710	883	2336
Elective cases	955	1172	1093	3220
Mean stay (days):				
Emergencies	7.2	6.3	7.4	7.0
Elective cases	4.2	5.6	4.3	4.8
No of bed days	9379	11 106	11 278	31 763
Emergencies	5375	4502	6573	16 450
Elective cases	4004	6604	4705	15 313

*Includes outpatient operations.

tions by 14.5% over this period. Patients admitted for emergency operations accounted for 42% of those admitted, stayed longer in hospital than patients undergoing elective operations, and accounted for 52% of the bed use. The average number of beds in use each day (34.5) was marginally below the nominal allocation of 38, but this average conceals fluctuations. These fluctuations determine when crises occur.

Bed use day by day for three years was reconstructed from dates of admission and discharge. Figure 1 shows numbers of beds used each day for one year starting 1 August 1986. The beds required for elective admissions showed a weekly periodicity not shown by the requirements for emergency cases. This reflects the fact that our firm has routine lists on the same days each week but takes emergencies on different days in a six week cycle. The number of beds used fluctuated considerably, with peaks 20 or even 30 beds above the troughs. Over the Christmas period elective operations stopped and only five patients admitted as emergencies and 10 other patients were left in our wards.

It is often suggested that elective admissions should be planned so that capacity for emergencies is avail-

able, and this is the policy of the surgical division in our hospital. If this policy had been strictly applied we could not have approached our actual workload. Emergency admissions in a single day varied from none to 14, or exceptionally 28 (after a riot in Tottenham). Figure 2 shows that 38 beds would have been adequate for our actual practice only 68% of the time, 49 beds 95% of the time, and 51 beds 99% of the time. Loss of beds would obviously have a deleterious effect on service.

FIG 2—Proportion of days on which a given allocation of beds would have been inadequate. Nominal allocation was 38 beds

Flexible use of a pool of vacant beds in the hospital allowed more than 38 beds to be occupied by the firm's patients on many occasions. This pool was accumulated from vacant beds allocated to all surgical (and sometimes medical) specialties. For the past three years maintaining this pool of vacant beds has depended on a general policy of early discharge. The larger the pool the more effectively fluctuations in bed requirements can be buffered for emergencies, which are unpredictable in terms of both severity of disease and time of the patient's arrival. Emergency cases, however, increased by 19% in three years: What if the increase in emergencies should squeeze out elective surgery and lead to an ever lengthening waiting list? So far this has been avoided at our hospital.

As the waiting list lengthens surgeons are tempted to retain control by keeping patients in "only till tomorrow." This compromises the efficiency of the whole hospital, and the waiting lists grow even more. As an example, to keep in for one extra day all 1127 patients who stayed for five to 15 days in 1985-7 would lengthen the waiting list by perhaps 235 patients over three years. Vacant beds are required to take the varying numbers of emergencies; filling these beds with patients having elective operations would increase the likelihood that a surge of emergencies would cause a bed crisis. The cancellation of booked admissions would follow, with distress to patients and a great deal of extra work for staff.

On a wider scale, where more than one hospital serves a population protection of beds for elective cases in one hospital overloads other hospitals with emergencies and compromises their ability to contain their waiting lists. This has implications for any internal market arrangement in the NHS.

INCREASING THROUGHPUT

We looked to see how beds might be saved, a faster throughput of patients achieved, and the day of breakdown deferred. The throughput of patients that has already been achieved suggests that the scope for improvement is limited. With 38 available beds (or with the mean of 34.5 beds used) the throughput was 48.7 (or 53.7) patients per bed each year. Table II shows the length of stay of patients admitted for emergency and elective operations in 1985-7. The trends for the individual years were similar. Many patients admitted as an emergency (962; 41.2%) and most admitted electively (2100; 65.2%) stayed for two days or less. Day cases accounted for half of the elective

FIG 1—Numbers of beds occupied each day for one year starting 1 August 1986. Each day case required 0.5 bed day. Surgical service was minimised over Christmas (week 22)

TABLE II—Distribution of length of hospital stay of surgical patients

No of days	Emergency admissions (n=2336)	Elective admissions (n=3220)
0	135	1036
1	482	343
2	345	721
3	288	252
4	223	136
5	117	87
6	98	68
7	87	69
8	64	51
9	54	59
10	40	31
11	46	40
12	45	26
13	27	20
14	31	30
15	28	15
16	20	16
17	19	14
18	17	18
19	10	9
20	11	23
21	8	16
22	12	7
23	2	9
24	6	3
25	9	7
26	9	5
27	4	10
28	7	7
29	6	6
30	6	3
>30	80	83
Mean (days)	7.0	4.8

patients. Nevertheless, 80 (3.4%) patients admitted as an emergency and 83 (2.6%) elective patients stayed in hospital for more than 30 days.

What could be done to shorten stays without endangering patients? Would it, for example, be worth while always to admit patients having elective operations on the day of operation, so reducing their stay in hospital by one day? There were 1676 patients who came for elective operations and stayed one to seven days; thus 1676 bed days might have been saved over three years (1.5 empty beds a day), which would allow for perhaps 100 additional cases a year.

Table III shows for 1987 the length of patients' stay in hospital before and after operation for seven groups of common diseases that accounted for one third of all admissions. Most patients were admitted on the day before operation. Admitting more patients on the day

of operation would require clerking the patients before admission and would put extra pressure on the nursing staff who prepare the patients for operation. Patients would not necessarily like such an admission policy, and if they had not fasted for the required time anaesthesia could be risky. On the other hand, a shorter stay might reduce the chance of infection and deep venous thrombosis. Admission several days before operation will always be needed for investigations for vascular, malignant, and pancreatic or biliary disease.

Length of stay after operation varied considerably among the different disease groups: 92% of patients recovered from varicose vein operations in one or two days, and 71% had recovered four days after operation for acute appendicitis. Length of stay was more variable in the other disease groups, and over half the patients with pancreatic or biliary disease stayed in hospital for more than a week after operation. As we had an explicit policy of early discharge these differences reflected differences in the pace of recovery after different operations.

At first sight there seems to be another source of potentially empty beds within the existing system: tabulations of the length of stay (table II) seem to die away to nothing after one month. In a sense, this was true: of 5556 patients admitted, only 163 (2.9%) stayed in hospital for more than 30 days. But these 163 patients had a mean stay of 54.6 days (range 31-278) and so consumed 8907 bed days, 28% of the total. We therefore looked critically at the records of these patients to see whether their stays might have been curtailed (table IV). Although the preponderance of vascular disease reflects our particular interest, we suspect that other features—namely, the importance of infections and the intractability of social delays—are widespread problems.

Of these prolonged admissions, 109 (61%) were of patients over 70 years old. Their death rate (12/100) was significantly higher than that of patients of the same age (106/1984) who did not have a prolonged stay ($p<0.01$; χ^2 test with Yates's correction). Peripheral vascular disease was the most common diagnosis in the long stay patients, and in many cases we used percutaneous transluminal angioplasty as the first manoeuvre to shorter stay. Infection was treated vigorously by antibiotics or surgery as appropriate.

TABLE III—Length of stay (days) of patients before and after operation. Figures are numbers (percentages) of patients

	Before operation						After operation										
	0	1	2	3	4	>4	0	1	2	3	4	5	6	7	>7		
Vascular diseases* (n=163)	36 (22)	83 (51)	11 (7)	10 (6)	7 (4)	16 (10)	11 (7)	26 (16)	21 (13)	3 (2)	4 (2)	6 (4)	6 (4)	6 (4)	80 (49)		
Hernia† (n=114)	32 (28)	76 (67)	2 (2)	0	0	4 (4)	14 (12)	31 (27)	16 (14)	22 (19)	8 (7)	8 (7)	5 (4)	1 (1)	9 (8)		
Varicose veins‡ (n=91)	19 (21)	68 (75)	2 (2)	1 (1)	0	1 (1)	7 (8)	66 (73)	10 (11)	3 (3)	2 (2)	0	0	0	3 (3)		
Cancers (n=83)	18 (22)	41 (49)	4 (5)	7 (8)	3 (4)	10 (12)	8 (10)	12 (14)	3 (4)	7 (8)	2 (2)	5 (6)	8 (10)	5 (6)	33 (39)		
Pancreatic-biliary diseases§ (n=80)	6 (8)	26 (33)	5 (6)	10 (13)	7 (9)	26 (33)	0	0	3 (4)	1 (1)	4 (5)	9 (11)	9 (11)	14 (18)	40 (50)		
Upper gastrointestinal diseases (n=77)	38 (49)	21 (27)	4 (5)	5 (6)	2 (3)	7 (9)	13 (17)	7 (9)	6 (8)	5 (6)	3 (4)	3 (4)	3 (4)	12 (16)	25 (32)		
Appendicitis (n=69)	55 (80)	11 (16)	1 (1)	1 (1)	0	1 (1)	0	0	4 (6)	23 (33)	21 (30)	10 (14)	5 (7)	1 (1)	5 (7)		

*Excludes varicose veins and leg ulcers.

†All types.

‡Excludes leg ulcers.

§Excludes cancers.

TABLE IV—Details of patients who stayed in hospital longer than 30 days

Age (years)	No of admissions	No who died	No with peripheral vascular disease	No with life-threatening infections	No affected by social delay*	Mean (range) length of stay (days)	No of bed days
<50	17		3	5	2	50.5 (31-121)	857
50-59	14	1	4	2	2	53.4 (32-138)	747
60-69	32	2	20	6	4	48.9 (31-115)	1565
70-79	53	4	32	4	9	57.3 (31-210)	3036
80-89	44	8	27	11	5	57.5 (31-278)	2528
90-99	3		1			57.3 (48-69)	172
Total	163	15	87	28	22	49.7 (31-278)	8907

*Delays in provision of geriatric beds, part III accommodation, wheelchairs, artificial limbs, and alterations to patients' homes.

Throughput was impaired by delays in the provision of geriatric beds, part III accommodation, wheelchairs, artificial limbs, and alterations to patients' homes. In the three years monitored 1047 bed days (roughly equivalent to one bed) passed while social factors delayed discharge of 22 patients. About half of this inappropriate use of beds was accounted for by just four patients. Many extra patients could have been treated if the blocked beds had been free. It will become even harder to discharge these patients if local authorities economise by cutting their services.

Table IV also shows that when younger people (aged under 50) had a condition—for example, head injury, necrotising fasciitis, Crohn's disease, pancreatitis, or

premature peripheral vascular disease—that necessitated a hospital stay of more than 30 days they took nearly as long as the elderly (aged 70 and over) to recover sufficiently to leave hospital (mean 50.5 days *v* 57.3 days).

Conclusions

If efficiency in a surgical service means a high throughput of patients and a low rate of cancellation we have three suggestions.

(1) A surgical service might be provided with fewer beds if patients for elective operations were always admitted on the day of operation and discharged at the earliest opportunity and if elderly patients with chronic illness were transferred to other care. The potential for such economy is, however, only 10% of the average number of beds occupied (3.5 of 34.5 in our study).

(2) Numbers of beds should not be decreased further: physical removal of beds runs the risk that

fluctuations in the need for beds can be accommodated only by cancelling elective admissions.

(3) If all surgeons in a district were to increase their throughput the overall cost of health care would be increased and there would be a risk of bankrupting the district health authority. A new system of funding should be set up to encourage efficient surgical units with short waiting lists.

We thank the many house surgeons who entered the data—Richard Baker, Mike Catchpole, Sam Chong, Elizabeth Clifford, Iona Cobb, Paula Domizio, Ellie Dow, Claire Etherington, Paul Evers, Emily Finch, Susan Fish, Alison Grant, Andrea Hughes, Roy Love, Gisella Martinez, Chris Mattock, Marek Miller, Leonie Penna, Melanie Robson, Eileen Rosenfelder, Emma Rowley-Conwy, Andrew Sadler, Alison Seidlin, Peter Stone, Richard Sturgess, John Winchester, Chris Yerbury—and Miss J Oyston for secretarial help.

(Accepted 16 August 1988)

Ambulatory pH monitoring of gastro-oesophageal reflux in “morning dipper” asthmatics

Robyn A Nagel, Paul Brown, Warren H Perks, R S E Wilson, Graeme D Kerr

Abstract

A causal relation between gastro-oesophageal reflux and nocturnal asthma has been postulated. Forty four adult asthmatics underwent ambulatory monitoring of their oesophageal pH over 24 hours to find out if there was such a relation. Of these 21 showed significant “morning dipping” in which the peak expiratory flow falls during the night.

Asthmatics with morning dipping had a history of nocturnal wheeze and a higher incidence of reflux symptoms, but measurement of oesophageal pH showed no significant difference in the amount or pattern of reflux when compared with “non-dippers.”

Overall, 15 asthmatics had gastro-oesophageal reflux, and these participated in a randomised, double blind crossover trial of ranitidine versus placebo. No significant difference was found in the peak expiratory flow rates or subjective evaluation of well being of the patients.

Introduction

Gastro-oesophageal reflux has been reported to be more common in asthmatics¹ and has been postulated to cause worsening of asthma, particularly nocturnal wheezing.² The cause of this is unknown but micro-aspiration of the contents of the stomach might be one possible mechanism, and improvement of asthma has been reported after surgical repair of hiatus hernia.³ Another possible mechanism is that the presence of acid in the oesophagus may stimulate vagal nerve endings which can cause reflex bronchoconstriction.⁴

This study was performed to find out if there was a difference in the degree of gastro-oesophageal reflux

TABLE II—Gastro-oesophageal reflux scores as defined by Goodall *et al*⁶

Gastro-oesophageal reflux score	Heartburn	Regurgitation
0	None	None
1	Occasional	Occasional (on straining or posture change)
2	Needing antacids or medical advice	Predictable
3	Constant (interfering with daily activities)	Occurrence of pulmonary aspiration

between asthmatics with “morning dipping,” in which the peak expiratory flow rate falls during the night, and asthmatics in which this does not occur (“non-dippers”). We also gave ranitidine and placebo to asthmatics with gastro-oesophageal reflux to see whether they would benefit from reduction of acidity in the stomach.

Patients and methods

Forty four adult asthmatics attending the respiratory clinic agreed to participate in the study. All patients showed a greater than 25% increase in peak expiratory flow rate or forced expiratory volume in one second after treatment with bronchodilators. No patient had had gastric or oesophageal operations. One patient who had been prescribed ranitidine for gastro-oesophageal reflux stopped taking the drug for the duration of the study. All patients continued their usual medications for asthma. We recorded the weight of each patient as percentage deviation from the population mean,⁵ taking account of age, sex, and height. A patient was considered to have a history of reflux if symptoms occurred at least once every six weeks.

The patients kept a diary for six days and returned to hospital on day 7 for ambulatory monitoring of their oesophageal pH. They recorded type, dose, and timing of drugs as well as their asthma and reflux scores⁶ for day and night (tables I and II). To determine the peak expiratory flow rate of each patient the best of three

TABLE I—Asthma scores as defined by Goodall *et al*⁶

Asthma score	Daytime	Night time
0	No wheeze	No wheeze
1	Wheeze 1-2 hours	Slept well, slight wheeze
2	Wheeze most of the day	Woke 2-3 times wheezing
3	Wheeze all day	Bad night, much wheeze

Royal Shrewsbury Hospital, Shrewsbury, Shropshire
Robyn A Nagel, FRACP, research registrar
Paul Brown, FRCP, consultant physician
Warren H Perks, MRCP, consultant physician
R S E Wilson, FRCP, consultant physician
Graeme D Kerr, FRACP, consultant physician

Correspondence to:
Dr R Nagel, Flat 14
Surbiton Court, Surbiton,
Surrey.