Maintenance treatment with H₂ receptor antagonists in patients with peptic ulcer disease: reduces morbidity in a significant minority of patients

K G Wormsley

The aim of all treatment is as far as possible to restore the full health of people suffering from disease. This aim is best achieved by eliminating the cause of the disease. Unfortunately, in the case of ulcers the cause is not known¹ and therefore ulcer disease cannot be therapeutically "cured." Under these circumstances the objective of treatment is to alter the natural course of the disease to minimise its symptomatic, functional, and morphological sequelae² and, of course, prevent a lethal outcome. In the case of ulcer disease the principal clinical problems result because ulcers tend to recur, often two, three, or more times each year for many years—and perhaps during the whole lifetime of the affected patient.³ Most recurrences are painful, which prevents work and sleep. During the course of a lifetime, however, about one third of patients with an ulcer bleed from it and nearly 10% suffer from perforation of the ulcer; recurrent relapse and re-healing may result in gross duodenal scarring and pyloric stenosis. These complications or the resulting requirement for operative treatment directly cause the death of 1-2% of patients with ulcers.

Treatment must therefore maintain remission after an ulcer has been healed to minimise interference with the patient's life and work and prevent potentially lethal complications. These objectives cannot be achieved by intermittent treatment—that is, by treating each symptomatic relapse as it occurs—as this type of treatment does not influence the natural course of ulcer disease because it depends on a symptomatic relapse to trigger the next course of treatment. As there are no means of predicting when or how the next recurrence will be manifest in any patient intermittent treatment cannot prevent complications and be expected to reduce the mortality from ulcer disease.

Two therapeutic regimens have been used to sustain remission of ulcers: gastric surgery and long term, continuous (maintenance) treatment with drugs. Gastric surgery is invariably accompanied by mortality and postoperative sequelae in some patients, the sequelae giving rise to occasional irreversible and clinically unacceptable disability. There is, therefore, nowadays no place for the surgical treatment of uncomplicated ulcer disease, except for a few (<3%) patients whose ulcers do not heal during medical treatment; patients whose occupation precludes continuous treatment with drugs; and, occasionally,

patients receiving large doses of non-steroidal anti-inflammatory drugs.

Advantages of maintenance treatment

Maintenance treatment with drugs has beneficially altered the natural course of ulcer disease by keeping most ulcers in remission and altering the clinical characteristics of any relapses that do occur.⁴ For example, during six years of continuous treatment with H₂ receptor antagonists such as ranitidine and cimetidine 84% of 388 patients remained free from symptoms and only 16% suffered from symptomatic recurrence. During the six years only 1% of the patients given maintenance treatment who had recurrences presented with haemorrhage compared with 15% of untreated patients. In other words, most recurrences during maintenance treatment were uncomplicated and therefore posed no threat to the wellbeing or life of the affected patients. In comparison, among patients whose ulcers had relapsed when treatment had been discontinued (in a trial of intermittent treatment) more than 80% of recurrences were symptomatic and more than a quarter of the patients presented with haemorrhage from the recurrent ulcer.⁵ Similar findings were reported from general practice in a study comparing maintenance treatment with intermittent healing treatment. It was concluded that "continuous treatment would benefit many patients treated in general practice."⁶

Asymptomatic recurrences that are discovered endoscopically during maintenance treatment may be ignored because the patients come to no harm as long as maintenance treatment is continued.⁷ Symptomatic recurrences during maintenance treatment are easily rehealed by doubling the dose of the drug used in the maintenance regimen. If a patient relapses more than twice during maintenance treatment the higher (healing) dose of the drug should be used in further maintenance treatment as some patients (especially heavy smokers) need the higher dose of the drug to maintain remission.

The treatment of gastric and duodenal ulcers is identical. Although endoscopic monitoring of duodenal ulcers is unnecessary during maintenance treatment,

continued on page 1394

EDITORIAL COMMENT

Here we have two quite obviously conflicting views between lifelong treatment for everybody with a diagnosis of duodenal ulcer on the one hand and long term maintenance treatment for carefully selected groups on the other. As Dr Wormsley would have everybody with a duodenal ulcer receive maintenance treatment he would obviously agree with using such treatment in the special groups of patients (the elderly patients, those taking non-steroidal anti-inflammatory drugs, those with a previous complication from ulcers, and those with frequent relapses of ulcer disease) mentioned by Dr Howden. I would certainly agree that long term use of H₂ receptor

antagonists in these groups of patients seems eminently reasonable, although even then the argument is based more on theory than evidence. Drs Wormsley and Howden diverge substantially on their approach to managing the large number of patients who do not fall into one of these special categories. What do you do with the patient who presents for the first time with a peptic ulcer which is uncomplicated except for its symptoms? According to the evidence presented here, my own view would be not to embark on lifelong treatment but to wait and see whether the ulcer becomes recurrently troublesome, when I would have no difficulty in embarking on maintenance treatment.—PETER RUBIN, *professor of therapeutics, University Hospital, Nottingham NG7 2UH*

Controversies in Therapeutics

K G Wormsley
continued from page 1392

patients with healed gastric ulcers should be endoscopically re-examined every three months during the first year of maintenance treatment as the original healed gastric ulcer may have been malignant, and recurrence, which is usually fairly rapid, may be asymptomatic. Rapidly recurring gastric ulcers should therefore be treated surgically.

The duration of the relapsing tendency is not known, but in many or most patients with ulcer disease recurrences probably occur throughout life. Maintenance treatment must therefore also be life long, as with most other chronic diseases. Compliance with long term maintenance treatment is excellent because patients develop pain when treatment is stopped. The compliance can be reinforced by pointing out that "a tablet a day keeps the surgeon away."

Conclusions

Long term, and even life long, treatment with drugs is not objectionable provided that the treatment

is effective and safe. Maintenance treatment with currently available H₂ receptor antagonists is effective in keeping ulcers in remission and is safe because unwanted effects during maintenance treatment with these drugs have been so rare as to be truly remarkable.⁸

Maintenance treatment with H₂ receptor antagonists is, therefore, the current optimal treatment for patients with ulcer disease.

- 1 Wormsley KG. Aetiology of ulcers. *Baillieres Clin Gastroenterol* 1988;2:555-71.
- 2 Boyd EJS, Wormsley KG. Natural history of duodenal ulcer. *Surveys of Digestive Diseases* 1985;3:230-9.
- 3 Boyd EJS, Wilson JA, Wormsley KG. Recurrent ulcer disease. In: Misiewicz JJ, Wood JR, eds. *Ranitidine: therapeutic advances*. Amsterdam: Excerpta Medica, 1984:14-42.
- 4 Wormsley KG. Long term treatment of duodenal ulcer. *Postgrad Med J* 1988;64(suppl 1):47-53.
- 5 Boyd EJS, Wilson JA, Wormsley KG. Safety of ranitidine maintenance treatment of duodenal ulcer. *Scand J Gastroenterol* 1984;19:394-400.
- 6 Wade AG, Rowley-Jones D. Long term management of duodenal ulcer in general practice: How best to use cimetidine? *Br Med J* 1988;296:971-4.
- 7 Boyd EJS, Wilson JA, Wormsley KG. The natural history of asymptomatic duodenal ulcer recurrences. *Scand J Gastroenterol* 1984;19:808-12.
- 8 Penston J, Wormsley KG. Adverse reactions and interactions with H₂-receptor antagonists. *Medical Toxicology* 1986;1:192-216.

C W Howden
continued from page 1393

Gastric ulcer perforation rates in England and Wales for men and women aged 65-74 and 75 and over.²⁰ Reproduced with permission

complications in elderly patients.^{19,20} H₂ receptor antagonists are often prescribed to patients who are being treated with non-steroidal anti-inflammatory drugs, and they have been recommended routinely and for long term use in patients with a history of peptic ulcer who need to take non-steroidal anti-inflammatory drugs. H₂ receptor antagonists are effective in healing ulcers in patients being treated with such drugs,²¹ although their long term usefulness in maintaining healing has not been evaluated.²²⁻²⁴

Patients with a previous ulcer complication should be offered maintenance treatment as they are at a greater risk of further complications. Despite the lack of evidence that maintenance treatment with H₂ receptor antagonists reduces complications in general its selective use in patients with a more severe ulcer diathesis may be beneficial.

Patients with frequent episodes of relapse are also suitable for maintenance treatment. At the outset of the disease predicting which patients will relapse frequently is not possible, although certain risk factors, particularly cigarette smoking, may be identified.²² Bardhan has found that only 17% of patients will have three or more relapses a year.²⁵ Such patients may be quickly identified if treated intermittently and transferred to maintenance treatment. Currently most authorities agree that universal maintenance treatment after the first attack of duodenal ulcer entails treating many patients unnecessarily.^{5,16,17,25}

Summary

Maintenance treatment reduces the frequency of

ulcer relapse. Its routine use is unnecessary because the condition is benign and most patients do not relapse frequently. Routine maintenance treatment does not reduce the incidence of complications of ulcers overall, although it may do so in specific groups mentioned above. Routine maintenance treatment for all patients with ulcers would be expensive as well as largely unnecessary.

I thank Drs R I Russell and K D Bardhan for their helpful comments, and Miss R Berry for typing the manuscript.

- 1 Jones DB, Howden CW, Burget DW, et al. Acid suppression in duodenal ulcer: a meta-analysis to define optimal dosing with antisecretory drugs. *Gut* 1987;28:1120-7.
- 2 Howden CW, Jones DB, Peace KE, et al. The treatment of gastric ulcer with antisecretory drugs: relationship of pharmacological effect to healing rates. *Dig Dis Sci* 1988;33:619-24.
- 3 Gough KR, Korman MG, Bardhan KD, et al. Ranitidine and cimetidine in prevention of duodenal ulcer relapse. *Lancet* 1984;ii:659-62.
- 4 Boyd EJS, Wormsley KG. Natural history of duodenal ulcer. *Surveys of Digestive Diseases* 1985;3:230-9.
- 5 Bardhan KD. Intermittent treatment of duodenal ulcer with cimetidine. *Br Med J* 1980;281:20-2.
- 6 Fry J. Peptic ulcer: a profile. *Br Med J* 1964;iii:809-12.
- 7 Elashoff JD, DeVenter GV, Reedy TJ, et al. Long-term follow-up of duodenal ulcer patients. *J Clin Gastroenterol* 1983;5:509-15.
- 8 McManus J. No cure at CURE. *Gastroenterology* 1984;87:731-2.
- 9 Goldberg MA. Medical treatment of peptic ulcer: Is it truly efficacious? *Am J Med* 1984;77:589-91.
- 10 Lee S, Iida M, Yao T, et al. Long-term follow-up of 2529 patients with gastric and duodenal ulcer: survival rate and causes of death. *Gastroenterology* 1988;94:381-6.
- 11 Bonnevie O. Causes of death in duodenal and gastric ulcer. *Gastroenterology* 1977;73:1000-4.
- 12 Bonnevie O. Survival in peptic ulcer. *Gastroenterology* 1978;75:1055-60.
- 13 Schiller LR, Fordtran JS. Ulcer complications during short-term therapy of duodenal ulcer with active agents and placebo. *Gastroenterology* 1986;90:478-81.
- 14 Sonnenberg A. Comparison of different strategies for treatment of duodenal ulcer. *Br Med J* 1985;290:1185-7.
- 15 Jones DB, Howden CW, Hunt RH. Comparison of different strategies for treating duodenal ulcer. *Br Med J* 1985;290:1904.
- 16 Wade AG, Rowley-Jones D. Long-term management of duodenal ulcer in general practice: How best to use cimetidine? *Br Med J* 1988;296:971-4.
- 17 Misiewicz JJ. Maintenance treatment of duodenal ulcer. In: *Peptic ulcer disease. Proceedings of the seventh BSG/SK&F International Workshop*, 1986. Smith, Kline and French Laboratories, 1986:69-72.
- 18 Penston JG, Wormsley KG. Long-term treatment of duodenal ulcers. *Gastroenterology* 1988;94:A349.
- 19 Somerville KW, Faulkner G, Langman M. Non-steroidal anti-inflammatory drugs and bleeding peptic ulcer. *Lancet* 1986;i:462-4.
- 20 Walt RP, Katschinski B, Logan RFA, et al. Rising frequency of ulcer perforation in elderly people in the United Kingdom. *Lancet* 1986;i:489-92.
- 21 Manniche C, Malchow-Moller A, Andersen JR, et al. Randomised study of the influence of non-steroidal anti-inflammatory drugs on the treatment of peptic ulcer in patients with rheumatic disease. *Gut* 1987;28:226-9.
- 22 Sontag S, Graham DY, Belsito A, et al. Cimetidine, cigarette smoking and recurrence of duodenal ulcer. *N Engl J Med* 1984;311:689-93.
- 23 Doherty M. Non-steroidal anti-inflammatory drugs in patients with peptic ulcer disease: to be considered in certain situations. *Br Med J* (in press).
- 24 Hawkey C. Non-steroidal anti-inflammatory drugs in patients with peptic ulcer disease: dangerous in some patient groups. *Br Med J* (in press).
- 25 Bardhan KD. Intermittent treatment of duodenal ulcer for long term medical management. *Postgrad Med J* 1988;64(suppl 1):40-6.